

## **MILK OF MAGNESIA- magnesium hydroxide suspension**

**CVS Pharmacy**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **CVS Pharmacy, Inc. Milk of Magnesia Drug Facts**

#### **Active ingredient (in each 15 mL)**

Magnesium hydroxide 1200 mg

#### **Purpose**

Saline laxative

#### **Uses**

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

#### **Warnings**

##### **Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

##### **Ask a doctor or pharmacist before use if you are**

taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

##### **Stop use and ask a doctor if**

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

##### **If pregnant or breast-feeding,**

ask a health professional before use.

##### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

#### **Directions**

- do not exceed the maximum recommended daily dose in a 24 hour period

- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- mL = milliliter

adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

**Other information**

- each 15 mL contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 3 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F)
- close cap tightly after use
- avoid freezing

**Inactive ingredients**

anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

**Questions or comments?**

1-800-719-9260

**Principal Display Panel**

Compare to the active ingredient in Phillips'® Milk of Magnesia

Milk of Magnesia

MAGNESIUM HYDROXIDE 1200 mg per 15 mL

Saline Laxative

STIMULANT FREE

Comfortable, cramp free relief

Cherry Flavor

ALCOHOL 0.4%

26 FL OZ (1 pt 10 OZ) 769 mL



Compare to the active ingredient in Phillips® Milk of Magnesia\*

NDC 59779-949-35

# Milk of Magnesia

**MAGNESIUM HYDROXIDE**  
1200 mg per 15 mL  
Saline Laxative

**STIMULANT FREE**

- Comfortable, cramp free relief



**Cherry Flavor**  
ALCOHOL 0.4%

26 FL OZ (1 pt 10 OZ) 769 mL

: 94935 17 F4

## Drug Facts

**DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.**

### Active ingredient (in each 15 mL)

Magnesium hydroxide 1200 mg.....Saline laxative

- Uses**
- relieves occasional constipation (irregularity)
  - generally produces bowel movement in ½ to 6 hours

### Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug.

Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

### Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### Directions

- do not exceed the maximum recommended daily dose in a 24 hour period
- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- mL = milliliter

adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

### Other information

- each 15 mL contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 3 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F) ■ close cap tightly after use ■ avoid freezing

**Inactive ingredients** anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

**Questions or comments? 1-800-719-9260**

\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Phillips® Milk of Magnesia.

hB L1 5E6h6 :

Distributed by: **CVS Pharmacy, Inc.**  
One CVS Drive, Woonsocket, RI 02895  
© 2016 CVS/pharmacy  
CVS.com® 1-800-SHOP CVS  
V-16467



#690036



0 50428 34611 2

## MILK OF MAGNESIA

magnesium hydroxide suspension

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:59779-949
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	1200 mg in 15 mL

### Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	

GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0K00R)	
SORBITOL (UNII: 506T60A25R)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	

### Product Characteristics

<b>Color</b>	PINK (Opaque Light pink)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY (wild cherry)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59779-949-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:59779-949-35	769 mL in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	05/05/2005	

**Labeler** - CVS Pharmacy (062312574)