

**ALKA-SELTZER- aspirin, citric acid, sodium bicarbonate effervescent tablet tablet, effervescent
Lil' Drug Store Products, Inc.**

Alka-Seltzer ®

Pouch Header

Alka-Seltzer ®

Anhydrous citric acid / Antacid

Aspirin (**NSAID**) / Analgesic

Sodium bicarbonate / Antacid

[Bayer logo]

ORIGINAL Fast Relief

Do not use if pouch is torn or open

2 EFFERVESCENT TABLETS

Active ingredients (in each tablet)	Purpose
Anhydrous citric acid 1000 mg	Antacid
Aspirin 325 mg (NSAID)*	Analgesic
Sodium bicarbonate (heat-treated) 1916 mg	Antacid

*nonsteroidal anti-inflammatory drug

Uses

for the temporary relief of:

- heartburn, acid indigestion, and sour stomach when accompanied with headache or body aches and pains
- upset stomach with headache from overindulgence in food or drink
- headache, body aches, and pain alone

Warnings

Reye's syndrome

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older • have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic • you have asthma
- you have a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.
- taking a prescription drug for diabetes, gout, or arthritis

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
- feel faint • vomit blood • have bloody or black stools
- have stomach pain that does not get better
- symptoms get worse or last more than 10 days
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs
- new symptoms occur

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Fully dissolve 2 tablets in 4 ounces of water before taking.

Adults and children 12 years and over: take 2 tablets every 4 hours, or as directed by a doctor. Do not exceed 8 tablets in 24 hours. Adults 60 years and over: take 2 tablets every 4 hours, or as directed by a doctor. Do not exceed 4 tablets in 24 hours. Children under 12 years: consult a doctor.

Other information

Each tablet contains: sodium 567 mg. Store at room temperature. Avoid excessive heat. Alka-Seltzer in water contains principally the antacid sodium citrate and the analgesic sodium acetylsalicylate.

Inactive ingredients

none

Questions or comments? 1-800-986-0369 (Mon - Fri 9AM - 5PM EST) or www.alkaseltzer.com

Bayer HealthCare LLC, P.O. Box 1910, Morristown, NJ 07962-1910

Alka-

Seltzer ®

Anhydrous citric acid / Antacid

Aspirin (NSAID) / Analgesic

Sodium bicarbonate / Antacid

[Bayer logo]

ORIGINAL

15 PACKETS OF

2 EFFERVESCENT TABLETS

www.lildrugstore.com

[UPC]

MADE IN MEXICO

• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed **Do not use** • if you are allergic to aspirin or any other pain reliever/fever reducer • if you have ever had an allergic reaction to this product or any of its ingredients **Ask a doctor before use** if • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma • you have a sodium-restricted diet **Ask a doctor or pharmacist before use** if you are presently taking a prescription drug (antacids may interact with certain prescription drugs), or taking a prescription drug for diabetes, gout, or arthritis. **When using this product** do not exceed recommended dosage. **Stop use and ask a doctor** if • an allergic reaction occurs (seek medical help right away) • you experience any of the following signs of stomach bleeding (feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better) • symptoms get worse or last more than 10 days • redness or swelling is present • ringing in the ears or a loss of hearing occurs • new symptoms occur **If pregnant or breast-feeding**, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.** Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** Fully dissolve 2 tablets in 4 ounces of water before taking. Adults and Children 12 years and over: take 2 tablets every 4 hours, or as directed by a doctor. Do not exceed 8 tablets in 24 hours. Adults 60 years and over: take 2 tablets every 4 hours, or as directed by a doctor. Do not exceed 4 tablets in 24 hours. Children under 12 years: consult a doctor **Other information** Each tablet contains: sodium 567 mg. Store at room temperature. Avoid excessive heat. Alka-Seltzer in water contains principally the antacid sodium citrate and the analgesic sodium acetylsalicylate. **Inactive ingredients** None **Questions or comments?** 1-800-986-0369 (Mon-Fri 9AM - 5PM EST) or www.alkaseltzer.com
Bayer HealthCare LLC, P.O. Box 1910, Morristown, NJ 07962-1910

81441596

81441596/1175-01-01E

Alka-Seltzer

Anhydrous citric acid / Antacid
Aspirin (NSAID) / Analgesic
Sodium bicarbonate / Antacid



ORIGINAL Fast Relief

Do not use if pouch is torn or open.

2 EFFERVESCENT TABLETS

Active ingredients (in each tablet)

	Purpose
Anhydrous citric acid 1000 mg.....	Antacid
Aspirin 325 mg (NSAID)*.....	Analgesic
Sodium bicarbonate (heat-treated) 1916 mg.....	Antacid

*nonsteroidal anti-inflammatory drug

Uses for the relief of • heartburn, acid indigestion, and sour stomach when accompanied with headache or body aches and pains • upset stomach with headache from overindulgence in food or drink • headache, body aches, and pain alone **Warnings** **Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug

READ ALL WARNINGS AND DIRECTIONS BEFORE USE. ▼

X23T70 EXP: APR/22 6

81441618

81441618/1176-01-01E



ALKA-SELTZER

aspirin, citric acid, sodium bicarbonate effervescent tablet tablet, effervescent

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-5106
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Base of

Ingredient Name	Basis of Strength	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) (ANHYDROUS CITRIC ACID - UNII:XF417D3PSL)	ANHYDROUS CITRIC ACID	1000 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
SODIUM BICARBONATE (UNII: 8MDF5V39QO) (BICARBONATE ION - UNII:HN1Z RA3Q20)	SODIUM BICARBONATE	1916 mg

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	25mm
Flavor		Imprint Code	ALKA;SELTZER
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-5106-0	15 in 1 BOX, UNIT-DOSE	01/17/2008	06/01/2026
1		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	01/17/2008	06/01/2026

Labeler - Lil' Drug Store Products, Inc. (093103646)

Revised: 12/2023

Lil' Drug Store Products, Inc.