

**TYLOVET 100 - tylosin phosphate powder**  
**Huvepharma, Inc.**

-----  
**Tylovet 100 Type A Medicated Article**

**Tylovet 100 Front Panel**

For use in Swine and Beef Cattle Feeds Only

**Tylovet® 100**

Tylosin Phosphate

**Type A Medicated Article**

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

**For Use in Swine and Beef Cattle Feeds Only**

**Do Not Feed Undiluted**

**Equivalent to 100 g Tylosin per Pound**

**Swine:**

For the reduction in severity of effects of atrophic rhinitis.

For control of swine dysentery associated with *Brachyspira hyodysenteriae*.

For the treatment and control of swine dysentery associated with *Brachyspira hyodysenteriae* immediately after medicating with **Tylovet Soluble** (tylosin) drinking water.

For control of porcine proliferative enteropathies (PPE, ileitis) associated with *Lawsonia intracellularis*.

For control of porcine proliferative enteropathies (PPE, ileitis) associated with *Lawsonia intracellularis* immediately after medicating with **Tylovet Soluble** (tylosin) drinking water.

**Beef Cattle:**

For reduction of incidence of liver abscesses associated with *Fusobacterium necrophorum* and *Arcanobacterium pyogenes*.

**Important: Must be Thoroughly Mixed in Feeds Before Use. Restricted Drug (California) Use Only as Directed. ANADA# 200-484, Approved by FDA**

**Manufactured By: Biovet AD**  
**Distributed By: Huvepharma, Inc. 39, Petar Rakov Str. 4550**  
**525 Westpark Drive, Suite**

**230 Bulgaria**

**Peachtree, GA 30269**

Huvepharma®, and Tylovet® are trademarks of Huvepharma EOOD.

To report adverse effects, access medical information, or obtain additional product information, call 1-877-426-7765.

**Net weight: 50 lb (22.68 kg)**

**HUVEPHARMA®**

## Directions for Use

### Read all Directions Carefully Before Mixing and Feeding

Tylovet® 100 Type A Medicated Article

Net Weight: 50 lbs (22.68 kgs)

### Do Not Feed Undiluted

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

**Active Drug Ingredient:** Tylosin (as tylosin phosphate)...100 g per lb

**Ingredients:** Roughage products, pregelatinized starch and dipotassium phosphate.

**Caution:** To ensure adequate mixing, an intermediate blending step should be used prior to manufacturing a complete feed. Do not use in any

liquid feed containing sodium metabisulfite or in any finished feed (supplement, concentrate or complete feed) containing in excess of 2%

bentonite.

**Warning: Tylovet 100** may be irritating to unprotected skin and eyes. When mixing and handling **Tylovet 100** use protective clothing, impervious gloves and a dust respirator. In case of accidental exposure, flush eyes with plenty of water. Exposed skin should be washed with plenty of soap and water. Remove and wash contaminated clothing. Seek medical attention if irritation becomes severe or persists. The material safety data sheet (MSDS) contains more detailed occupational safety information. To report adverse effects, access medical information, or obtain additional product information, call 1-877-426-7765.

**For the reduction in severity of effects of atrophic rhinitis.** Feed 100 g of tylosin per ton (1.0 pound **Tylovet 100** per ton) of complete feed. Feed continuously as the sole ration.

**For control of swine dysentery.** Feed 100 g of tylosin per ton (1.0 pound **Tylovet 100** per ton) of complete feed for at least three weeks. Follow with 40 g tylosin per ton (0.4 pounds **Tylovet 100** per ton) of complete feed until pigs reach market weight.

**For the treatment and control of swine dysentery.** Feed 40 to 100 grams of tylosin (0.4 to 1.0 pounds of **Tylovet 100**) per ton of complete feed for 2 to 6 weeks immediately after medicating with 250 mg tylosin (as **Tylosin Soluble**) per gallon of drinking water for 3 to 10 days.

**For the control of porcine proliferative enteropathies. (PPE, ileitis).** Feed 100 g tylosin per ton (1.0 pound **Tylovet 100** per ton) of complete feed for 21 days. Alternatively, feed 100 g of tylosin per tone (1.0 pound **Tylovet 100** per ton) of complete feed for at least three weeks, followed by 40 g tylosin per ton of complete feed until pigs reach market weight. Alternatively, feed 40 to 100 grams of tylosin (0.4 to 1.0 pounds of **Tylovet 100**) per ton of complete feed for 2 to 6 weeks immediately after medicating with 250 mg tylosin (as **Tylovet Soluble**) per gallon in drinking water for 3 to 10 days. Feed continuously as the sole ration when feeding Tylovet.

**NOTICE: Organisms vary in their degree of susceptibility to any chemotherapeutic. If no improvement is observed after recommended treatment, diagnosis and susceptibility should be reconfirmed.**

---

## Mixing and Feeding

### Directions for Beef

#### Cattle Feeds

For reduction of incidence of liver abscesses associated with

*Fusobacterium necrophorum* and *Arcanobacterium pyogenes*

#### Tylovet 100 per Ton of Type C Feed

0.08 to 0.10 lbs

To be fed so that each animal receives not more than 90 mg per head per day and not less than 60 mg per head per day.

Feed continuously as the sole ration.

**Tylosin Per Ton of Type C Feed**  
8 to 10 g

---

---

**Limitations :**

1. pH must be between 4.5 and 6.0
  2. For liquid feeds stored in recirculating tank systems: Recirculate immediately prior to use for not less than 10 minutes, moving not less than 1 percent of the tank contents per minute from the bottom of the tank to the top.
  3. For liquid feeds stored in mechanical, air, or other agitation - type tank systems: Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily as described even when not used.
  4. Liquid Type B Cattle Feeds must bear an expiration date of 31 days after the date of manufacture.
- 

**☐Mixing****Directions for  
Liquid Type B  
Cattle Feeds**

Recirculate daily as described even when not used.

**Store at Room Temperature, 25°C (77°F). Avoid moisture.  
Not to be used after the date printed on the bottom of the bag.**

**Back Panel****Tylovet® 100**

Tylosin Phosphate

**Type A Medicated Article****Tylovet 100 Bag label**

# TYLOVET 100

tylosin phosphate powder

## Product Information

VED TYPE A MEDICATED ARTICLE ANIMAL

Item Code

NDC 22242

**Tylovet® 100**  
Type A Medicated Article

HUVEPHARMA

For use in Swine and Beef Cattle Feeds Only

**Tylovet® 100**  
Tylosin Phosphate

**Type A Medicated Article**

**CAUTION:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

**For Use in Swine and Beef Cattle Feeds Only**

**Do Not Feed Undiluted**

Equivalent to 100 g Tylosin per Pound

**Swine:**  
 An investigational study of efficacy of Tylosin Phosphate for the treatment and control of enteric disease associated with *Archaeobacter* spp. in swine was conducted with Tylovet 100. In this study, Tylovet 100 was administered to swine in the form of a medicated feed. The results of this study are pending.

**Beef Cattle:**  
 An investigational study of efficacy of Tylosin Phosphate for the treatment and control of enteric disease associated with *Archaeobacter* spp. in beef cattle was conducted with Tylovet 100. In this study, Tylovet 100 was administered to beef cattle in the form of a medicated feed. The results of this study are pending.

**Net weight: 50 lb (22.68 kg)**

HUVEPHARMA

**Tylovet® 100**

**Tylovet® 100**  
Type A Medicated Article

HUVEPHARMA

**Tylovet® 100**  
Tylosin Phosphate  
Type A Medicated Article

HUVEPHARMA

**Directions for Use**

Tylovet® 100 Tylosin Phosphate is a medicated feed additive. It is used to control enteric disease associated with *Archaeobacter* spp. in swine and beef cattle. Tylovet 100 is administered to swine and beef cattle in the form of a medicated feed. The results of this study are pending.

**CAUTION:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

**Do Not Feed Undiluted**

Equivalent to 100 g Tylosin per Pound

**Swine:**  
 An investigational study of efficacy of Tylosin Phosphate for the treatment and control of enteric disease associated with *Archaeobacter* spp. in swine was conducted with Tylovet 100. In this study, Tylovet 100 was administered to swine in the form of a medicated feed. The results of this study are pending.

**Beef Cattle:**  
 An investigational study of efficacy of Tylosin Phosphate for the treatment and control of enteric disease associated with *Archaeobacter* spp. in beef cattle was conducted with Tylovet 100. In this study, Tylovet 100 was administered to beef cattle in the form of a medicated feed. The results of this study are pending.

**Net weight: 50 lb (22.68 kg)**

HUVEPHARMA

USDCV4111062260 USM410

<b>Product Type</b>	VFD TYPE A MEDICATED ARTICLE ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:23243-2350	
<b>Route of Administration</b>	ORAL			
<b>Active Ingredient/Active Moiety</b>				
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>	
	TYLOSIN PHOSPHATE (UNII: NR75X12AFF) (TYLOSIN - UNII: YEF4JXN031)	TYLOSIN	100 g in 0.45 kg	
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:23243-2350-1	22.68 kg in 1 BAG		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
ANADA	ANADA200484	03/01/2012		

**Labeler** - Huvepharma, Inc. (619153559)

Revised: 4/2017

Huvepharma, Inc.