

ACTINEL- dextromethorphan hbr, guaifenesin, pseudoephedrine hcl solution
ACTIPHARMA, LLC

ACTINEL®

Drug Facts

Active Ingredients (in each 5 mL tsp)

Dextromethorphan HBr, 15 mg
Guaifenesin, 200 mg
Pseudoephedrine HCl, 30 mg

Purposes

Cough Suppressant
Expectorant
Nasal Decongestant

Uses

• temporarily relieves cough due to minor throat and bronchial irritation associated with the common cold • helps control the cough reflex that causes coughing • helps loosen phlegm (mucus) and thin bronchial secretions to make cough more productive • temporarily relieves nasal congestion due to the common cold, hay fever, or other upper respiratory allergies (allergic rhinitis).

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

• heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to an enlarged prostate gland • cough that occurs with too much phlegm (mucus) • cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

When using this product do not use more than directed.

Stop use and ask a doctor if

• you get nervous, dizzy, or sleepless

- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 4 doses in any 24-hour period.

AGE	DOSE
Adults and children 12 years of age and over	Take 2 teaspoonfuls (10 mL) every 6 hours
Children under 12 years of age	do not use

Other information

- Tamper Evident Feature: Do not use if inner seal is torn, cut, or opened
- Store at controlled room temperature 15°- 30°C (59° - 86°F)
- Avoid excessive heat or humidity.

Inactive ingredients

Citric Acid, Flavor, Methylparaben, Potassium Citrate, Propylene Glycol, Propylparaben, Purified Water, Sorbitol, Sucralose

Questions or comments?

call weekdays from 8AM to 4PM AST at **1.787.608.0882**

NEW FORMULA

Contains the same active ingredients as Tusnel®*

SUGAR FREE

DYE FREE

ALCOHOL FREE

SACCHARIN FREE

COMMITTED TO HEALTH AND WELL-BEING

WWW.ACTIPHARMA.NET

Manufactured in USA with imported ingredients for ActiPharma, Inc.
San Juan, PR 00917.

* Tusnel® is a registered trademark of Llorens Pharmaceutical Corp. This product is not manufactured, distributed or marketed by Llorens Pharmaceutical Corp.

Packaging

NDC 63102-101-16

ACTINEL®

NEW FORMULA

Contains the same active ingredients as Tusnel®*

**COUGH SUPPRESSANT
EXPECTORANT
NASAL DECONGESTANT**

SUGAR FREE
DYE FREE
ALCOHOL FREE
SACCHARIN FREE



16 Fl.oz. (473 mL)

Drug Facts

Active Ingredients (in each 5 mL tsp) **Purposes**
Dextromethorphan HBr, 15 mg..... Cough Suppressant
Guaifenesin, 200 mg..... Expectorant
Pseudoephedrine HCl, 30 mg..... Nasal Decongestant

Uses
temporarily relieves cough due to minor throat and bronchial irritation associated with the common cold ■ helps control the cough reflex that causes coughing ■ helps loosen phlegm (mucus) and thin bronchial secretions to make cough more productive ■ temporarily relieves nasal congestion due to the common cold, hay fever, or other upper respiratory allergies (allergic rhinitis).

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
■ heart disease ■ high blood pressure ■ thyroid disease
■ diabetes ■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus)
■ cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

When using this product do not use more than directed.

Stop use and ask a doctor if
■ you get nervous, dizzy, or sleepless
■ symptoms do not get better within 7 days or are accompanied by fever

Drug Facts (continued)

■ cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not take more than 4 doses in any 24-hour period.

AGE	DOSE
Adults and children 12 years of age and over	Take 2 teaspoonfuls (10 mL) every 6 hours
Children under 12 years of age	do not use

Other information
■ Tamper Evident Feature: Do not use if inner seal is torn, cut, or opened
■ Store at controlled room temperature 15° - 30°C (59° - 86°F)
■ Avoid excessive heat or humidity.

Inactive ingredients Citric Acid, Flavor, Methylparaben, Potassium Citrate, Propylene Glycol, Propylparaben, Purified Water, Sorbitol, Sucralose

Questions or comments?
call weekdays from 8AM to 4PM AST at **1.787.608.0882**

Manufactured in USA with imported ingredients for ActiPharma, Inc. San Juan, PR 00917. Rev 9/23

* Tusnel® is a registered trademark of Llorens Pharmaceutical Corp. This product is not manufactured, distributed or marketed by Llorens Pharmaceutical Corp.



ACTINEL

dextromethorphan hbr, guaifenesin, pseudoephedrine hcl solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63102-101
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg in 5 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 5 mL
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	

METHYLPARABEN (UNII: A2I8C7HI9T)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63102-101-16	474 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/24/2014	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/24/2014	

Labeler - ACTIPHARMA, LLC (079340948)

Revised: 6/2024

ACTIPHARMA, LLC