# MEIJER MAXIMUM STRENGTH ORAL PAIN RELIEF- benzocaine, and benzalkonium chloride gel

MEIJER DISTRIBUTION INC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

### Active ingredients Purpose

Benzocaine 20%.....Oral pain reliever

Benzalkonium chloride 0.1%......Antiseptic

#### Use

- for the temporary relief of pain due to toothaches
- to help protect against infection in minor oral irritation

#### **Warnings**

**Allergy alert:** do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics

#### Do not use

- more than directed
- for more than 7 days unless directed by a physician or healthcare provider

### Stop use and ask a physician if

- swelling, rash or fever develops
- irritation, pain or redness persists or worsens
- symptoms do not improve in 7 days
- allergic reaction occurs

#### Keep out of reach of children.

In case of overdose or allergic reaction, get medical help or contact a Poison Control Center right away.

#### **Directions**

• cut open tip of tube on score mark

Adults and children 2 years of age and over	Apply a small amount of product to the cavity around gum surrounding the teeth.
Children under 12 years of age	Should be supervised in the use of this product
Children under 2 years of age	Ask a physician or healthcare provider

- do not use if tube tip is cut prior to opening
- this preparation is intended for use in cases of toothache, only as a temporary expedient until a physician can be consulted
- do not use continuously
- avoid using toothpaste or drinking soft drinks or fruit juices for at least one hour after applying


**Inactive ingredients** glycerin, peppermint oil, polyethylene glycol 400, polyethylene glycol 4000, saccharin sodium, sorbitol

DIST. BY MEIJER DISTRIBUTION, INC.

**GRAND RAPIDS, MI 49544** 

www.meijer.com

Made in Korea



# ralPainReliefGel

Benzocaine 20% • Benzalkonium Chloride 0.1%

**Double Medicated for Fast Relief** 

- Benzocaine Works To Relieve Oral Pain
- Benzalkonium Chloride Helps Protect Against Oral Infection

## meijer

**Maximum Strength** 

Benzocaine 20% • Benzalkonium Chloride 0.1%

NET WT 0.42 OZ (11.9 g)

**Double Medicated for Fast Relief** 







DIST. BY MEIJER DISTRIBUTION, INC. GRAND RAPIDS, MI 49544 www.meijer.com

Made in Korea

\*This product is not manufactured or distributed by Church & Dwight, Inc., owner of the registered trademark Orajel®.

Lot & Exp.

<b>nestions or comments?</b> call 1-800-814-8028			
Inactive ingredients glycerin, peppermint oil, polyethylene glycol 400, polyethylene glycol 4000, saccharin sodium, sorbitol			
<b>Other information</b> = do not use if tube tip is cut prior to opening = this preparation is intended for use in cases of toothache, only as a femporary expedient until a physician can be consulted = do not use continuously = avoid using toothpaste or drinking soft drinks or fruit juices for at least one hour after applying			
ildren under 2 years of age Ask a physician or healthcare provider			
illdren under 12 years of age Should be supervised in the use of this product			
ults and children 2 years Apply a small amount of product to the cavity and around gum surrounding the teeth.  Use up to 4 times daily or as directed by a physician or healthcare provider			
irections ■ cut open tip of tube on score mark			
e <b>p out of reach of children.</b> ep out <b>of reach of children.</b> case of overdose or allergic reaction, get medical help or contact a Polson Control Center right away.			
Stop use and ask a physician if ■ swelling, rash or fever develops ■ lirritation, pain or redness persists or worsens ■ slymptoms do not improve in 7 days ■ sllergic reaction occurs			
Do not use ■ more than directed ■ for more than 7 days unless directed by a physician or healthcare provider			
<b>/armings</b> l <b>ergy alert:</b> do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, nzocaine, or other "caine" anesthetics			
S • for the temporary relief of pain due to toothaches • to help protect against infection in minor oral irritation			
Virye ingredients cial particular cial particular cial particular cial particular cial particular cial particular cial cial cial cial cial cial cial cial			

#### MEIJER MAXIMUM STRENGTH ORAL PAIN RELIEF

benzocaine, and benzalkonium chloride gel

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-782
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
BENZO CAINE (UNII: U3RS Y48 JW5) (BENZO CAINE - UNII: U3RS Y48 JW5)	BENZOCAINE	200 mg in 1 g	
<b>BENZALKO NIUM CHLO RIDE</b> (UNII: F5UM2KM3W7) (BENZALKO NIUM - UNII:7N6 JUD5X6 Y)	BENZALKONIUM CHLORIDE	1 mg in 1 g	

Inactive Ingredients		
Ingredient Name	Strength	
GLYCERIN (UNII: PDC6A3C0OX)		
PEPPERMINT OIL (UNII: AV092KU4JH)		
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)		
POLYETHYLENE GLYCOL 4000 (UNII: 4R4HFi6 D95)		
SACCHARIN SO DIUM (UNII: SB8 ZUX40 TY)		
SORBITOL (UNII: 506T60A25R)		

F	Packaging			
#	Item Code	Package Description	<b>Marketing Start Date</b>	Marketing End Date
1	NDC:41250-782-42	1 in 1 BOX	12/08/2016	
1		11.9 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	12/08/2016	

### Labeler - MEJER DISTRIBUTION INC (006959555)

Revised: 12/2016 MEIJER DISTRIBUTION INC