

STOOL SOFTENER LAXATIVE PLUS STIMULANT- docusate sodium and sennosides tablet
Walgreens

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each tablet)

Docusate sodium 50 mg

Sennosides 8.6 mg

Purpose

Stool softener

Stimulant laxative

Uses

- for overnight relief from occasional constipation (irregularity)
- generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

- laxative products for longer than 1 week unless told to do so by a doctor
- if you are presently taking mineral oil, unless told to do so by a doctor

Ask a doctor before use if you have

- stomach pain
- nausea
- vomiting
- noticed a sudden change in bowel habits that lasts over a period of 2 weeks

Stop use and ask a doctor if

you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- Take only by mouth. Doses may be taken as a single daily dose, preferably in the evening, or in divided doses.

adults and children 12 years and older	take 2-4 tablets daily
children 6 to under 12 years of age	take 1-2 tablets daily
children 2 to under 6 years of age	take up to 1 tablet daily
children under 2	ask a doctor

Other information

- **each tablet contains:** calcium 30 mg
- **each tablet contains:** sodium 6 mg VERY LOW SODIUM
- store at 25°C (77°F); excursions permitted between 15-30°C (59-86°F)

Inactive ingredients

carnauba wax*, croscarmellose sodium, dibasic calcium phosphate dihydrate, FD&C blue #2 aluminum lake, FD&C red #40 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol*, silicon dioxide, sodium benzoate*, stearic acid, talc*, titanium dioxide

*contains one or more of these ingredients

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredients in Colace® 2-IN-1††

Stool Softener + Stimulant Laxative

DOCUSATE SODIUM 50 mg / STOOL SOFTENER

SENNOSIDES 8.6 mg / STIMULANT LAXATIVE

GENTLE DEPENDABLE

- Relief in 6 to 12 hours

TABLETS

††This product is not manufactured or distributed by AVrio Health L.P., distributor of

Colace® 2-IN1

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS
BROKEN OR MISSING**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT
INFORMATION**

DISTRIBUTED BY: WALGREENS CO,
200 WILMOT RD., DEERFIELD, IL 60015
walgreens.com

Product Label

Walgreens

Compare to the active ingredients in Colace® 2-IN-1[†]

NDC 0363-0827-60

Stool Softener + Stimulant Laxative

DOCUSATE SODIUM 50 mg /
STOOL SOFTENER
SENNOSIDES 8.6 mg /
STIMULANT LAXATIVE

GENTLE DEPENDABLE

• Relief in 6 to 12 hours

60
TABLETS



ACTUAL SIZE

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2018 Walgreen Co.

ITEM 484739 W10024-1018-L



PLD-F9U
FC005448

Lot No.:

Exp. Date:

Drug Facts

Active ingredients (in each tablet)

Docusate sodium 50 mg.....Stool softener
Sennosides 8.6 mg.....Stimulant laxative

Purpose

Uses

- for overnight relief from occasional constipation (irregularity)
- generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

- laxative products for longer than 1 week unless directed by a doctor
- if you are now taking mineral oil, unless directed by a doctor

Ask a doctor before use if you have

- stomach pain ■ nausea ■ vomiting
- noticed a sudden change in bowel habits that lasts over a period of 2 weeks

Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take only by mouth. Doses may be taken as a single daily dose, preferably in the evening, or in divided doses.

Drug Facts (continued)

adults and children 12 years and older	take 2-4 tablets daily
children 6 to under 12 years of age	take 1-2 tablets daily
children 2 to under 6 years of age	take up to 1 tablet daily
children under 2	ask a doctor

Other information

- each tablet contains: calcium 30 mg
- each tablet contains: sodium 6 mg VERY LOW SODIUM
- store at 25°C (77°F); excursions permitted between 15-30°C (59-86°F)

Inactive ingredients carnauba wax*, croscarmellose sodium, dibasic calcium phosphate dihydrate, FD&C blue #2 aluminum lake*, FD&C red #40 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium benzoate*, stearic acid*, talc*, titanium dioxide

*contains one or more of these ingredients

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

[†]Walgreens Pharmacist Survey

^{††}This product is not manufactured or distributed by Avrio Health L.P., distributor of Colace® 2-IN-1.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

WALGREENS Stool Softener plus Stimulant Laxative

STOOL SOFTENER LAXATIVE PLUS STIMULANT

docusate sodium and sennosides tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0827
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

DOCUSATE SODIUM (UNII: F05Q2T2JA0) (DOCUSATE - UNII:M7P27195AG)		DOCUSATE SODIUM	50 mg	
SENNOSIDES (UNII: 3FYP5M0IJX) (SENNOSIDES - UNII:3FYP5M0IJX)		SENNOSIDES	8.6 mg	
Inactive Ingredients				
Ingredient Name			Strength	
CARNAUBA WAX (UNII: R12CBM0EIZ)				
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)				
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
HYPROMELLOSES (UNII: 3NXW29V3WO)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
TALC (UNII: 7SEV7J4R1U)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
ALUMINUM OXIDE (UNII: LMI26O6933)				
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)				
Product Characteristics				
Color	red	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	TCL097;PSD21;S44	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0827-60	1 in 1 BOX	09/30/2018	
1		60 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0363-0827-20	200 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/30/2018	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC monograph not final	part334		09/30/2018	

Labeler - Walgreens (008965063)

