

MOTRIN ARTHRITIS PAIN- diclofenac sodium gel
Johnson & Johnson Consumer Inc.

MOTRIN ARTHRITIS PAIN

Drug Facts

Active ingredient

Diclofenac sodium (NSAID*) 1% (equivalent to 0.93% diclofenac)

*nonsteroidal anti-inflammatory drug

Purpose

Arthritis pain reliever

Uses

- for the temporary relief of arthritis pain ONLY in the following areas:
- hand, wrist, elbow (upper body areas)
- foot, ankle, knee (lower body areas)
- this product may take up to 7 days to work for arthritis pain; it is not for immediate relief. If no pain relief in 7 days, stop use.

Warnings

For external use only

Allergy alert

Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- blisters
- facial swelling
- shock
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Liver warning

This product contains diclofenac. Liver damage may occur if you apply

- more or for a longer time than directed
- when using other drugs containing diclofenac

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is small but higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- apply more or for longer than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever or to a fever reducer
- for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.
- right before or after heart surgery
- on more than 2 body areas at the same time
- in the eyes, nose or mouth

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you are under age 18 years. It is not known if this drug works or is safe in children under age 18 years.

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- avoid contact with eyes, nose, or mouth
- if eye contact occurs, rinse thoroughly with water

Stop use and ask a doctor if

- pain gets worse or lasts more than 21 days
- redness or swelling is present in the painful area
- fever occurs

- skin irritation occurs
- any new symptoms appear. These could be signs of a serious condition.
- you experience any of the following signs of stomach bleeding:
- feel faint
- have bloody or black stools
- vomit blood
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- leg swelling
- weakness in one part or side of body
- slurred speech

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use diclofenac at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

Use up to 21 days unless directed by your doctor

Not for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.

Daily	Per Dose
<p>For your arthritis pain:</p> <ul style="list-style-type: none"> • Use 4 times per day every day • Do not use on more than 2 body areas at the same time 	<p>Use ENCLOSED DOSING CARD to measure a dose</p> <ul style="list-style-type: none"> • For each upper body area (hand, wrist, or elbow) – Squeeze out 2.25 inches (2 grams) • For each lower body area (foot, ankle or knee) – Squeeze out 4.5 inches (4 grams)

Read the enclosed User Guide for complete instructions:

- use only as directed
- do not use more than directed or for longer than directed
- apply only to clean, dry skin that does not have any cuts, open wounds, infections or rashes
- do not apply in same area as any other product
- do not apply with external heat such as heating pad
- do not apply a bandage over the treated area
- store ENCLOSED DOSING CARD with your Motrin® Arthritis Pain product. The dosing card is re-usable.

Other information

- store at 20-25°C (68-77°F). Keep from freezing.
- read all product information before using. Keep the dosing card, this carton and accompanying User Guide for important information.

Inactive ingredients

carbomer homopolymer Type C, cocoyl caprylocaprate, isopropyl alcohol, mineral oil, polyoxyl 20 cetostearyl ether, propylene glycol, purified water, strong ammonia solution

Questions or comments?

Call **1-800-962-5357** (toll-free) or **215-273-8755** (collect) or visit www.motrin.com

PRINCIPAL DISPLAY PANEL

NDC 50580-574-02

Original

Prescription

Strength

New!

Different Ingredient

Diclofenac Sodium

Motrin®

ARTHRITIS PAIN

Diclofenac Sodium

Topical Gel, 1% (NSAID)

ARTHRITIS PAIN RELIEVER

Anti-inflammatory

For daily treatment of
arthritis pain

Fragrance

Free**

For external use only

NET WT. 3.53 OZ (100 g)

Code area

Drug Facts (continued)
Directions
 Use up to 21 days unless directed by your doctor. Not for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.

Daily	Per Dose
For your arthritis pain: ■ Use 4 times per day every day ■ Do not use on more than 2 body areas at the same time	Use ENCLOSED DOSING CARD to measure a dose ■ For each upper body area (hand, wrist, or elbow) – Squeeze out 2.25 inches (2 grams) ■ For each lower body area (foot, ankle or knee) – Squeeze out 4.5 inches (4 grams)

Read the enclosed **User Guide** for complete instructions:
 ■ use only as directed ■ do not use more than directed or for longer than directed

Drug Facts (continued)
 ■ apply only to clean, dry skin that does not have any cuts, open wounds, infections or rashes
 ■ do not apply in same area as any other product ■ do not apply with external heat such as heating pad
 ■ do not apply a bandage over the treated area ■ store ENCLOSED DOSING CARD with your Motrin® Arthritis Pain product. The dosing card is re-usable.

Other information
 ■ store at 20-25°C (68-77°F). Keep from freezing.
 ■ read all product information before using. Keep the dosing card, this carton and accompanying User Guide for important information.

Inactive ingredients carbomer homopolymer Type C, cetyl caprylocaprate, isopropyl alcohol, mineral oil, polyoxy 20 ceteary ether, propylene glycol, purified water, strong ammonia solution

Questions or comments? Call 1-800-962-5387 (toll-free) or 215-273-8735 (collect) or visit www.motrin.com

NDC 50580-574-02

Original Prescription Strength

Motrin®

ARTHROSIS PAIN

Diclofenac Sodium Topical Gel, 1% (NSAID)

ARTRHOSIS PAIN RELIEVER

Anti-inflammatory
For daily treatment of arthritis pain

NEW!
Different Ingredient
Diclofenac Sodium

Fragrance Free**

For external use only

NET WT. 3.53 OZ (100 g)

Use Enclosed Dosing Card to Measure a Dose

Motrin ARTHROSIS PAIN Diclofenac Sodium Topical Gel, 1% (NSAID) Arthritis Pain Reliever

Dosing Card (Not Actual Size)
Read product User Guide before use
Not for immediate pain relief

Upper Body Dose 2.25 inches long (2 grams)
Lower Body Dose 4.5 inches long (4 grams)

Treatment Areas

Hand Wrist Elbow
Foot Ankle Knee

How to Use
 ● For arthritis pain only ● Use 4 times a day ● For use on no more than 2 body areas
 ● May take up to 7 days to work for your arthritis pain ● Use up to 21 days unless directed by your doctor

Not for use on any other body area (Such as back, hip or shoulder)
Medicated Gel Clinically Proven to Relieve Arthritis Pain

**No Fragrance Added to Product

NDC 50580-574-02

Original Prescription Strength

Motrin®

ARTHROSIS PAIN

Diclofenac Sodium Topical Gel, 1% (NSAID)

ARTRHOSIS PAIN RELIEVER

Anti-inflammatory
For daily treatment of arthritis pain

NEW!
Different Ingredient
Diclofenac Sodium

Fragrance Free**

For external use only

NET WT. 3.53 OZ (100 g)

Drug Facts
Active ingredient
 Diclofenac sodium (NSAID)* 1% (equivalent to 0.93% diclofenac)..... Arthritis pain reliever
 *nonsteroidal anti-inflammatory drug

Purpose
 Arthritis pain reliever

Uses
 ■ for the temporary relief of arthritis pain ONLY in the following areas:
 ■ hand, wrist, elbow (upper body areas) ■ foot, ankle, knee (lower body areas)
 ■ this product may take up to 7 days to work for arthritis pain; it is not for immediate relief.
 If no pain relief in 7 days, stop use.

Warnings
For external use only
Allergy alert: Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 ■ hives ■ asthma (wheezing) ■ skin reddening ■ blisters ■ facial swelling ■ shock ■ rash
 If an allergic reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)
Liver warning: This product contains diclofenac. Liver damage may occur if you apply more or for a longer time than directed. ■ when using other drugs containing diclofenac.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is small but higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product.
 ■ apply more or for longer than directed.
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Do not use ■ if you have ever had an allergic reaction to any other pain reliever or to a fever reducer ■ for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.

LIFT HERE

Lift Flap for Important Directions & Warnings

(42H) EFB 2547MV

Drug Facts (continued)
 ■ right before or after heart surgery ■ on more than 2 body areas at the same time
 ■ in the eyes, nose or mouth

Ask a doctor before use if
 ■ you have problems or serious side effects from taking pain relievers or fever reducers
 ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn
 ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 ■ you are taking a diuretic
 ■ you are under age 18 years. It is not known if this drug works or is safe in children under age 18 years.

Ask a doctor or pharmacist before use if you are
 ■ under a doctor's care for any serious condition ■ taking any other drug

When using this product
 ■ avoid contact with eyes, nose, or mouth ■ if eye contact occurs, rinse thoroughly with water

Drug Facts (continued)
Stop use and ask a doctor if ■ pain gets worse or lasts more than 21 days
 ■ redness or swelling is present in the painful area ■ fever occurs ■ skin irritation occurs
 ■ any new symptoms appear. These could be signs of a serious condition.
 ■ you experience any of the following signs of stomach bleeding:
 ■ feel faint ■ have bloody or black stools ■ vomit blood ■ have stomach pain that does not get better
 ■ you have symptoms of heart problems or stroke:
 ■ chest pain ■ trouble breathing
 ■ leg swelling ■ weakness in one part or side of body ■ slurred speech

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use diclofenac at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

USER GUIDE

Motrin® & NDC# = PMS 2765 overprinting background
 All dark violet PDP & side panel text = PMS 2617 overprinting background

Motrin[®] ARTHRITIS PAIN

Diclofenac Sodium Topical Gel, 1% (NSAID) Arthritis Pain Reliever

Use Guide

How to use Motrin[®] Arthritis Pain Gel with answers to frequently asked questions

Quick Start

How to get started right away with Motrin[®] Arthritis Pain Gel
 Understanding Motrin[®] Arthritis Pain Gel
 Motrin[®] Arthritis Pain (Diclofenac Sodium Topical Gel, 1% (NSAID) —arthritis pain reliever) is an arthritis pain relief medication that you apply to your skin.

Before You Use Motrin[®] Arthritis Pain Gel
 Read the Drug Facts Label, which appears on the carton.



Remove the dosing card from the inside of the carton. You should always use the dosing card to measure out the correct dose of Motrin[®] Arthritis Pain Gel.

The first time you use a tube of Motrin[®] Arthritis Pain Gel, you will have either one of the tube caps shown below.



1. Take the cap off the tube.




2. Open the safety seal by firmly pressing the point on the top of the cap onto the seal on the tube. Do not open the safety seal with scissors or other sharp objects.



3. After use, put the cap back on the end of the tube.


Using Motrin[®] Arthritis Pain Gel
 How often to use Motrin[®] Arthritis Pain Gel
 Use 4 times a day for best results.

For example:




Where to apply Motrin[®] Arthritis Pain Gel
 Motrin[®] Arthritis Pain Gel can be used on up to 2 body areas from the following list: a hand, a wrist, an elbow, a foot, an ankle, a knee. Do not use on more than 2 body areas at once.

Some examples include:



Measuring the correct amount using the dosing card




For each upper body area (a hand, a wrist, or an elbow): Squeeze gel from the tube equal to the length shown on the upper body section of the dosing card (2.25 inches).




For each lower body area (a foot, an ankle, or a knee): Squeeze gel from the tube equal to the length shown on the lower body section of the dosing card (4.5 inches).

Lost Your Dosing Card? Don't worry!
 Call 1-800-969-5357 (toll-free) or 215-273-8755 (collect) to get a free replacement.

Applying Motrin[®] Arthritis Pain Gel
 Gently rub Motrin[®] Arthritis Pain Gel into the skin using your hand. Apply 4 times a day for best results. Do not use on more than 2 body areas at once.
 Apply only to clean, dry skin that doesn't have any cuts, open wounds, infections or rashes. Do not apply in the same area as any other medicines or products that are applied to the skin.



After You Use Motrin[®] Arthritis Pain Gel



Wash your hands after applying Motrin[®] Arthritis Pain Gel. If the treated area is the hands wait up to 1 hour to wash the hands.



Wash the dosing card with water. Store dosing card with your Motrin[®] Arthritis Pain Gel and keep out of reach of children.



Wait ten minutes before covering the treated areas with clothes or gloves.

Avoid:
 • showering or bathing for at least 1 hour after use
 • exposing the treated area to sunlight or artificial light (such as in tanning booths)
 • use in same area as other products applied to the skin
 • applying a bandage over the treated area
 • eye, nose, or mouth contact
 • if eye contact occurs, rinse thoroughly with water

Facts you need to know
Important information about Motrin[®] Arthritis Pain Gel
Do not use Motrin[®] Arthritis Pain Gel...
 • if you have ever had an allergic reaction to any other pain reliever or fever reducers
 • for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.
 • right before or after heart surgery
 • on more than 2 body areas at the same time
 • in the eyes, nose or mouth

Ask a doctor before use if...
 • you have problems or serious side effects from taking pain relievers or fever reducers
 • stomach bleeding warning applies to you
 • you have a history of stomach problems such as heartburn
 • you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 • you are taking a diuretic
 • you are under age 18 years. It is not known if this drug works or is safe in children under age 18 years.

Ask a doctor or pharmacist before use if...
 • you are under a doctor's care for any serious condition
 • you are taking any other drug
 • you are pregnant or breast-feeding

Turn over for more information

It is especially important not to use diclofenac at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Warnings to consider before use...

Allergy alert
 Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 • hives • blisters
 • asthma (wheezing) • facial swelling
 • skin reddening • shock
 • rash
 If an allergic reaction occurs, stop use and seek medical help right away.

Liver warning
 This product contains diclofenac. Liver damage may occur if you apply:
 • more or for a longer time than directed
 • when using other drugs containing diclofenac

Stomach bleeding warning
 This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:
 • are age 60 or older
 • have had stomach ulcers or bleeding problems
 • take a blood thinning (anticoagulant) or steroid drug
 • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 • have 3 or more alcoholic drinks every day while using this product
 • apply more or for longer than directed

Heart attack and stroke warning
 NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. This can be fatal. The risk is higher if you use more than directed or for longer than directed.

Follow these important guidelines when using Motrin[®] Arthritis Pain Gel...
 • use only as directed
 • apply only to clean, dry skin that does not have any cuts, open wounds, infections or rashes
 • do not apply in the same area as any other product used on the skin
 • do not apply with external heat such as heating pad
 • do not apply a bandage over the treated area
 • do not get gel in eyes, nose or mouth
 • store ENCLOSED DOSING CARD with your Motrin[®] Arthritis Pain product. The dosing card is re-usable.
Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Stop using Motrin[®] Arthritis Pain Gel and ask a doctor if...

- pain gets worse or lasts more than 21 days
- redness or swelling is present in the painful area
- fever occurs
- skin irritation occurs
- any new symptoms appear. These could be signs of a serious condition.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - leg swelling
 - weakness in one part or side of body
 - slurred speech

Questions and Answers

Answers to questions you may ask about Motrin[®] Arthritis Pain Gel

Can I use Motrin[®] Arthritis Pain Gel for the immediate relief of pain like that associated with sprains, strains, bruises, or sports injuries?
No.
 Do not use for sprains, sprains, bruises, or sports injuries. This product has not been shown to work for these types of injuries. It's intended only for the temporary relief of arthritis pain.

How quickly will Motrin[®] Arthritis Pain Gel work for arthritis pain?
 Motrin[®] Arthritis Pain Gel may take up to 7 days to work for your arthritis pain; it is not for immediate relief. With 4 times-a-day use, you may start to feel relief within a few days. You should feel significant pain relief within 7 days of continuous use. If no pain relief in 7 days, stop use.

What is Motrin[®] Arthritis Pain Gel used for?
 Motrin[®] Arthritis Pain Gel can be used for the temporary relief of arthritis pain in the hands, wrists, elbows, feet, ankles, and knees. Osteoarthritis (OA) is the most common form of arthritis. It mainly affects a type of tissue called cartilage, which cushions the joints and prevents the bones from rubbing against each other. With OA, cartilage begins to break down and wear away, resulting in joint pain and stiffness. OA occurs more frequently as you get older. Pain develops slowly and can gradually worsen over time.



How does Motrin[®] Arthritis Pain Gel work?
 Diclofenac sodium, the medicine in Motrin[®] Arthritis Pain Gel, is a nonsteroidal anti-inflammatory drug (NSAID). NSAIDs are used to treat pain from medical conditions such as arthritis. Motrin[®] Arthritis Pain Gel works similarly to oral NSAIDs like ibuprofen or naproxen by temporarily blocking the production of pain signaling chemicals called prostaglandins. However, when taking a pill, the medication is distributed to the site of pain through the bloodstream. Compared to oral diclofenac, only 6% of

the medicine in Motrin[®] Arthritis Pain Gel is absorbed in the bloodstream. It works by penetrating through your skin at the application site to deliver arthritis pain relief.

How does over-the-counter Diclofenac Sodium Topical Gel, 1% differ from prescription Diclofenac Sodium Topical Gel, 1%?
 Over-the-counter Diclofenac Sodium Topical Gel, 1% is the same strength and formula as original full prescription strength Diclofenac Sodium Topical Gel, 1% but is available without a prescription.

Apply Motrin[®] Arthritis Pain Gel 4 times a day to ensure you get the full effect of the medicine.

Can I apply Motrin[®] Arthritis Pain Gel to upper and lower body areas at the same time?

Yes.
 You can use Motrin[®] Arthritis Pain Gel on up to 2 body areas at the same time. The areas can include the foot, ankle, knee, hand, wrist, and elbow.

Some correct examples include:



Do not use Motrin[®] Arthritis Pain Gel on more than 2 body areas at once.

Can I apply Motrin[®] Arthritis Pain Gel without measuring it out on the dosing card first?

No.
 Use the dosing card to make sure you are getting the correct amount of medicine.

Can I use Motrin[®] Arthritis Pain Gel on my spine, hips, or shoulders?
No.
 The use of Motrin[®] Arthritis Pain Gel on the spine, hips, and shoulders has not been studied.

Can I take any other pain medications with Motrin[®] Arthritis Pain Gel?
 Ask your doctor or pharmacist before using Motrin[®] Arthritis Pain Gel if you are taking any other pain reliever, especially prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others).

Can I take any other pain medications during the first 7 days I use Motrin[®] Arthritis Pain Gel?
 Taking other pain medications is generally not recommended. Speak with your doctor or pharmacist before taking other pain medications when using Motrin[®] Arthritis Pain Gel.

How long can I use Motrin[®] Arthritis Pain Gel?
 Use Motrin[®] Arthritis Pain Gel 4 times a day every day for up to 21 days for treatment of arthritis pain or as directed by your doctor.

Motrin[®] Arthritis Pain Gel should be part of an overall program for managing your arthritis pain. The program should include activities such as appropriate movement or exercise, specific lifestyle changes, and weight maintenance. Regular visits and discussions with your doctor can help incorporate Motrin[®] Arthritis Pain Gel into a complete program to help manage your arthritis pain.

What do I do if I get Motrin[®] Arthritis Pain Gel in my eyes?
 If eye contact occurs, rinse thoroughly with water and consult a doctor if irritation lasts more than an hour.

How do I clean the dosing card?
 Wash the dosing card with water, and store dosing card and tube out of reach of children.

What if I miss a dose of Motrin[®] Arthritis Pain Gel?
 If you forget to apply Motrin[®] Arthritis Pain Gel don't double your dose. Just apply the next dose as scheduled.

How should I store Motrin[®] Arthritis Pain Gel?
 • Store at 20-25°C (68-77°F).
 • Do not freeze Motrin[®] Arthritis Pain Gel.
 • Store the dosing card with your Motrin[®] Arthritis Pain Gel.
 • Keep out of reach of children.

Additional Questions?
 Call 1-800-969-5357 (toll-free) or 215-273-8755 (collect) or visit www.motrin.com

Made in Israel
 Distributed by:
 JOHNSON & JOHNSON CONSUMER INC.
 McNeil Consumer Healthcare Division
 Fort Washington, PA 19034 USA © J&JCI 2021
 3051316

ENR08R 11

MOTRIN ARTHRITIS PAIN

diclofenac sodium gel

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:50580-574

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DICLOFENAC SODIUM (UNII: QTG126297Q) (DICLOFENAC - UNII:144O8QL0L1)	DICLOFENAC SODIUM	9.3 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
CARBOMER HOMOPOLYMER TYPE C (UNII: 4Q93RCW27E)	
COCOYL CAPRYLOCAPRATE (UNII: 8D9H4QU99H)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
AMMONIA (UNII: 5138Q19F1X)	
MINERAL OIL (UNII: T5L8T28FGP)	
POLYOXYL 20 CETOSTEARYL ETHER (UNII: YRC528SWUY)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-574-01	1 in 1 CARTON	06/20/2022	
1		50 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:50580-574-02	1 in 1 CARTON	06/20/2022	
2		100 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA211253	06/20/2022	

Labeler - Johnson & Johnson Consumer Inc. (878046358)

