

**ALKA-SELTZER ORIGINAL- anhydrous citric acid, aspirin, and sodium bicarbonate tablet, effervescent**

**R J General Corporation**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Alka-Seltzer® Original**

**Drug Facts**

<b>Active ingredients (in each tablet)</b>	<b>Purpose</b>
Anhydrous citric acid 1000 mg	Antacid
Aspirin 325 mg (NSAID)*	Analgesic
Sodium bicarbonate (heat-treated) 1916 mg	Antacid

\* nonsteroidal anti-inflammatory drug

**Uses**

for the temporary relief of:

- heartburn, acid indigestion, and sour stomach when accompanied with headache or body aches and pains
- upset stomach with headache from overindulgence in food or drink
- headache, body aches, and pain alone

**Warnings**

**Reye's syndrome**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.
- taking a prescription drug for diabetes, gout, or arthritis

**When using this product** do not exceed recommended dosage

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- symptoms get worse or last more than 10 days
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs
- new symptoms occur

**If pregnant or breast-feeding**, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

Directions

- fully dissolve 2 tablets in 4 ounces of water before taking

---

adults and children 12 years and over	2 tablets every 4 hours, or as directed by a doctor	do not exceed 8 tablets in 24 hours
adults 60 years and over	2 tablets every 4 hours, or as directed by a doctor	do not exceed 4 tablets in 24 hours
children under 12 years	consult a doctor	

---

### **Other information**

- **each tablet contains:** sodium 567 mg

- store at room temperature. Avoid excessive heat.
- Alka-Seltzer in water contains principally the antacid sodium citrate and the analgesic sodium acetylsalicylate

### **Inactive ingredients**

none

### **Questions or comments?**

**1-800-986-0369** (Mon – Fri 9AM – 5PM EST)

Dist. by: Bayer HealthCare LLC

Whippany, NJ 07981

### **PRINCIPAL DISPLAY PANEL - 100 Tablet Pouch Carton**

***Alka-  
Seltzer®***

*Anhydrous citric acid / Antacid*

*Aspirin (NSAID) / Analgesic*

*Sodium bicarbonate / Antacid*

***Fast Relief of:***

***Heartburn***

***Upset Stomach***

***Acid Indigestion***

***with Headache***

***Body Aches***

***ORIGINAL***



**Fast Relief of:**  
Heartburn  
Upset Stomach  
Acid Indigestion  
with Headache  
Body Aches

**ORIGINAL**



**Fast Relief of:**  
Heartburn  
Upset Stomach  
Acid Indigestion  
with Headache  
Body Aches

**ORIGINAL**



**50** POUCHES OF **2** TABLETS

60 TABLETS OF 2 TABLETS



This product is repackaged by RJ General Corp, 2350 Burnet Ave., Cincinnati, Ohio 45219 (513)621-3738 (Mon-Fri 9AM-3PM EST) from a product manufactured by Bayer. Bayer®, the Bayer Cross® and Alka-Seltzer® is a registered trademark of Bayer, used with permission.

**Warnings:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: hives • facial swelling • asthma (wheezing) • shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

**Do not use if you are allergic to aspirin or any other pain reliever/fever reducer**

**Ask a doctor before use if • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma • you have a sodium-restricted diet**

**Ask a doctor or pharmacist before use if you are**

- presently taking a prescription drug. Antacids may interact with certain prescription drugs. • taking a prescription drug for diabetes, gout, or arthritis

**When using this product do not exceed recommended dosage**

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
- feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better
- symptoms get worse or last more than 10 days • redness or swelling is present • ringing in the ears or a loss of hearing occurs
- new symptoms occur

**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.**

**Directions**

• fully dissolve 2 tablets in 4 ounces of water before taking

adults and children 12 years and over	2 tablets every 4 hours, or as directed by a doctor	do not exceed 8 tablets in 24 hours
adults 60 years and over	2 tablets every 4 hours, or as directed by a doctor	do not exceed 4 tablets in 24 hours
children under 12 years	consult a doctor	

**Other information • each tablet contains: sodium 567 mg • store at room temperature. Avoid excessive heat. • Alka-Seltzer® in water contains principally the antacid sodium citrate and the analgesic sodium acetylsalicylate**

**Inactive ingredients** none

**Questions or comments? 1-800-986-0369 (Mon-Fri 9AM-5PM EST) or www.alkaseltzer.com**

**Drug Facts**

**Active Ingredients (in each tablet)**

Anhydrous citric acid 1000 mg.....  
 Aspirin 325 mg (NSAID).....  
 Sodium bicarbonate (heat-treated) 1916 mg.....  
 nonsteroidal anti-inflammatory drug

**Purpose**

Analgesic.....  
 Antacid.....

**Uses for the relief of:**

- heartburn, acid indigestion, and sour stomach when accompanied with headache or body aches and pains
- upset stomach with headache from overindulgence in food or drink
- headache, body aches, and pain alone



**Alka-Seltzer® ORIGINAL**

Place two (2) tablets in 4 oz of water. Dissolve tablets completely.

You do not have to drink any of the residue that may be on the bottom of the glass. The medicines of Alka-Seltzer® are in the water.

Take Alka-Seltzer® any time – morning, noon, or night – when you need relief from heartburn, upset stomach, acid indigestion with headache or body aches.



**Alka-Seltzer® ORIGINAL**

Place two (2) tablets in 4 oz of water. Dissolve tablets completely.

You do not have to drink any of the residue that may be on the bottom of the glass. The medicines of Alka-Seltzer® are in the water.

Take Alka-Seltzer® any time – morning, noon, or night – when you need relief from heartburn, upset stomach, acid indigestion with headache or body aches.

 Bayer HealthCare  
 Bayer HealthCare LLC  
 Whippany, NJ 07981 USA

Bayer, the Bayer Cross and Alka-Seltzer are registered trademarks of Bayer, used with permission. Bayer

**ALKA-SELTZER ORIGINAL**

anhydrous citric acid, aspirin, and sodium bicarbonate tablet, effervescent

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:70264-014(NDC:0280-4000)
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) (ANHYDROUS CITRIC ACID - UNII:XF417D3PSL)	ANHYDROUS CITRIC ACID	1000 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
SODIUM BICARBONATE (UNII: 8MDF5V39QO) (BICARBONATE ION - UNII:HN1ZRA3Q20)	SODIUM BICARBONATE	1916 mg

**Product Characteristics**

Color	white	Score	no score
Shape	ROUND	Size	25mm
Flavor		Imprint Code	ALKA;SELTZER
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70264-014-01	50 in 1 CARTON	03/02/2010	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	03/02/2010	

**Labeler** - R J General Corporation (122542830)**Establishment**

Name	Address	ID/FEI	Business Operations
R J General Corporation		122542830	REPACK(70264-014)

Revised: 8/2018

R J General Corporation