

**ALKA-SELTZER ORIGINAL- anhydrous citric acid, aspirin, and sodium bicarbonate tablet, effervescent**  
**R J General Corporation**

-----  
**Alka-Seltzer® Original**

**Drug Facts**

<b>Active ingredients (in each tablet)</b>	<b>Purpose</b>
Anhydrous citric acid 1000 mg	Antacid
Aspirin 325 mg (NSAID) *	Analgesic
Sodium bicarbonate (heat-treated) 1916 mg	Antacid

\* nonsteroidal anti-inflammatory drug

**Uses**

for the temporary relief of:

- heartburn, acid indigestion, and sour stomach when accompanied with headache or body aches and pains
- upset stomach with headache from overindulgence in food or drink
- headache, body aches, and pain alone

**Warnings**

**Reye's syndrome**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.
- taking a prescription drug for diabetes, gout, or arthritis

**When using this product** do not exceed recommended dosage

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- symptoms get worse or last more than 10 days
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs
- new symptoms occur

**If pregnant or breast-feeding**, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

Directions

- fully dissolve 2 tablets in 4 ounces of water before taking

---

adults and children 12      2 tablets every 4 hours, or as      do not exceed 8 tablets

years and over	directed by a doctor	in 24 hours
adults 60 years and over	2 tablets every 4 hours, or as directed by a doctor	do not exceed 4 tablets in 24 hours
children under 12 years	consult a doctor	

---

### **Other information**

- **each tablet contains:** sodium 567 mg
- store at room temperature. Avoid excessive heat.
- Alka-Seltzer in water contains principally the antacid sodium citrate and the analgesic sodium acetylsalicylate

### **Inactive ingredients**

none

### **Questions or comments?**

**1-800-986-0369** (Mon – Fri 9AM – 5PM EST)

Dist. by: Bayer HealthCare LLC  
Whippany, NJ 07981

### **PRINCIPAL DISPLAY PANEL - 100 Tablet Pouch Carton**

**Alka-  
Seltzer** ®

*Anhydrous citric acid / Antacid  
Aspirin ( **NSAID**) / Analgesic  
Sodium bicarbonate / Antacid*

**Fast Relief of:**  
**Heartburn**  
**Upset Stomach**  
**Acid Indigestion**  
**with Headache**  
**Body Aches**

**ORIGINAL**



**Fast Relief of:**  
Heartburn  
Upset Stomach  
Acid Indigestion  
with Headache  
Body Aches

**ORIGINAL**



**Fast Relief of:**  
Heartburn  
Upset Stomach  
Acid Indigestion  
with Headache  
Body Aches

**ORIGINAL**



**50** POUCHES OF **2** TABLETS

60 TABLETS OF 2 TABLETS



This product is repackaged by RJ General Corp, 2350 Burnet Ave., Cincinnati, Ohio 45219 (513)621-3738 (Mon-Fri 9AM-3PM EST) from a product manufactured by Bayer. Bayer®, the Bayer Cross® and Alka-Seltzer® is a registered trademark of Bayer, used with permission.

**Warnings:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: hives • facial swelling • asthma (wheezing) • shock. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

**Do not use if you are allergic to aspirin or any other pain reliever/fever reducer**

**Ask a doctor before use if • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma • you have a sodium-restricted diet**

**Ask a doctor or pharmacist before use if you are**

- presently taking a prescription drug. Antacids may interact with certain prescription drugs. • taking a prescription drug for diabetes, gout, or arthritis

**When using this product do not exceed recommended dosage**

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
- feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better
- symptoms get worse or last more than 10 days • redness or swelling is present • ringing in the ears or a loss of hearing occurs
- new symptoms occur

**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.**

**Directions**

• fully dissolve 2 tablets in 4 ounces of water before taking

adults and children 12 years and over	2 tablets every 4 hours, or as directed by a doctor	do not exceed 8 tablets in 24 hours
adults 60 years and over	2 tablets every 4 hours, or as directed by a doctor	do not exceed 4 tablets in 24 hours
children under 12 years	consult a doctor	

**Other information • each tablet contains: sodium 567 mg • store at room temperature. Avoid excessive heat. • Alka-Seltzer® in water contains principally the antacid sodium citrate and the analgesic sodium acetylsalicylate**

**Inactive ingredients** none

**Questions or comments? 1-800-986-0369 (Mon-Fri 9AM - 5PM EST) or www.alkaseltzer.com**

**Drug Facts**

**Active Ingredients (in each tablet)**

Anhydrous citric acid 1000 mg.....Antacid

Aspirin 325 mg (NSAID).....Analgesic

Sodium bicarbonate (heat-treated) 1916 mg.....Antacid

**Uses for the relief of:**

- heartburn, acid indigestion, and sour stomach when accompanied with headache or body aches and pains
- upset stomach with headache from overindulgence in food or drink
- headache, body aches, and pain alone

**Nonsteroidal anti-inflammatory drug**



**Alka-Seltzer® ORIGINAL**

Place two (2) tablets in 4 oz of water. Dissolve tablets completely.

You do not have to drink any of the residue that may be on the bottom of the glass. The medicines of Alka-Seltzer® are in the water.

Take Alka-Seltzer® any time – morning, noon, or night – when you need relief from heartburn, upset stomach, acid indigestion with headache or body aches.



**Alka-Seltzer® ORIGINAL**

Place two (2) tablets in 4 oz of water. Dissolve tablets completely.

You do not have to drink any of the residue that may be on the bottom of the glass. The medicines of Alka-Seltzer® are in the water.

Take Alka-Seltzer® any time – morning, noon, or night – when you need relief from heartburn, upset stomach, acid indigestion with headache or body aches.

**Bayer HealthCare**  
Bayer HealthCare LLC  
Whippany, NJ 07981 USA

Bayer, the Bayer Cross and Alka-Seltzer are registered trademarks of Bayer, used with permission. Bayer

<b>ALKA-SELTZER ORIGINAL</b>			
anhydrous citric acid, aspirin, and sodium bicarbonate tablet, effervescent			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:70264-014(NDC:0280-4000)
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO) (BICARBONATE ION - UNII:HN1ZRA3Q20)	SODIUM BICARBONATE	1916 mg
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL) (ANHYDROUS CITRIC ACID - UNII:XF417D3PSL)	ANHYDROUS CITRIC ACID	1000 mg

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	25mm
<b>Flavor</b>		<b>Imprint Code</b>	ALKA;SELTZER
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70264-014-01	50 in 1 CARTON	03/02/2010	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	03/02/2010	

**Labeler** - R J General Corporation (122542830)

## Establishment

Name	Address	ID/FEI	Business Operations
R J General Corporation		122542830	repack(70264-014)

Revised: 10/2023

R J General Corporation