

**MUCINEX FAST-MAX DAY COLD AND FLU AND MUCINEX NIGHTSHIFT NIGHT SEVERE COLD AND FLU MAXIMUM STRENGTH- acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride, and triprolidine hydrochloride**  
**RB Health (US) LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Mucinex® Fast-Max® Day Cold & Flu and Mucinex Nightshift Night Severe Cold & Flu**

**Maximum Strength**

**Drug Facts**

---

| <b>Active ingredients (in each caplet)</b> | <b>Purposes</b>                    |
|--|------------------------------------|
| <b>Mucinex Fast-Max Day Cold &amp; Flu</b> |                                    |
| <b>Acetaminophen 325 mg</b>                | <b>Pain reliever/fever reducer</b> |
| Dextromethorphan HBr 10 mg                 | Cough suppressant                  |
| Guaifenesin 200 mg                         | Expectorant                        |
| Phenylephrine HCl 5 mg                     | Nasal decongestant                 |

---

---

| <b>Active ingredients (in each caplet)</b>            | <b>Purposes</b>                    |
|---|------------------------------------|
| <b>Mucinex Nightshift Night Severe Cold &amp; Flu</b> |                                    |
| <b>Acetaminophen 325 mg</b>                           | <b>Pain reliever/fever reducer</b> |
| Dextromethorphan HBr 10 mg                            | Cough suppressant                  |
| Phenylephrine HCl 5 mg                                | Nasal decongestant                 |
| Triprolidine HCl 1.25 mg                              | Antihistamine                      |

---

**Uses**

**Mucinex Fast-Max Day Cold & Flu**

- temporarily relieves these common cold and flu symptoms:
  - cough
  - nasal congestion
  - minor aches and pains
  - sore throat
  - headache

- stuffy nose
- sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

### **Mucinex Nightshift Night Severe Cold & Flu**

- temporarily relieves these common cold and flu symptoms:
  - cough
  - nasal congestion
  - minor aches and pains
  - sore throat
  - headache
  - sneezing
  - sinus congestion and pressure
  - runny nose
  - itching of the nose or throat
  - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

### **Warnings**

#### **Liver warning**

This product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

#### **Allergy alert**

Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

#### **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### **Ask a doctor before use if you have**

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma (***Nightshift Night Severe Cold & Flu only***)
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis (***Nightshift Night Severe Cold & Flu only***)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

#### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (***Nightshift Night Severe Cold & Flu only***)

#### **When using this product**

- **do not use more than directed**
- excitability may occur, especially in children (***Nightshift Night Severe Cold & Flu only***)
- marked drowsiness may occur (***Nightshift Night Severe Cold & Flu only***)
- alcohol, sedatives, and tranquilizers may increase drowsiness (***Nightshift Night Severe Cold & Flu only***)
- avoid alcoholic drinks (***Nightshift Night Severe Cold & Flu only***)
- use caution when driving a motor vehicle or operating machinery (***Nightshift Night Severe Cold & Flu only***)

#### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

## **Overdose warning**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## **Directions**

### **Mucinex Fast-Max Day Cold & Flu**

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

### **Mucinex Nightshift Night Severe Cold & Flu**

- **do not take more than directed (see Overdose warning)**
- do not take more than 8 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

## **Other information**

- store at 20-25°C (68-77°F)

## **Inactive ingredients Mucinex Fast-Max Day Cold & Flu**

croscarmellose sodium, crospovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

## **Inactive ingredients Mucinex Nightshift Night Severe Cold & Flu**

croscarmellose sodium, crospovidone, ferric oxide, hydroxypropyl cellulose, mica, microcrystalline cellulose, polyvinyl alcohol, polyvinyl alcohol polyethylene glycol copolymer, povidone, silicon dioxide, stearic acid, talc, titanium dioxide

## **Questions?**

**1-866-MUCINEX (1-866-682-4639)** You may also report side effects to this phone number.

Dist. by: RB Health (US)  
Parsippany, NJ 07054-0224

## **PRINCIPAL DISPLAY PANEL - Kit Carton**

FAST RELEASE. POWERFUL SYMPTOM RELIEF!

NDC 72854-241-20

MAXIMUM STRENGTH

Mucinex®

FAST-MAX® NIGHTSHIFT

DAY

COLD & FLU

Acetaminophen – Pain Reliever/

Fever Reducer

Dextromethorphan HBr – Cough Suppressant

Guaifenesin – Expectorant

Phenylephrine HCl – Nasal Decongestant

HEADACHE

SORE THROAT

CHEST CONGESTION

BODY PAIN

FEVER

COUGH

ALL IN

ONE\*

NASAL CONGESTION

SINUS CONGESTION

SINUS PRESSURE

ACTUAL SIZE

FOR AGES 12+

12 CAPLETS

FAST

RELEASE

NIGHT

SEVERE COLD & FLU

Acetaminophen – Pain Reliever/

Fever Reducer

Dextromethorphan HBr – Cough Suppressant

Phenylephrine HCl – Nasal Decongestant

Tripolidine HCl – Antihistamine

HEADACHE

SORE THROAT

ITCHY THROAT

BODY PAIN

FEVER

COUGH

ALL IN

ONE\*

NASAL CONGESTION

SNEEZING  
RUNNY NOSE  
ACTUAL SIZE  
FOR AGES 12+  
8 CAPLETS  
TOTAL 20 CAPLETS

Tamper evident: Do not use if carton is damaged or if printed seal on blister is broken or missing.

MAXIMUM STRENGTH

**Mucinex**  
FAST-MAX NIGHTSHIFT

**DAY**  
COLD & FLU

**NIGHT**  
SEVERE COLD & FLU

FAST RELEASE. POWERFUL SYMPTOM RELIEF!

MAXIMUM STRENGTH

NDC 72854-241-20

**Mucinex**

FAST-MAX NIGHTSHIFT

**DAY**  
COLD & FLU

Acetaminophen - Pain Reliever/  
Fever Reducer  
Dextromethorphan HBr - Cough Suppressant  
Guaifenesin - Expectorant  
Phenylephrine HCl - Nasal Decongestant

**NIGHT**  
SEVERE COLD & FLU

Acetaminophen - Pain Reliever/  
Fever Reducer  
Dextromethorphan HBr - Cough Suppressant  
Phenylephrine HCl - Nasal Decongestant  
Triprolidine HCl - Antihistamine



ACTUAL SIZE  
FOR AGES 12+



FAST  
RELEASE



ACTUAL SIZE  
FOR AGES 12+



12 CAPLETS

TOTAL 20 CAPLETS

8 CAPLETS

MAXIMUM STRENGTH  
**Mucinex**  
FAST-MAX NIGHTSHIFT

**DAY** **NIGHT**  
COLD & FLU SEVERE COLD & FLU  
Maximum Strength per 4-hour dose  
Keep carton for full information.

Fast-Max Day Cold & Flu helps to relieve these symptoms day or night. Nightshift Night Severe Cold & Flu helps to relieve these symptoms at night. Do not take the Mucinex Fast-Max Day Cold & Flu and Mucinex Nightshift Night Severe Cold & Flu caplets at the same time. Always wait at least 4 hours before taking another dose of Mucinex caplets. TAKE ONLY AS DIRECTED.

**PARENTS:**  
Learn about their medicine advice  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**Questions?**  
1-866-MUCINEX (1-866-682-4639)  
You may also report side effects to this phone number.




HEALTH • HYGIENE • HOME

[www.mucinex.com](http://www.mucinex.com)

Patents: [www.rb.com/patents](http://www.rb.com/patents)

Dist. by: RB Health (US)  
Parsippany, NJ 07054-0224  
©2021 RB Health

3176344 012721



6382499531 1

LOT:

EXP.:

MADE IN:

3176344



Do not take the Mucinex Fast-Max Day Cold & Flu and Mucinex Nightshift Night Severe Cold & Flu caplets at the same time. Always wait at least 4 hours before taking another dose of Mucinex caplets. TAKE ONLY AS DIRECTED.

3176344

**Drug Facts**

Active ingredients (in each caplet)

Purposes



010621 PEEL CORNER TO READ COMPLETE DRUG FACTS AND INFORMATION

**Mucinex Fast-Max Day Cold & Flu**

|                                 |                             |
|---------------------------------|-----------------------------|
| Acetaminophen 325 mg.....       | Pain reliever/fever reducer |
| Dextromethorphan HBr 10 mg..... | Cough suppressant           |
| Guaifenesin 200 mg.....         | Expectorant                 |
| Phenylephrine HCl 5 mg.....     | Nasal decongestant          |

**Active ingredients (in each caplet)**

**Purposes**

**Mucinex Nightshift Night Severe Cold & Flu**

|                                 |                             |
|---------------------------------|-----------------------------|
| Acetaminophen 325 mg.....       | Pain reliever/fever reducer |
| Dextromethorphan HBr 10 mg..... | Cough suppressant           |
| Phenylephrine HCl 5 mg.....     | Nasal decongestant          |
| Triprolidine HCl 1.25 mg.....   | Antihistamine               |

**Uses**

**Mucinex Fast-Max Day Cold & Flu**

- temporarily relieves these common cold and flu symptoms:
  - cough
  - nasal congestion
  - minor aches and pains
  - sore throat
  - headache
  - stuffy nose
  - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Mucinex Nightshift Night Severe Cold & Flu**

- temporarily relieves these common cold and flu symptoms:
  - cough
  - nasal congestion
  - minor aches and pains
  - sore throat
  - headache
  - sneezing
  - sinus congestion and pressure
  - runny nose
  - itching of the nose or throat
  - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

**Warnings**

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:
- more than 4,000 mg in 24 hours, which is the maximum daily amount
  - with other drugs containing acetaminophen
  - 3 or more alcoholic drinks daily while using this product
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
- If a skin reaction occurs, stop use and seek medical help right away.

**Reseal Area**

**Drug Facts (continued)**

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma (*Nightshift Night Severe Cold & Flu only*)
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis (*Nightshift Night Severe Cold & Flu only*)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (*Nightshift Night Severe Cold & Flu only*)

**When using this product**

- do not use more than directed
- excitability may occur, especially in children (*Nightshift Night Severe Cold & Flu only*)
- marked drowsiness may occur (*Nightshift Night Severe Cold & Flu only*)
- alcohol, sedatives, and tranquilizers may increase drowsiness (*Nightshift Night Severe Cold & Flu only*)
- avoid alcoholic drinks (*Nightshift Night Severe Cold & Flu only*)
- use caution when driving a motor vehicle or operating machinery (*Nightshift Night Severe Cold & Flu only*)



Hinge

Hinge

**Drug Facts (continued)**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

**Mucinex Fast-Max Day Cold & Flu**

- do not take more than directed (see Overdose warning)
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

**Mucinex Nightshift Night Severe Cold & Flu**

- do not take more than directed (see Overdose warning)
- do not take more than 8 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

**Other information** ■ store at 20-25°C (68-77°F)

**Inactive ingredients Mucinex Fast-Max Day Cold & Flu**

croscarmellose sodium, crospovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

**Inactive ingredients Mucinex Nightshift Night Severe Cold & Flu**

croscarmellose sodium, crospovidone, ferric oxide, hydroxypropyl cellulose, mica, microcrystalline cellulose, polyvinyl alcohol, polyvinyl alcohol polyethylene glycol copolymer, povidone, silicon dioxide, stearic acid, talc, titanium dioxide

**Questions?** 1-866-MUCINEX (1-866-682-4639) You may also report side effects to this phone number.

**MUCINEX FAST-MAX DAY COLD AND FLU AND MUCINEX NIGHTSHIFT NIGHT SEVERE COLD AND FLU MAXIMUM STRENGTH**

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride, and triprolidine hydrochloride kit

**Product Information**

|                     |                |                           |               |
|---------------------|----------------|---------------------------|---------------|
| <b>Product Type</b> | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:72854-241 |
|---------------------|----------------|---------------------------|---------------|

**Packaging**

| # | Item Code        | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:72854-241-10 | 1 in 1 CARTON       | 07/01/2021           |                    |
| 1 |                  | 1 in 1 KIT          |                      |                    |
| 2 | NDC:72854-241-20 | 2 in 1 CARTON       | 07/01/2021           |                    |

|   |                  |               |            |
|---|------------------|---------------|------------|
| 2 |                  | 1 in 1 KIT    |            |
| 3 | NDC:72854-241-40 | 4 in 1 CARTON | 07/01/2021 |
| 3 |                  | 1 in 1 KIT    |            |

### Quantity of Parts

| Part # | Package Quantity | Total Product Quantity |
|--------|------------------|------------------------|
| Part 1 | 1 BLISTER PACK   | 6                      |
| Part 2 | 1 BLISTER PACK   | 4                      |

### Part 1 of 2

#### MAXIMUM STRENGTH MUCINEX FAST-MAX COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, film coated

### Product Information

|                         |               |
|-------------------------|---------------|
| Item Code (Source)      | NDC:63824-575 |
| Route of Administration | ORAL          |

### Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength             | Strength |
|--|-------------------------------|----------|
| <b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)                    | ACETAMINOPHEN                 | 325 mg   |
| <b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg    |
| <b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)                        | GUAIFENESIN                   | 200 mg   |
| <b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)      | PHENYLEPHRINE HYDROCHLORIDE   | 5 mg     |

### Inactive Ingredients

| Ingredient Name  | Strength |
|--|----------|
| <b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)             |          |
| <b>CROSPVIDONE</b> (UNII: 2S7830E561)                      |          |
| <b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)              |          |
| <b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)            |          |
| <b>ALUMINUM OXIDE</b> (UNII: LMI26O6933)                   |          |
| <b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)               |          |
| <b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)       |          |
| <b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A) |          |
| <b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)   |          |
| <b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)            |          |
| <b>TALC</b> (UNII: 7SEV7J4R1U)                             |          |
| <b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)                 |          |

## Product Characteristics

|                 |      |                     |          |
|-----------------|------|---------------------|----------|
| <b>Color</b>    | red  | <b>Score</b>        | no score |
| <b>Shape</b>    | OVAL | <b>Size</b>         | 20mm     |
| <b>Flavor</b>   |      | <b>Imprint Code</b> | MSC;VW   |
| <b>Contains</b> |      |                     |          |

## Packaging

| # | Item Code | Package Description                                    | Marketing Start Date | Marketing End Date |
|---|-----------|--|----------------------|--------------------|
| 1 |           | 6 in 1 BLISTER PACK; Type 0: Not a Combination Product |                      |                    |

## Marketing Information

| Marketing Category  | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part341                                  | 07/01/2021           |                    |

## Part 2 of 2

### MUCINEX NIGHTSHIFT SEVERE COLD AND FLU MAXIMUM STRENGTH

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride, and triprolidine hydrochloride tablet, coated

## Product Information

|                                |               |
|--------------------------------|---------------|
| <b>Item Code (Source)</b>      | NDC:72854-234 |
| <b>Route of Administration</b> | ORAL          |

## Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength             | Strength |
|--|-------------------------------|----------|
| <b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)                    | ACETAMINOPHEN                 | 325 mg   |
| <b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg    |
| <b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)      | PHENYLEPHRINE HYDROCHLORIDE   | 5 mg     |
| <b>TRIPROLIDINE HYDROCHLORIDE</b> (UNII: YAN7R5L890) (TRIPROLIDINE - UNII:2L8T9S52QM)        | TRIPROLIDINE HYDROCHLORIDE    | 1.25 mg  |

## Inactive Ingredients

| Ingredient Name                                | Strength |
|--|----------|
| <b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48) |          |

|  |
|--|
| <b>CROSPROVIDONE</b> (UNII: 2S7830E561)  |
| <b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)   |
| <b>HYDROXYPROPYL CELLULOSE, UNSPECIFIED</b> (UNII: 9XZ8H6N6OH)                                   |
| <b>MICA</b> (UNII: V8A1AW0880)   |
| <b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)   |
| <b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)   |
| <b>POLYVINYL ALCOHOL GRAFT POLYETHYLENE GLYCOL COPOLYMER (3:1; 45000 MW)</b> (UNII: 23ZQ42JZ ZH) |
| <b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)  |
| <b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)  |
| <b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)   |
| <b>TALC</b> (UNII: 7SEV7J4R1U)   |
| <b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)   |

| Product Characteristics |        |                     |                         |
|-------------------------|--------|---------------------|-------------------------|
| <b>Color</b>            | yellow | <b>Score</b>        | no score                |
| <b>Shape</b>            | OVAL   | <b>Size</b>         | 19mm                    |
| <b>Flavor</b>           |        | <b>Imprint Code</b> | VW;LOGOcrescentmoonplus |
| <b>Contains</b>         |        |                     |                         |

| Packaging |           |  |                      |                    |
|-----------|-----------|--|----------------------|--------------------|
| #         | Item Code | Package Description                                    | Marketing Start Date | Marketing End Date |
| 1         |           | 4 in 1 BLISTER PACK; Type 0: Not a Combination Product |                      |                    |

| Marketing Information |  |                      |                    |
|-----------------------|--|----------------------|--------------------|
| Marketing Category    | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| OTC monograph final   | part341                                  | 07/01/2021           |                    |

| Marketing Information |  |                      |                    |
|-----------------------|--|----------------------|--------------------|
| Marketing Category    | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| OTC monograph final   | part341                                  | 07/01/2021           |                    |

**Labeler** - RB Health (US) LLC (081049410)