HEMRID ANORECTAL CREAM- lidocaine cream KKA & Associates LLC

Benzyl Alcohol, Carbomer, Cholesterol, Hydrogenated Lecithin (Soya), Isopropyl Myristate, Polysorbate 80, Propylene Glycol, Purified Water, Triethanolamine, Vitamin-E Acetate.

Lidocaine 5% W/W

Local Anesthetic

When practical, cleanse the affected area with mild soap and warm water and rinse thoroughly, gently dry by patting or blotting with toilet tissue or a soft cloth before application of this product.

apply externally to the affected area up to 6 times a day.

Children under 12 years of age: consult a doctor.

To use with finger cot. Roll one finger cot over finger. Gently squeeze cream onto finger Smooth a layer of the cream over the affected area.

Keep out of reach of children and pets. If swallowed, get medical help or contact a Poison Control Center right away.

FOR EXTERNAL USE ONLY

When using this product

use only as directed.

Avoid contact with eyes.

do not exceed recommended dosage unless directed by a doctor.

do not out this product into the rectum by using fingers or any mechanical device or applicator.

Stop use and ask a doctor if

rectal bleeding occurs.

condition worsens or does not improve within 7 days.


symptoms clear up and occur again within a few days.

an allergic reaction develops to ingredients in this product.

symptom being treated does not subside or if redness, irritation, swelling, pain or other symptoms develop or increase.

If pregnant or breast-feeding, ask a health professional before use.

Helps relieve the pain, itching, and burning, associated with hemorrhoids and anorectal disorders.



5% LIDOCAINE ANORECTAL CREAM

RÁPIDA FÓRMULA ALIVIADORA DE DOLOR • PREVIENE IRRITACIÓN ADICIONAL

CREMA ANORRECTAL

5% LIDOCAINI

- · Alivio de ardor, picazón e irritación
- Mayor concentración de lidocaína sin receta disponible
- · Garantía de devolución de dinero de 120 días

FAST PAIN RELIEVING FORMULA · PREVENTS FURTHER IRRITATION

5% LIDOCAINE
ANORECTAL CREAM

- Relief from Burning, Itching, and Irritation
- Highest NonRx Lidocaine Strength Available
- 120 Day Money Back Guarantee

NET WT 1 OZ (28.3 g)

5% LIDOCAINE
ANORECTAL CREAM



поэ.Бітель

Distributed by KKA & Associates, LLC, 10510 Marino Point Drive, Suite 603, Estero, FL 33913

Gnespous 1-888-832-8001

Purified Water, Trolamine, Vitamin E Acetate

Inactive ingredients Benzyl Alcohol, Carbomer, Cholesterol, Hydrogenated Lecitinin, Isopropyl Myntstate, Polysorbate 80, Propylene Glycol,

Other information Store at controlled room temperature 68-777F (20-25°C).

over affected area

When practical, dean area with mild soap and warm water and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or soft cloth
before applying.
 Adults and children 12 years and older: apply externally to the affected area up to 6 times a day.
 Children under 12 years of
age: consult a doctor.
 To use finger cots: Adult one finger cot over finger. Gently squeeze cream onto finger cot. Smooth a layer of the cream

Directions

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a health professional before use.

increase • symptoms clear up and return within a few days

Stop use and ask a doctor if • rectal bleeding occurs • condition worsens or does not improve within 7 days • allergic reaction occurs to ingredients in this product • symptom being treated does not subside or if redness, infitation, swelling, pain or other symptoms develop or

do not put this product into the rectum by using fingers or any mechanical device or applicator

For external use only • When using this product • avoid contact with eyes • do not exceed recommended dosage unless directed by a doctor

Uses Helps relieve the pain, itching, and burning associated with hemorrhoids and other anorectal disorders.

Local anesthetic

Active ingredient Lidocaine 5% w/w

Drug Facts

5% LIDOCAINE

MABRO DATE OF THE PROPERTY OF TH



OTOGOU

HEMRID ANORECTAL CREAM

lidocaine cream

| Product Information | oduct Information | | | | | |
|----------------------------|-------------------|--------------------|---------------|--|--|--|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:82620-001 | | | |
| Route of Administration | TOPICAL | | | | | |

| Active Ingredient/Active Moiety | | | |
|--|-------------------|--------------|--|
| Ingredient Name | Basis of Strength | Strength | |
| LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987) | LIDOCAINE | 50 mg in 1 g | |

| Inactive Ingredients | | |
|--|----------|--|
| Ingredient Name | Strength | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | | |
| TROLAMINE (UNII: 903K93S3TK) | | |
| .ALPHATOCOPHEROL CALCIUM SUCCINATE, DL- (UNII: H5A1374A6R) | | |
| CHOLESTEROL (UNII: 97C5T2UQ7J) | | |
| WATER (UNII: 059QF0KO0R) | | |
| ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS) | | |
| CARBOMER 940 (UNII: 4Q93RCW27E) | | |
| BENZYL ALCOHOL (UNII: LKG8494WBH) | | |
| HYDROGENATED SOYBEAN LECITHIN (UNII: H1109Z9J4N) | | |
| POLYSORBATE 80 (UNII: 6OZP39ZG8H) | | |

| F | Packaging | | | | |
|---|----------------------|---|-------------------------|-----------------------|--|
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date | |
| 1 | NDC:82620-001- 01 | 1 in 1 CARTON | 04/04/2022 | | |
| 1 | | 28.3 g in 1 TUBE; Type 0: Not a Combination Product | | | |

| Marketing Information | | | | |
|-----------------------|---|-------------------------|-----------------------|--|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | |
| OTC Monograph Drug | M015 | 04/04/2022 | | |
| | | | | |

Labeler - KKA & Associates LLC (050979549)

Revised: 12/2023 KKA & Associates LLC