

**NIGHTTIME COLD AND FLU- acetaminophen diphenhydramine hcl  
phenylephrine hcl liquid  
Rite Aid Corporation**

-----

**Drug Facts**

**Active ingredients (in each 20 mL)**

**Acetaminophen 650 mg**

Diphenhydramine HCl 25 mg

Phenylephrine HCl 10 mg

**Purposes**

**Pain reliever/fever reducer**

Antihistamine/Cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves these common cold/flu symptoms:
  - cough
  - nasal congestion
  - minor aches and pains
  - sore throat
  - headache
  - runny nose
  - sneezing
- temporarily reduces fever
- controls cough to help you get to sleep

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- with any other drug containing diphenhydramine, even one used on the skin
- for children under 12 years of age

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

### **Ask a doctor or pharmacist before use if you are taking**

- the blood thinning drug warfarin
- sedatives or tranquilizers

### **When using this product**

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24-hours period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL= milliliter
- keep dosing cup with product
- dose as follows or as directed by a doctor
- adults and children 12 years and over: 20 mL every 4 hours while symptoms last
- children under 12 years of age: do not use

**Other information**

- **each 20 mL contains:** Sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients**

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

**Principal Display Panel**

Compare to the active ingredients in Maximum Strength Mucinex® Fast-Max® Night Time Cold & Flu\*

**MAXIMUM STRENGTH**

**NIGHT TIME**

**COLD & FLU**

**MULTI-SYMPTOM RELIEF**

PAIN RELIEVER / FEVER REDUCER

ANTIHISTAMINE / COUGH SUPPRESSANT

NASAL DECONGESTANT

ACETAMINOPHEN 650 mg

DIPHENHYDRAMINE HCl 25 mg

PHENYLEPHRINE HCl 10 MG

Temporarily relieves aches, fever, sore throat, nasal congestion, runny nose & sneezing

Controls cough

For ages 12 & over

FL OZ (mL)

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor or Maximum Strength Mucinex® Fast-Max® Night Time Cold & Flu

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

DISTRIBUTED BY:

RITE AID, 30 HUNTER LANE,

CAMP HILL, PA 17011

[www.riteaid.com](http://www.riteaid.com)

**Product Label**

Compare to the active ingredients in Maximum Strength Mucinex® Fast-Max® Night Time Cold & Flu\*

NDC 11822-6859-7

**MAXIMUM STRENGTH**

# NIGHT TIME COLD & FLU

**MULTI-SYMPTOM RELIEF**

PAIN RELIEVER / FEVER REDUCER  
ANTIHISTAMINE / COUGH SUPPRESSANT  
NASAL DECONGESTANT

**ACETAMINOPHEN 650 mg**  
**DIPHENHYDRAMINE HCl 25 mg**  
**PHENYLEPHRINE HCl 10 mg**

Temporarily relieves aches, fever, sore throat, nasal congestion, runny nose & sneezing  
Controls cough  
For ages 12 & over



6 FL OZ (177 mL)

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Night Time Cold & Flu.

**DISTRIBUTED BY:**

RITE AID, 30 HUNTER LANE,  
CAMP HILL, PA 17011

[www.riteaid.com](http://www.riteaid.com)

**SATISFACTION GUARANTEE**

If you're not satisfied, we'll happily refund your money.

PLD-A377D  
LB007033

**Drug Facts**

**Active ingredients (in each 20 mL) Purposes**

Acetaminophen 650 mg.....	Pain reliever/fever reducer
Diphenhydramine HCl 25 mg.....	Antihistamine/cough suppressant
Phenylephrine HCl 10 mg.....	Nasal decongestant

PEEL CORNER FOR MORE DRUG FACTS

**Drug Facts (continued)**

**Uses** ■ temporarily relieves these common cold and flu symptoms

- cough ■ nasal congestion
- minor aches and pains ■ sore throat
- headache ■ runny nose ■ sneezing
- temporarily reduces fever
- controls cough to help you get to sleep

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

■ with any other drug containing acetaminophen

**Drug Facts (continued)**

(prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- with any other drug containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

**Ask a doctor before use if you have**

- liver disease ■ heart disease
- high blood pressure ■ thyroid disease
- diabetes ■ glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if**

- you are taking the blood thinning drug warfarin
- you are taking sedatives or tranquilizers

**Drug Facts (continued)**

**When using this product**

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter ■ keep dosing cup with product
- dose as follows or as directed by a doctor
- adults and children 12 years and older: 20 mL every 4 hours while symptoms last
- children under 12 years of age: do not use

**Other information**

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

PEEL CORNER FOR MORE DRUG FACTS

**RITE AID Maximum Strength Night Time Cold & Flu**

# Package label

Compare to the active ingredients in  
**Maximum Strength Mucinex®  
 Fast-Max® Severe Cold\***

NDC 11822-6859-7

**MAXIMUM STRENGTH**

## MUCUS RELIEF SEVERE COLD

**MULTI-SYMPTOM RELIEF**

PAIN RELIEVER / FEVER REDUCER  
 COUGH SUPPRESSANT • EXPECTORANT  
 NASAL DECONGESTANT

**ACETAMINOPHEN 650 mg**  
**DEXTROMETHORPHAN HBr 20 mg**  
**GUAIFENESIN 400 mg**  
**PHENYLEPHRINE HCl 10 mg**

Temporarily relieves aches, fever,  
 sore throat, nasal & chest congestion  
 Controls cough  
 Thins & loosens mucus  
 For ages 12 & over

 **6 FL OZ (177 mL)**

PLD-A346B LB007034

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL  
 AROUND OR UNDER CAP IS BROKEN OR MISSING.**

\*This product is not manufactured or distributed  
 by Reckitt Benckiser, distributor of Maximum  
 Strength Mucinex® Fast-Max® Severe Cold.

PLD-A346B  
 LB007034

**PARENTS:**  
 Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

### Drug Facts

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg.....	Nasal decongestant

PEEL CORNER FOR MORE DRUG FACTS

### Drug Facts (continued)

#### Uses

- temporarily relieves these common cold and flu symptoms
  - cough
  - nasal congestion
  - minor aches and pains
  - sore throat
  - headache
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

#### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
  - blisters
  - rash
- If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or

### Drug Facts (continued)

vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

#### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur

PEEL CORNER FOR MORE DRUG FACTS

### Drug Facts (continued)

- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

do not take more than directed (see Overdose warning)

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL every 4 hours
- children under 12 years of age: do not use

### Drug Facts (continued)

#### Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

#### DISTRIBUTED BY:

RITE AID, 30 HUNTER LANE,  
 CAMP HILL, PA 17011

[www.riteaid.com](http://www.riteaid.com)

#### SATISFACTION GUARANTEE

If you're not satisfied, we'll  
 happily refund your money.

# RITE AID Maximum Strength Mucus Relief Severe Cold

## Package label



## RITE AID Maximum Strength Night Time Cold & Flu, Maximum Strength Mucus Relief Severe Cold

### NIGHTTIME COLD AND FLU

acetaminophen diphenhydramine hcl phenylephrine hcl liquid

#### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-6859
<b>Route of Administration</b>	ORAL		

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg in 20 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

#### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE CALCIUM DISODIUM</b> (UNII: 25IH6R45GF)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYL GALLATE</b> (UNII: 8D4SNN7V92)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	

<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-6859-7	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/31/2021	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/31/2021	

**Labeler** - Rite Aid Corporation (014578892)

Revised: 5/2024

Rite Aid Corporation