VITAMIN E-AD- tocopherol vitamin a vitamin d3 injection MWI/VetOne

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Vitamin E-AD

Vitamin E-AD

Injectable d-alphaTocopherol with AD

For animal use only.

Keep out of reach of children.

Natural vitamin A (carotenes), and tocopherols can be destroyed in feedstuffs through processing, ensiling and storage. Due to these losses, reduced intakes of fat-soluble vitamins can occur in animals maintained in continual confinement compared to animals allowed to graze in lush pasture. Intramuscular or subcutaneous injections offer an efficient and rapid method to increase vitamin A, vitamin D and Vitamin E status of animals.

Indications: Vitamin E-AD injectable d-alpha-tocopherol with AD is a clear, sterile sterile water emulsifiable solution of vitamin A, vitamin D_3 , and vitamin E. The product is intended as a supplemental source of vitamins A, D and E.

EACH mL CONTAINS:

ACTIVE INGREDIENTS: 300 I.U. vitamin E (as d-alpha-tocopherol, a natural source of vitamin E); 100,000 I.U. Vitamin A Propionate; 10,000 I.U. Vitamin D₃

INACTIVE INGREDIENTS: Compounded with 2% benzyl alcohol (preservative) in a water emulsifiable base.

CAUTION:

Do not exceed recommended dosage. Anaphylactoid or allergic reactions have been reported in individual animals as well as entire herds. Should such reactions or hypersensitivity occur, treat immediately with injection of epinephrine and/or antihistamines. Administration of this product to well-nourished animals may cause hypervitaminosis D, which may result in hypercalcemia and hyperphosphatemia.

WARNING:

Do not add water to solution.

STORAGE:

Store between 15°-30°C (59°-86°F) in a dark place.

ADMINISTRATION AND DOSAGE:

Intramuscular or Subcutaneous Administration Only. May be repeated. If dose is greater than 5 mL, equally divide the dose and inject at two different sites.

SUGGESTED DO SAGE

CATTLE (Dairy and Beef)	mL
Cows and Heifers	
2-3 wks pre-partum	8-10
At calving	8-10
Calves At birth	
Weaning	. 4–6
Yearlings	. 5-6

SUGGESTED DO SAGE

CATTLE (Dairy and Beef)	mL
Cows and Heifers 2-3 wks pre-partum	8–10
At calving	8–10
Calves At birth Weaning	4-6
Yearlings	5-6

Natural vitamin A (carotenes), and tocopherols can be destroyed in feedstuffs through processing, ensiling and storage. Due to these losses, reduced intakes of fat-soluble vitamins can occur in animals maintained in continual confinement compared to animals allowed to graze lush pasture. Intramuscular or subcutaneous injections offer an efficient and rapid method to increase vitamin A, vitamin D and vitamin E status of animals. INDICATIONS: Vitamin E-AD injectable d-alpha-

tocopherol with AD is a clear, sterile water emulsifiable solution of vitamin A, vitamin D3, and vitamin E. This product is intended as a supplemental source of vitamins A, D and E

EACH mL CONTAINS:
ACTIVE INGREDIENT: 300 I.U. of vitamin E (as d-alpha-tocopherol, a natural source of vitamin E); 100,000 I.U. vitamin A Propionate; 10,000 I.U. vitamin D3 INACTIVE INGREDIENTS: Compounded with 2% benzyl alcohol (preservative) in a water emulsifiable base.



V-3234-05 Rev. 06/16

Distributed by: MWI Boise, ID 83705 www.VetOne.net

NDC 13985-058-04



Injection

Injectable d-alpha Tocopherol with AD

For animal use only. Keep out of reach of children.

V1 501058 Net Contents: 250 mL (8.5 fl. oz.)

DO SAGE AND ADMINISTRATION: Inframuscular or subcutaneous administration only. May be repeated as needed. If dose is greater than 5 mL, equally divide the dose and inject at two different sites.

SUGGESTED DOSAGE

CATTLE (Dairy and Beef)

250 mL

Cows and Heifers											
2-3 wks pre-par	tun	n.									8-10
At calving											8-10
End of lactation									-		8-10
Calves At birth											
Weaning			8			67					. 4-6

CAUTION: Do not exceed recommended dosage Anaphylactoid or allergic reactions have been reported in individual animals as well as entire herds. Should such reactions or hypersensitivity occur, treat immediately with epinephrine and/or antihistamines Administration of this product to well-nourished animals may cause hypervitaminosis D, which may result in hypercalcemia and hyperphosphatemia.

WARNING: Do not add water to solution

STORAGE: Store between 15°-30°C (59°-86°F) in a dark place.



Lot No.: Exp. Date:

VITAMIN E-AD

Product Information			
Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:13985-058
Route of Administration	INTRAMUSCULAR, SUBCUTANEOUS		

Active Ingredient/Active Moiety							
Ingredient Name	Basis of Strength	Strength					
.ALPHATOCOPHEROL, D- (UNII: N9PR3490H9) (.ALPHATOCOPHEROL, DUNII:N9PR3490H9)	.ALPHA TOCOPHEROL, D-	201 mg in 1 mL					
VITAMIN A (UNII: 81G40H8B0T) (VITAMIN A - UNII:81G40H8B0T)	VITAMIN A	3.6 mg in 1 mL					
VITAMIN D (UNII: 9VU1KI44GP) (CHOLECALCIFEROL - UNII:1C6V77QF41)	VITAMIN D	2.5 mg in 1 mL					

P	ackaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:13985-058-04	250 mL in 1 VIAL		

Marketing End Date

Labeler - MWI/VetOne (019926120)

Revised: 12/2023 MW/VetOne