

E-VAC PLUS- docusate sodium and benzocaine enema
ZURICH MEDICAL LABS, LLC

E-Vac Plus

Drug Facts

Active ingredient (in each 5cc disposable tube)	Purpose
Docusate Sodium 283mg	Stool Softener Enema
Benzocaine 20mg	Topical Analgesic

Uses

- For the relief of occasional constipation and/or aid in preventing constipation
- Helps to reduce pain or rectal damage caused by hard stools or by straining during bowel movements
- usually produces bowel movement in 2 to 15 minutes

Warnings

Ask a doctor before use if you have

- appendicitis
- symptoms of appendicitis (e.g., nausea, vomiting, abdominal pain)
- bleeding from the rectum
- undiagnosed bleeding
- congestive heart failure
- high blood pressure
- fecal impaction
- intestinal obstruction
- been taking other prescriptions

Ask a doctor or pharmacist before use if you are taking mineral oils.

Stop use and ask a doctor if

- bleeding from the rectum
- you fail to have a bowel movement after using this product.

These may indicate a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental ingestion or overdose, get medical help or contact a Poison Control Center right away.

Directions

E-VAC™ MINI-ENEMA & E-VAC™ PLUS MINI-ENEMA is to be used as an enema. **This is not a suppository.** Do not take orally.

Administering the mini enema

Twist off and remove tip. Place water or a few drops of the product on the shaft of the tube to lubricate prior to insertion. It is also recommended that one of these lubricants be applied to the anus before

inserting the enema. With steady pressure, gently insert the enema into the rectum with care to prevent damage to the rectal wall. Squeeze to empty the contents. Keep the tube squeezed until it is removed from the rectum. After the contents have been emptied, remove the disposable tube and discard. A small amount of liquid may remain in the unit after use.

Positioning

- For best results, lay on left side with knees bent.
- Second position: Kneel, then lower head and chest forward until side of face resting on surface.

Other Information

- each mini-enema contains: 283mg sodium docusate
- Store at 25°C (77°F); excursions permitted to 15°C - 30°C (59° - 86°F).
- Protect from Freezing. [See USP Controlled Room Temperature.]
- Below 50% relative humidity is optimal.
- See box and tube crimp for Lot Number and Expiration Date.

Inactive ingredients

a medicinal soft soap base of Polyethylene Glycol and Glycerin.

PRINCIPAL DISPLAY PANEL - 30 Tube Box

E-VAC

Plus

MINI-ENEMA

Suavizante de Heces / Evacuante

zürich

medical

labs

30 x 5cc TUBOS DESECHABLES

E-VAC *Plus*

MINI-ENEMA

Suavizante de Heces / Evacuante



30 x 5cc TUBOS DESECHABLES

REGISTROS:

BOLIVIA: Ministerio de Salud Pública No.

COLOMBIA: Com. de Espc Farmacéuticas No.

COSTA RICA: Dept. de Control y Registros No.

ECUADOR: Dirección General de Sanidad No.

EL SALVADOR: Consejo Superior de Salud Pública, No.

USA National Drug Code No. 61357-179-01

GUATEMALA: Registro San. No

HAITI: Departement du Commerce et de L'Industries No. Reg.

HONDURAS: No.

PANAMA: Dir. de Fcias., Drogas y Alim. No.

PERU: Minist. de Salud Pub., y As. Soc., Aut. No.

REP. DOMINICANA: Registro Sanitario No.

NICARAGUA: Inspección General de Farmacia No.

UPC
85499400540

Modo de Empleo (continuación)

También se recomienda que uno de estos lubricantes deben aplicarse hasta el ano antes de insertar el enema. Con la presión constante, insertar suavemente el enema en el recto con cuidado para evitar daños a la pared rectal. Apriete para vaciar el contenido. Mantenga el tubo apretado hasta que se retira del recto. Después de que los contenidos se han vaciado, retire el tubo desechable y deseche. Una pequeña cantidad de líquido puede permanecer en la unidad después de su uso.

Posicionamiento:

- Para obtener los mejores resultados, coloque en el lado izquierdo con las rodillas dobladas.
- La segunda posición: Arrodillarse, entonces baje la cabeza y el pecho hacia delante hasta que un lado de la cara este apoyada en la superficie.

Otra información

- cada mini-enema contiene: docusato de sodio 283 mg
- Almacenar a 25 °C (77 °F), se permiten variaciones de 15 °C - 30 °C (59 ° - 86 ° F).
- Proteger de la congelación. [Ver USP controlado de temperatura ambiente.]
- A continuación 50% de humedad relativa es óptima.
- Vea la caja y el tubo para el número de lote y fecha de caducidad.

Ingredientes inactivos

una base de jabón medicinal suave de polietilenglicol y glicerina.

PC-240.00

30 x 5cc TUBOS DESECHABLES



Suavizante de Heces / Evacuante

MINI-ENEMA

E-VAC *Plus*

E-VAC *Plus*

MINI-ENEMA

Suavizante de Heces / Evacuante



30 x 5cc TUBOS DESECHABLES

Información del Medicamento

<i>Ingrediente activo (en cada tubo desechable 5cc)</i>	<i>Propósito</i>
Docusato 283 mg de sodio	Suavizante de Heces
Benzocaina 20mg	Analgésico tópico

Usos ■ Para el alivio del estreñimiento y / o ayuda a prevenir el estreñimiento
 ■ Ayuda a reducir el dolor y el daño rectal causado por heces duras o por el esfuerzo durante la defecación ■ generalmente produce evacuación intestinal en 2 a 15 minutos

Advertencias

Pregúntele a su médico antes de usar si usted tiene
 ■ apendicitis ■ síntomas de apendicitis (por ejemplo, náuseas, vómitos, dolor abdominal)
 ■ sangrado del recto ■ hemorragia sin diagnosticar ■ insuficiencia cardíaca congestiva
 ■ Presión arterial alta ■ impactación fecal ■ obstrucción intestinal
 ■ estado tomando otros medicamentos

Pregúntele a su médico o farmacéutico antes de usar si usted está tomando aceites minerales.

Suspenda el uso y pregúntele a un doctor si

■ sangrado del recto ■ deja de tener una evacuación intestinal después de usar este producto. Estos pueden indicar una enfermedad grave.

Si está embarazada o en periodo de lactancia, consulte a un profesional de la salud antes de usar. **Mantener fuera del alcance de los niños.** En caso de ingestión accidental o sobredosis, obtenga ayuda médica o contacte un Centro de Control de Intoxicación inmediatamente.

Modo de Empleo

E-VAC™ MINI-ENEMA y *E-VAC™ PLUS MINI-ENEMA* se utilizan como enema. Esto no es un supositorio. No lo tome por vía oral.

Al administrar el mini enema:

Desenrosque y retire la punta. Colocar el agua o unas pocas gotas de producto en el eje del tubo a lubricar antes de la inserción.

DRUG FACTS (TRANSLATION)

Drug Facts

<i>Active ingredient (in each 5cc disposable tube)</i>	<i>Purpose</i>
Docusate Sodium 283mg	Stool Softener Enema
Benzocaine 20mg	Topical Analgesic

Uses ■ For the relief of occasional constipation and/or aid in preventing constipation
 ■ Helps to reduce pain or rectal damage caused by hard stools or by straining during bowel movements ■ usually produces bowel movement in 2 to 15 minutes

Warnings

Ask a doctor before use if you have

■ appendicitis ■ symptoms of appendicitis (e.g., nausea, vomiting, abdominal pain)
 ■ bleeding from the rectum ■ undiagnosed bleeding ■ congestive heart failure
 ■ high blood pressure ■ fecal impaction ■ intestinal obstruction
 ■ been taking other prescriptions

Ask a doctor or pharmacist before use if you are taking mineral oils.

Stop use and ask a doctor if

■ bleeding from the rectum ■ you fail to have a bowel movement after using this product.

Drug Facts (continued)

Directions (continued)

lubricate prior to insertion. It is also recommended that one of these lubricants be applied to the anus before inserting the enema. With steady pressure, gently insert the enema into the rectum with care to prevent damage to the rectal wall. Squeeze to empty the contents. Keep the tube squeezed until it is removed from the rectum. After the contents have been emptied, remove the disposable tube and discard. A small amount of liquid may remain in the unit after use.

Positioning:

■ For best results, lay on left side with knees bent.
 ■ Second position: Kneel, then lower head and chest forward until side of face resting on surface.

Other Information

■ each mini-enema contains: 283mg sodium docusate
 ■ Store at 25°C (77°F); excursions permitted to 15°C - 30°C (59° - 86°F).
 ■ Protect from Freezing. [See USP Controlled Room Temperature.]
 ■ Below 50% relative humidity is optimal.
 ■ See box and tube crimp for Lot Number and Expiration Date.

These may indicate a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental ingestion or overdose, get medical help or contact a Poison Control Center right away.

Directions

E-VAC™ MINI-ENEMA & E-VAC™ PLUS MINI-ENEMA is to be used as an enema. **This is not a suppository.** Do not take orally.

Administering the mini enema.

Twist off and remove tip. Place water or a few drops of the product on the shaft of the tube to

Inactive ingredients

a medicinal soft soap base of Polyethylene Glycol and Glycerin.

E-VAC PLUS

docusate sodium and benzocaine enema

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:6 1357-179
Route of Administration	RECTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DOCUSATE SODIUM (UNII: F05Q2T2JA0) (DOCUSATE - UNII:M7P27195AG)	DOCUSATE SODIUM	283 mg in 5 mL
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	20 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
GLYCERIN (UNII: PDC6A3C0OX)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:6 1357-179-01	30 in 1 BOX		
1		5 mL in 1 TUBE, WITH APPLICATOR		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
EXPORT ONLY		01/01/2010	

Labeler - ZURICH MEDICAL LABS, LLC (071904097)

Establishment

Name	Address	ID/FEI	Business Operations
ZURICH MEDICAL LABS, LLC		071904097	MANUFACTURE(6 1357-179)