ACETAMINOPHEN- acetaminophen tablet Rapid Care, Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

RapidCare Pain Reliever

Active ingredient (in each tablet)

Acetaminophen 500mg

Pain reliever/fever reducer

USES

temporarily relieves minor aches and pains due to:

- the common cold
- headache
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash If a skin reaction occurs, stop use and seek medical help right away.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present These could be signs of a serious condition
- You may report side effects to 1-888-952-0050

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

do not take more than directed (see overdose warning)

Adults and children 12 years and over

- take 2 tablets every 6 hours while symptoms last
- do not take more than 8 tablets in 24 hours, unless directed by a doctor
- do not take for more than 10 days unless directed by a doctor


Children under 12 years of age

ask a doctor

Other information

store between 20°-25°C (68°-77°F)

corn starch, hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide





ACETAMINOPHEN

acetaminophen tablet

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:73659-013

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII: 36209ITL9D) ACETAMINOPHEN 500 mg

Inactive Ingredients

l	Ingredient Name	Strength

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
HYPROMELLOSES (UNII: 3NXW29V3WO)

STARCH, CORN (UNII: O8232NY3SJ)
STEARIC ACID (UNII: 4ELV7Z65AP)

Product Characteristics					
Color	white	Score	score with uneven pieces		
Shape	ROUND	Size	12mm		
Flavor		Imprint Code			
Contains					

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73659-013- 01	50 in 1 BOX	04/18/2022	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:73659-013- 02	125 in 1 BOX	04/18/2022	
2		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information						
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
OTC monograph not final	part343	04/18/2022				

Labeler - Rapid Care, Inc (877432778)

Revised: 4/2022 Rapid Care, Inc