

**CIRCLE K HEADACHE RELIEF- acetaminophen, aspirin, and caffeine tablet**  
**Lil' Drug Store Products, Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Circle K™ Headache Relief**

***Drug Facts***

<b><i>Active ingredients (in each tablet)</i></b>	<b><i>Purpose</i></b>
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID*)	Pain reliever
Caffeine 65 mg	Pain reliever aid

\* nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - a cold
  - arthritis
  - muscular aches
  - sinusitis
  - toothache
  - premenstrual & menstrual cramps

**Warnings**

**Reye's syndrome**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

**Allergy alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### **Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Caffeine warning**

The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

### **Ask a doctor before use if**

- you have liver disease
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

### **Ask a doctor or pharmacist before use if you are taking**

- any other drug containing an NSAID (prescription or nonprescription)
- a blood thinning (anticoagulant) drug such as warfarin
- a steroid drug
- a prescription drug for diabetes, gout, or arthritis
- any other drug, or are under a doctor's care for any serious condition

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint

- vomit blood
- have bloody or black stools
- have stomach pain that does not get better.
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- **do not use more than directed**
- drink a full glass of water with each dose

adults and children 12 years and over	take 2 tablets every 6 hours; not more than 8 tablets in 24 hours.
children under 12 years	ask a doctor

### Other information

- store at room temperature 59-86°F (15-30°C)
- avoid excessive heat and humidity

### Inactive ingredients

hypromellose, maltodextrin, microcrystalline cellulose, polyethylene glycol, povidone, sodium starch glycolate, starch, stearic acid, titanium dioxide

### Questions or comments?

call toll-free **1-877-507-6516 (M-F 8 AM-4:30PM CST)**

**Proudly distributed by Circle K Stores Inc.**

### PRINCIPAL DISPLAY PANEL - 2 Tablet Pouch Carton

**Compare to the Active  
Ingredients in Excedrin®\***

**CIRCLE K™**

### Headache

**Acetaminophen, Aspirin (NSAID) and Caffeine  
Pain Reliever/Pain Reliever Aid**

**Extra-Strength**

**ACTUAL SIZE**

2 Tablets

1 POUCH, 2 TABLETS PER POUCH

Compare to the Active Ingredients in Excedrin®\*



# Headache

Acetaminophen, Aspirin (NSAID) and Caffeine  
Pain Reliever/Pain Reliever Aid



ACTUAL SIZE

Extra-Strength

2 Tablets

1 POUCH, 2 TABLETS PER POUCH

Headache 2 Tablets



DO NOT USE IF INNER POUCH OR CARTON IS OPEN OR DAMAGED!

### Drug Facts

Active ingredients (in each tablet)	Purpose
Acetaminophen 250 mg.....	Pain reliever
Aspirin 250 mg (NSAID)*.....	Pain reliever
Caffeine 65 mg.....	Pain reliever aid

\*nonsteroidal anti-inflammatory drug

**Uses** • temporarily relieves minor aches and pains due to:

- headache • a cold • arthritis • muscular aches
- sinusitis • toothache • premenstrual & menstrual cramps

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives • facial swelling • asthma (wheezing) • shock

### Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older • have had stomach ulcers or bleeding problems

LIFT PANEL FOR CONTINUED DRUG FACTS

### Drug Facts (continued)

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

**Do not use** • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer

• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if** • you have liver disease

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic • you have asthma

**Ask a doctor or pharmacist before use if you are taking**

- any other drug containing an NSAID (prescription or nonprescription)
- a blood thinning (anticoagulant) drug such as warfarin • a steroid drug
- a prescription drug for diabetes, gout, or arthritis
- any other drug, or are under a doctor's care for any serious condition

**When using this product** • take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint • vomit blood
  - have bloody or black stools • have stomach pain that does not get better
  - ringing in the ears or loss of hearing occurs • painful area is red or swollen

# Headache



Headache 2 Tablets

### Drug Facts (continued)

- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days • any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions • do not use more than directed

• drink a full glass of water with each dose

adults and children 12 years and over	take 2 tablets every 6 hours; not more than 8 tablets in 24 hours.
children under 12 years	ask a doctor

### Other information

• store at room temperature 59-86°F (15-30°C) • avoid excessive heat and humidity

### Inactive ingredients

hypromellose, maltodextrin, microcrystalline cellulose, polyethylene glycol, povidone, sodium starch glycolate, starch, stearic acid, titanium dioxide

### Questions or comments?

call toll-free 1-877-507-6516 (M-F 8 AM-4:30PM CST)

### Product manufactured for:

Lil' Drug Store Products, Inc., 1201 Continental Place NE, Cedar Rapids, IA 52402  
www.lildrugstore.com 57381C-US-02-16

Proudly distributed by Circle K Stores Inc.

\*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, LP, owners of the Excedrin® trademark.





# CIRCLE K HEADACHE RELIEF

acetaminophen, aspirin, and caffeine tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:66715-5738
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Acetaminophen</b> (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)	Acetaminophen	250 mg
<b>Aspirin</b> (UNII: R16CO5Y76E) (Aspirin - UNII:R16CO5Y76E)	Aspirin	250 mg
<b>Caffeine</b> (UNII: 3G6A5W338E) (Caffeine - UNII:3G6A5W338E)	Caffeine	65 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>Maltodextrin</b> (UNII: 7CVR7L4A2D)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>Sodium Starch Glycolate Type A Corn</b> (UNII: AG9B65PV6B)	
<b>Starch, Corn</b> (UNII: O8232NY3SJ)	
<b>Stearic Acid</b> (UNII: 4ELV7Z65AP)	
<b>Titanium Dioxide</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	11mm
<b>Flavor</b>		<b>Imprint Code</b>	FR9
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-5738-1	1 in 1 CARTON	11/01/2014	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part343	11/01/2014	

**Labeler** - Lil' Drug Store Products, Inc (093103646)

Revised: 3/2018

Lil' Drug Store Products, Inc