

**CARING MILL NIGHTTIME- acetaminophen, dextromethorphan hbr,
doxylamine succinate capsule, liquid filled
FSA Store Inc.**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

FSA Store Inc. Nighttime Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- minor aches and pains
- runny nose and sneezing
- sore throat
- headache
- fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening

- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- be careful when driving a motor vehicle or operating machinery
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- marked drowsiness may occur

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see overdose warning
- do not exceed 4 doses per 24 hours

adults & children 12 yrs & over	2 softgels with water every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Inactive ingredients

D&C yellow no. 10, edible ink*, FD&C blue no. 1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution *may contain this ingredient

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to Vicks[®] NyQuil[®] Cold & Flu active ingredients

Nighttime

Pain Reliever, Fever Reducer, Cough Suppressant, Antihistamine

Acetaminophen

Dextromethorphan HBr

Doxylamine succinate

Cold & Flu

Night Time Relief

16 Softgels

Actual Size

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org



NDC 81522-056-73



caring mill™

**Compare to Vicks® NyQuil® Cold & Flu active ingredients

Nighttime

Pain Reliever, Fever Reducer,
Cough Suppressant, Antihistamine

Acetaminophen
Dextromethorphan HBr
Doxylamine succinate

Cold & Flu
Night Time Relief

16 Softgels



Actual Size

CONVENIENT RECYCLING TAB

Drug Facts

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

Active ingredients (in each softgel)

Purpose

Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 15 mg.....	Cough suppressant
Doxylamine succinate 6.25 mg.....	Antihistamine

Uses temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- minor aches and pains
- runny nose and sneezing
- sore throat
- headache
- fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription).

If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have ■ liver disease ■ glaucoma

- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product ■ excitability may occur, especially in children

- be careful when driving a motor vehicle or operating machinery
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- marked drowsiness may occur

Stop use and ask a doctor if ■ pain or cough gets worse or lasts more than 7 days

- redness or swelling is present
- fever gets worse or lasts more than 3 days
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. ▶

OPEN OTHER END

05673 JV C1

Drug Facts (continued)

Directions ■ take only as directed - see overdose warning

- do not exceed 4 doses per 24 hours

adults & children 12 yrs & over	2 softgels with water every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Inactive ingredients D&C yellow no. 10, edible ink*, FD&C blue no. 1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution *may contain this ingredient ▶

Drug Facts (continued)

Questions or comments? 1-800-719-9260

Store at 20-25°C (68-77°F).

**This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® NyQuil® Cold & Flu.

Manufactured for **FSA Store Inc.**
240 W. 37th St 6th Floor, New York, NY 10018
Customer Care Help Line 1-800-719-9260

Made in Canada

CARING MILL NIGHTTIME

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:81522-056
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	GREEN (clear)	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	056
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:81522-056-73	8 in 1 CARTON	09/03/2021	
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/03/2021	

Labeler - FSA Store Inc. (049283340)

Revised: 9/2021

FSA Store Inc.