# ADVIL- ibuprofen tablets, coated tablet, coated Lil' Drug Store Products, Inc.

-----

Advil ®

### **Active Ingredient**

### Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

### **Purpose**

Pain reliever/Fever reducer

#### Uses

- temporarily relieves minor aches and pains due to:
- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

### Warnings Section

### Warnings

# **Allergy Alert**

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

# Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance

is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### Heart attack and stroke warning:

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

#### Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

#### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

# Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

# When using this product

take with food or milk if stomach upset occurs

# Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days

- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

#### Other information

- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

# Inactive ingredients

acetylated monoglycerides, colloidal silicon dioxide, corn starch, croscarmellose sodium, methylparaben, microcrystalline cellulose,

pharmaceutical glaze, pharmaceutical ink, povidone, pregelatinized starch, propylparaben, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, synthetic iron oxide, titanium dioxide, white wax

#### **Questions or comments?**

call toll free 1-800-88-ADVIL

For most recent product information, visit www.Advil.com

Dist. by GSK Consumer Healthcare, Warren, NJ 07059 ©2020 GSK or licensor

### **Primary Display Panel**

Advil ®

Ibuprofen Tablets, 200 mg

Pain Reliever/Fever Reducer (NSAID)

[tablet image]


**Tablets** 

### **30 PACKETS OF 2 TABLETS**

www.lildrugstore.com

[UPC]

3 66715 58030 6





of stomach problems, such as hearthurn wou have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke wou are taking a diuretic. Ask a doctor or pharmacist before use if you are wonder a doctor's care for any serious condition. It alking any other drug. When using this product lack evith load or milk if stomach upset occurs. Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: lefel faint womit blood have bloody or black stools. I have shoody or black stools in the pain to the stools because it may not have symptoms or sometimes of the swelling when the stools have some some shoody or well as the stools have shoody or well as the stool of the stools have some shood or seal as the stool of the stool of

L7451A EXP 11/23

#### **ADVIL**

ibuprofen tablets, coated tablet, coated

#### **Product Information**

NDC:66715-5803 **Product Type** HUMAN OTC DRUG **Item Code (Source)** 

**ORAL Route of Administration** 

### **Active Ingredient/Active Moiety**

**Basis of Strength** Strength **Ingredient Name** IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII: WK2XYI10QM) **IBUPROFEN** 200 mg

#### **Inactive Ingredients Ingredient Name** Strength CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)

**TITANIUM DIOXIDE** (UNII: 15FIX9V2JP)

STARCH, CORN (UNII: O8232NY3SJ)

**DIACETYLATED MONOGLYCERIDES (UNII: 5Z17386USF)** 

**SODIUM LAURYL SULFATE (UNII: 368GB5141))** 

STEARIC ACID (UNII: 4ELV7Z65AP)

SUCROSE (UNII: C151H8M554) WHITE WAX (UNII: 7G1J5DA97F)

**SODIUM BENZOATE** (UNII: OJ245FE5EU)

METHYLPARABEN (UNII: A2I8C7HI9T)

**SHELLAC** (UNII: 46N107B710)

POVIDONE, UNSPECIFIED (UNII: FZ 989GH94E)

FERRIC OXIDE RED (UNII: 1K09F3G675)

CROSCARMELLOSE SODIUM (UNII: M280L1HH48)

PROPYLPARABEN (UNII: Z8IX2SC10H) **SILICON DIOXIDE** (UNII: ETJ7Z6XBU4)

Product Characteristics						
Color	brown ((pinkish brown))	Score	no score			
Shape	ROUND	Size	11mm			
Flavor		Imprint Code	Advil			
Contains						

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:66715- 5803-0	30 in 1 BOX, UNIT-DOSE	11/16/2006			
1		2 in 1 POUCH; Type 0: Not a Combination Product				

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
NDA	NDA018989	11/16/2006			

Labeler - Lil' Drug Store Products, Inc. (093103646)

Revised: 4/2022 Lil' Drug Store Products, Inc.