

COLD FLU AND SORE THROAT RELIEF- acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride capsule, liquid filled
PuraCap Pharmaceutical LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

COLD, FLU & SORE THROAT RELIEF

Drug Facts

Active Ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

- temporarily relieves:
 - nasal congestion
 - headache
 - cough
 - minor aches and pains
 - sore throat
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 softgels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening

- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product, do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 liquid gels in any 24-hour

period

- adults and children 12 years of age and older: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

Other information

- store between 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients

FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, and white edible ink

Questions or Comments?

☐Call toll free: **1-855-215-8180**

PRINCIPAL DISPLAY PANEL

COLD, FLU & SORE THROAT RELIEF 16 SOFTGELS

NDC 51013-191-14

*Compare to the active ingredients in Mucinex® Fast-Max® Cold, Flu & Sore Throat



MAXIMUM STRENGTH*

Cold, Flu & Sore Throat

THIS PRODUCT IS PACKAGED IN A CHILD RESISTANT AND TAMPER EVIDENT PACKAGE. USE ONLY IF BLISTERS ARE INTACT.

NDC 51013-191-14



MAXIMUM STRENGTH*

Cold, Flu & Sore Throat

Acetaminophen – Pain Reliever/Fever Reducer
Dextromethorphan HBr – Cough Suppressant
Guaifenesin – Expectorant
Phenylephrine HCl – Nasal Decongestant

- Controls Cough, Thins & Loosens Mucus
- Relieves Nasal & Chest Congestion
- Relieves Headache & Fever



** Compare to the active ingredients in Mucinex® Fast-Max® Cold, Flu & Sore Throat

16 SOFTGELS

Non-Drowsy
For Ages 12+

MAXIMUM STRENGTH*
Cold, Flu & Sore Throat



*Per 4-hour dose.
Do not take more than a total of 12 softgels in a 24-hour period.



PLASTE RECYCLE

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

LOT NO:

EXP DATE:



0 93351 02398 9

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

www.discount-drugmart.com
DISTRIBUTED BY
DRUG MART-FOOD FAIR
MEDINA, OHIO 44256
MADE IN CHINA

SATISFACTION GUARANTEED
IF DISSATISFIED, RETURN UNUSED
PORTION AND PACKAGE TO STORE
WHERE PURCHASED. IF UNABLE TO
RETURN TO STORE, SEND REASON FOR
DISSATISFACTION, NAME, ADDRESS AND
EMPTY PACKAGE TO: DISCOUNT DRUG
MART, 211 COMMERCE DRIVE, MEDINA,
OHIO 44256

DDM33-00

Drug Facts

Active Ingredients (in each softgel)	Purposes
Acetaminophen 325 mg.....Pain reliever/fever reducer	
Dextromethorphan HBr 10 mg.....Cough suppressant	
Guaifenesin 200 mg.....Expectorant	
Phenylephrine HCl 5 mg.....Nasal decongestant	

Uses

- temporarily relieves these common cold and flu symptoms:
 - headache
 - nasal congestion
 - minor aches and pains
 - cough
 - sore throat
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 softgels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease

Drug Facts (continued)

- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

if pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 softgels in any 24-hour period
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use

Other information

- store between 15-30°C (59-86°F)
- avoid excessive heat

Inactive ingredients
FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution and white edible ink

Questions or comments?
Call toll free: 1-800-833-6278

**This product is not manufactured or distributed by Reckitt Benckiser, owner of the registered trademarks Mucinex® Fast-Max® Cold, Flu & Sore Throat.

COLD FLU AND SORE THROAT RELIEF			
acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride capsule, liquid filled			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:510 13-19 1
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	

GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	orange (clear)	Score	no score
Shape	capsule (oblong)	Size	25mm
Flavor		Imprint Code	PC26
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51013-191-14	2 in 1 CARTON	07/14/2017	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/14/2017	

Labeler - PuraCap Pharmaceutical LLC (962106329)

Establishment

Name	Address	ID/FEI	Business Operations
Humanwell PuraCap Pharmaceutical (Wuhan) Co., Ltd		421293287	manufacture(51013-191) , analysis(51013-191)