DOXEPIN HYDROCHLORIDE- doxepin hydrochloride cream Amneal Pharmaceuticals NY LLC

Doxepin Hydrochloride Cream, 5%

DESCRIPTION

Doxepin hydrochloride (HCl) cream, 5% is a topical cream. Each gram contains: 50 mg of doxepin hydrochloride, USP (equivalent to 44.3 mg of doxepin).

Doxepin hydrochloride, USP is one of a class of agents known as dibenzoxepin tricyclic antidepressant compounds. It is an isomeric mixture of N,N-dimethyldibenz[*b,e*]oxepin- $\Delta^{11(6H),\gamma}$ - propylaminehydrochloride. Doxepin hydrochloride, USP has an empirical formula of C₁₉H₂₁NO•HCl and a molecular weight of 315.84.



Doxepin HCl cream, 5% also contains sorbitol solution, cetyl alcohol, isopropyl myristate, glyceryl monostearate, PEG-100 stearate, white petrolatum, benzyl alcohol, titanium dioxide and purified water.

CLINICAL PHARMACOLOGY

Although doxepin HCl does have H₁ and H₂ histamine receptor blocking actions, the exact mechanism by which doxepin exerts its antipruritic effect is unknown. Doxepin HCl cream can produce drowsiness in significant numbers of patients, and this sedation may reduce awareness, including awareness of pruritic symptoms. In 19 pruritic eczema patients treated with doxepin HCl cream, plasma doxepin concentrations ranged from nondetectable to 47 ng/mL from percutaneous absorption. Plasma levels from topical application of doxepin HCl cream can result in CNS and other systemic side effects. Once absorbed into the systemic circulation, doxepin undergoes hepatic metabolism that results in conversion to pharmacologically-active desmethyldoxepin. Further glucuronidation results in urinary excretion of the parent drug and its metabolites. Desmethyldoxepin has a half-life that ranges from 28 to 52 hours and is not affected by multiple dosing. Plasma levels of both doxepin and desmethyldoxepin are highly variable and are poorly correlated with dosage. Wide distribution occurs in

body tissues including lungs, heart, brain, and liver. Renal disease, genetic factors, age, and other medications affect the metabolism and subsequent elimination of doxepin (see **PRECAUTIONS - DRUG INTERACTIONS**).

INDICATIONS AND USAGE

Doxepin HCl cream, 5% is indicated for the short-term (up to 8 days) management of moderate pruritus in adult patients with atopic dermatitis or lichen simplex chronicus (see **DOSAGE AND ADMINISTRATION**).

CONTRAINDICATIONS

Because doxepin HCl has an anticholinergic effect and because significant plasma levels of doxepin are detectable after topical doxepin HCl cream application, the use of doxepin HCl cream is contraindicated in patients with untreated narrow angle glaucoma or a tendency to urinary retention.

Doxepin HCl cream is contraindicated in individuals who have shown previous sensitivity to any of its components.

WARNINGS

Drowsiness occurs in over 20% of patients treated with doxepin HCl cream, especially in patients receiving treatment to greater than 10% of their body surface area. **Patients should be warned about the possibility of sedation and cautioned against driving a motor vehicle or operating hazardous machinery while being treated with doxepin HCl cream.**

The sedating effects of alcoholic beverages, antihistamines, and other CNS depressants may be potentiated when doxepin HCl cream is used.

If excessive drowsiness occurs it may be necessary to reduce the frequency of applications, the amount of cream applied, and/or the percentage of body surface area treated, or discontinue the drug. However, the efficacy with reduced frequency of applications has not been established. Keep this product away from the eyes.

PRECAUTIONS

General

Drowsiness

Since drowsiness may occur with the use of doxepin HCl cream, patients should be warned of the possibility and cautioned against driving a car or operating dangerous machinery while using this drug. Patients should also be cautioned that their response to alcohol may be potentiated.

Sedating drugs may cause confusion and oversedation in the elderly; elderly patients generally should be observed closely for confusion and oversedation when started on doxepin HCl cream (see **PRECAUTIONS - Geriatric Use**).

Use under occlusion

Occlusive dressings may increase the absorption of most topical drugs; therefore, occlusive dressings should not be utilized with doxepin HCl cream.

Contact sensitization

Use of doxepin HCl cream can cause Type IV hypersensitivity reactions (contact sensitization) to doxepin.

DRUG INTERACTIONS

Studies have not been performed examining drug interactions with doxepin HCl cream. However, since plasma levels of doxepin following topical application of doxepin HCl cream can reach levels obtained with oral doxepin HCl therapy, the following drug interactions are possible following topical doxepin HCl cream application:

Drugs Metabolized by P450 2D6

The biochemical activity of the drug metabolizing isozyme cytochrome P450 2D6 (debrisoquin hydroxylase) is reduced in a subset of the Caucasian population (about 7% to 10% of Caucasians are so-called "poor metabolizers"); reliable estimates of the prevalence of reduced P450 2D6 isozyme activity among Asian, African and other populations are not yet available. Poor metabolizers have higher than expected plasma concentrations of tricyclic antidepressants (TCAs) when given usual doses. Depending on the fraction of drug metabolized by P450 2D6, the increase in plasma concentration may be small, or quite large (8-fold increase in plasma AUC of the TCA).

In addition, certain drugs inhibit the activity of this isozyme and make normal metabolizers resemble poor metabolizers. An individual who is stable on a given dosage regimen of a TCA may become abruptly toxic when given one of these inhibiting drugs as concomitant therapy. The drugs that inhibit cytochrome P450 2D6 include some that are not metabolized by the enzyme (quinidine; cimetidine) and many that are substrates for P450 2D6 (many other antidepressants, phenothiazines, and the Type 1C antiarrhythmics propafenone and flecainide). While all the selective serotonin reuptake inhibitors (SSRIs), e.g., fluoxetine, sertraline, and paroxetine, inhibit P450 2D6, they may vary in the extent of inhibition. The extent to which SSRI-TCA interactions may pose clinical problems will depend on the degree of inhibition and the pharmacokinetics of the SSRI involved. Nevertheless, caution is indicated in the co-administration of TCAs with any of the SSRIs. Of particular importance, sufficient time must elapse before initiating TCA treatment in a patient being withdrawn from fluoxetine, given the long half-life of the parent and active metabolite (at least 5 weeks may be necessary).

Concomitant use of tricyclic antidepressants with drugs that can inhibit cytochrome P450 2D6 may require lower doses than usually prescribed for either the tricyclic antidepressant or the other drug. It is desirable to monitor TCA plasma levels whenever a TCA is going to be co-administered with another drug known to be an inhibitor of P450 2D6.

MAO Inhibitors

Serious side effects and even death have been reported following the concomitant use of certain drugs with MAO inhibitors. Therefore, MAO inhibitors should be discontinued at least two weeks prior to the cautious initiation of therapy with doxepin HCl cream. The exact length of time may vary and is dependent upon the particular MAO inhibitor being used, the length of time it has been administered, and the dosage involved.

Cimetidine

Serious anticholinergic symptoms (i.e., severe dry mouth, urinary retention and blurred vision) have been associated with elevations in the serum levels of tricyclic antidepressant when cimetidine therapy is initiated. Additionally, higher than expected tricyclic antidepressant levels have been observed when they are begun in patients already taking cimetidine.

Alcohol

Alcohol ingestion may exacerbate the potential sedative effects of doxepin HCl cream. This is especially important in patients who may use alcohol excessively.

Tolazamide

A case of severe hypoglycemia has been reported in a type II diabetic patient maintained on tolazamide (1 gm/day) 11 days after the addition of oral doxepin (75 mg/day).

Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis, mutagenesis, and impairment of fertility studies have not been conducted with doxepin hydrochloride.

Reproduction studies have been performed in which doxepin was orally administered to rats and rabbits at doses up to 0.6 and 1.2 times, respectively, the estimated exposure to doxepin that results from use of 16 grams of doxepin HCl cream per day (four applications of four grams of cream per day; dose multiples reflect comparisons made following normalization of the data on the basis of body surface area estimates) and have revealed no evidence of harm to rat or rabbit fetuses due to doxepin. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers

Doxepin is excreted in human milk after oral administration. It is possible that doxepin may also be excreted in human milk following topical application of doxepin HCl cream.

One case has been reported of apnea and drowsiness in a nursing infant whose mother was taking an oral dosage form of doxepin HCl.

Because of the potential for serious adverse reactions in nursing infants from doxepin, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use

The use of doxepin HCl cream in pediatric patients is not recommended. Safe conditions for use of doxepin HCl cream in children have not been established. One case has been reported of a 2.5 year old child who developed somnolence, grand mal seizure, respiratory depression, ECG abnormalities, and coma after treatment with doxepin HCl cream. A total of 27 grams had been applied over three days for eczema. He was treated with supportive care, activated charcoal, and systemic alkalization and recovered.

Geriatric Use

Clinical studies of doxepin HCl cream did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal or cardiac function, and of concomitant disease or other drug therapy.

The extent of renal excretion of doxepin has not been determined. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selections.

Sedating drugs may cause confusion and oversedation in the elderly; elderly patients generally should be observed closely for confusion and oversedation when started on doxepin HCl cream (see **WARNINGS**). An 80-year old male nursing home patient developed probable systemic anticholinergic toxicity which included urinary retention and delirium after doxepin HCl cream had been applied to his arms, legs and back three times daily for two days.

ADVERSE REACTIONS

Controlled Clinical Trials

Systemic Adverse Effects

In controlled clinical trials of patients treated with doxepin HCl cream, the most common systemic adverse event reported was drowsiness. Drowsiness occurred in 71 of 330 (22%) of patients treated with doxepin HCl cream compared to 7 of 334 (2%) of patients treated with vehicle cream. Drowsiness resulted in the premature discontinuation of the drug in approximately 5% of patients treated with doxepin HCl cream in controlled clinical trials.

Local Site Adverse Effects

In controlled clinical trials of patients treated with doxepin HCl cream, the most common local site adverse event reported was burning and/or stinging at the site of application. These occurred in 76 of 330 (23%) of patients treated with doxepin HCl cream compared to 54 of 334 (16%) of patients treated with vehicle cream. Most of these reactions were categorized as "mild"; however, approximately 25% of patients who reported burning and/or stinging reported the reaction as "severe". Four patients treated with doxepin HCl cream withdrew from the study because of the burning and/or stinging.

The table below presents the adverse events reported at an incidence of $\geq 1\%$ in either doxepin HCl cream or vehicle cream treatment groups during the trials:

Adverse Event	Doxepin HCl Cream N=330	Vehicle N=334	
Burning /Stinging	76 (23.0%)	54 (16.2%)	
Drowsiness	71 (21.5%)	7 (2.1%)	

Dry Mouth ¹	32 (9.7%)	4 (1.2%)		
Pruritus ²	13 (3.9%)	20 (6.0%)		
Fatigue/Tiredness	10 (3.0%)	5 (1.5%)		
Exacerbated Eczema	10 (3.0%)	8 (2.4%)		
Other Application Site Reaction ³	10 (3.0%)	16 (4.8%)		
Dizziness ⁴	7 (2.1%)	3 (0.9%)		
Mental Emotional Changes	6 (1.8%)	1 (0.3%)		
Taste Perversion ⁵	5 (1.5%)	1 (0.3%)		
Edema	4 (1.2%)	1 (0.3%)		
Headache	3 (0.9%)	14 (4.2%)		
¹ Includes reports of "dry lips", "dry throat", and "thirst"				
² Includes reports of "Pruritus Exacerbated"				
³ Includes report of "increased irritation at application site"				
⁴ Includes reports of "lightheadedness" and "dizziness/vertigo"				
⁵ Includes reports of "bitter taste" and "metallic taste in mouth"				

Adverse events occurring in 0.5% to < 1.0% of doxepin HCl cream treated patients in the controlled clinical trials included: nervousness/anxiety, tongue numbness, fever, and nausea.

Post Marketing Experience

Twenty-six cases of allergic contact dermatitis have been reported in patients using doxepin HCl cream, twenty of which were documented by positive patch test to doxepin 5% cream.

To report SUSPECTED ADVERSE REACTIONS, contact Amneal Pharmaceuticals at 1-877-835-5472 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

OVERDOSAGE

Deaths may occur from overdosage with this class of drugs. As the management is complex and changing, it is recommended that the physician contact a poison control center for current information on treatment. Signs and symptoms of toxicity develop rapidly after tricyclic antidepressant overdose; therefore, hospital monitoring is required as soon as possible.

Manifestations

Should overdosage with topical application of doxepin HCl cream occur, the signs and symptoms may include: cardiac dysrhythmias, severe hypotension, convulsions, and CNS depression, including coma. Changes in the electrocardiogram, particularly in QRS axis or width, are clinically significant indicators of tricyclic antidepressant toxicity.

Other signs of overdose may include: confusion, disturbed concentration, transient visual hallucinations, dilated pupils, agitation, hyperactive reflexes, stupor, drowsiness, muscle rigidity, vomiting, hypothermia, hyperpyrexia, or any of the symptoms listed under **ADVERSE REACTIONS**.

General Recommendations

General

Obtain an ECG and immediately initiate cardiac monitoring. Protect the patient's airway, establish an intravenous line and initiate gastric decontamination. A minimum of six hours of observation with cardiac monitoring and observation for signs of CNS or respiratory depression, hypotension, cardiac dysrhythmias and/or conduction blocks, and seizures is strongly advised. If signs of toxicity occur at any time during this period, extended monitoring is recommended. There are case reports of patients succumbing to fatal dysrhythmias late after overdose; these patients had clinical evidence of significant poisoning prior to death and most received inadequate gastrointestinal decontamination. Monitoring of plasma drug levels should not guide management of the patient.

Cardiovascular

A maximal limb-lead QRS duration of ≥ 0.10 seconds may be the best indication of the severity of the overdose. Intravenous sodium bicarbonate should be used to maintain the serum pH in the range of 7.45 to 7.55. If the pH response is inadequate, hyperventilation may also be used. Concomitant use of hyperventilation and sodium bicarbonate should be done with extreme caution, with frequent pH monitoring. A pH>7.60 or a pCO₂ <20 mm Hg is undesirable. Dysrhythmias unresponsive to sodium bicarbonate therapy/hyperventilation may respond to lidocaine, bretylium or phenytoin. Type 1A and 1C antiarrhythmics are generally contraindicated (e.g., quinidine, disopyramide, and procainamide).

In rare instances, hemoperfusion may be beneficial in acute refractory cardiovascular instability in patients with acute toxicity. However, hemodialysis, peritoneal dialysis, exchange transfusions, and forced diuresis generally have been reported as ineffective in tricyclic antidepressant poisoning.

CNS

In patients with CNS depression, early intubation is advised because of the potential for abrupt deterioration. Seizures should be controlled with benzodiazepines, or if these are ineffective, other anticonvulsants (e.g., phenobarbital, phenytoin). Physostigmine is not recommended except to treat life-threatening symptoms that have been unresponsive to other therapies, and then only in consultation with a poison control center.

Pediatric Management

The principles of management of child and adult overdosages are similar. It is strongly recommended that the physician contact the local poison control center for specific pediatric treatment.

DOSAGE AND ADMINISTRATION

A thin film of doxepin HCl cream, 5% should be applied four times each day with at least a 3 to 4 hour interval between applications. There are no data to establish the safety and effectiveness of doxepin HCl cream, 5% when used for greater than 8 days. Chronic use beyond eight days may result in higher systemic levels and should be avoided. Use of doxepin HCl cream, 5% for longer than 8 days may result in an increased likelihood of contact sensitization. The risk for sedation may increase with greater body surface area application of doxepin HCl cream, 5% (See **WARNINGS** section). Clinical experience has shown that drowsiness is significantly more common in patients applying doxepin HCl cream, 5% to over 10% of body surface area; therefore, patients with greater than 10% of body surface area (see **WARNINGS** section) affected should be particularly cautioned concerning possible drowsiness and other systemic adverse effects of doxepin. If excessive drowsiness occurs, it may be necessary to do one or more of the following: reduce the body surface area treated, reduce the number of applications per day, reduce the amount of cream applied, or discontinue the drug.

Occlusive dressings may increase the absorption of most topical drugs; therefore, occlusive dressings should not be utilized with doxepin HCl cream, 5%.

HOW SUPPLIED

Doxepin HCl cream, 5% is available as a soft homogeneous white to off-white cream free from lumps and foreign particles. It is supplied as follows:

45 g tubes (NDC 69238-1733-6)

Store at or below 27° C (80° F).

Distributed by: **Amneal Pharmaceuticals LLC** Bridgewater, NJ 08807

Rev. 02-2022-00

PRINCIPAL DISPLAY PANEL

eal'	NDC 69238- 1733 -6 Doxepin Hydrocl	hloride	
amn	Cream, 5 % FOR TOPICAL DERMATOLOGIC USE ONLY		
4	Rx only	NET WT 45 g	verage (no text)
Warnings Keep this Directions between a surface are	For External Use Only. Not For Oral, Intravaginal or O and all drugs out of the reach of children. Apply a thin film of doxepin hydrochloride cream, 5% four oplications. If excessive drowsiness occurs it may be necess a treated, reduce the number of applications per day, reduce	Dephthalmic Use. Keep away from eyes. times each day with at least a 3 to 4 hour interval any to do one or more of the following: reduce the body e the amount of cream applied, or discontinue the drug.	Full Co
Refer to pr Store at o	oduct insert for prescribing information and inactive ingredi r Below 27°C (80°F).	ent listing.	
See crimp Distributed Amneal P Bridgewate Rev. 02-20	for lot number and expiration date. by: harmaceuticals LLC r, NJ 08807 /22-01	N 69238 17336 9	



DOXEPIN HYDROCHLORIDE doxepin hydrochloride cream					
Product Information					
Product Type	HUMAN PRESCRIPTION DRUG Item Code (Source)			NDC:69238-1733	
Route of Administration	TOPICAL				
Active Ingredient/Active	Moiety				
Ingredient Name Basis of Str					Strength
DOXEPIN HYDROCHLORIDE (UNII: 3U9A0FE9N5) (DOXEPIN - UNII:5ASJ6HUZ7D) DOXEPIN			•	50 mg in 1 g	
Inactivo Ingradiante					
mactive ingredients					
Ingredient Name				St	trength
SORBITOL (UNII: 506T60A25R)					
CETYL ALCOHOL (UNII: 936JST6JCN)					
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)					
GLYCERYL MONOSTEARATE (UNII: 2300U9XXE4)					
POLYOXYL 100 STEARATE (UNII: YD01N1999R)					
PETROLATUM (UNII: 4T6H12BN9U)					
BENZYL ALCOHOL (UNII: LKG8494WBH)					
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)					

WATER (UNII: 059QF0KO0R)							
P	Product Characteristics						
Сс	Color white (white to off-white) Score						
Sł	Shape Size		Size				
Fla	lavor Imprint Code		Imprint Code				
Сс	Contains						
D	ackaging						
			Man	Lating Ctart	Mayleating		
#	Item Code	Package Description	Mar	keting Start Date	Marketing Date	End	
# 1	Item Code NDC:69238- 1733-6	Package Description 1 in 1 CARTON	Mar 08/21/2	keting Start Date	Marketing Date	End	
# 1 1	Item Code NDC:69238- 1733-6	Package Description 1 in 1 CARTON 45 g in 1 TUBE; Type 0: Not a Combination Product	Mar 08/21/2	keting Start Date 023	Marketing Date	End	
# 1 1	Item Code NDC:69238- 1733-6	Package Description 1 in 1 CARTON 45 g in 1 TUBE; Type 0: Not a Combination Product	Mar	keting Start Date 023	Marketing Date	End	
# 1 1	Item Code NDC:69238- 1733-6	Package Description 1 in 1 CARTON 45 g in 1 TUBE; Type 0: Not a Combination Product	Mar	keting Start Date 023	Marketing Date	End	
# 1 1	Item Code	Package Description 1 in 1 CARTON 45 g in 1 TUBE; Type 0: Not a Combination Product nformation	Mar	keting Start Date	Marketing Date	End	
# 1 1	Item Code NDC:69238- 1733-6	Package Description 1 in 1 CARTON 45 g in 1 TUBE; Type 0: Not a Combination Product Image: Colspan="2">Image: Colspan="2" Colspan="2">Image: Colspan="2" Colspa	Mar 08/21/2	keting Start Date 023 arketing Start Date	Marketing Date Marketing Date	End J End	
# 1 1 AN	Item Code NDC:69238- 1733-6 Iarketing I Marketing Category DA	Package Description 1 in 1 CARTON 45 g in 1 TUBE; Type 0: Not a Combination Product Product	Mar 08/21/2 M 08/2	keting Start Date 023 arketing Start Date 1/2023	Marketing Date Marketing Date	End J End	

Labeler - Amneal Pharmaceuticals NY LLC (123797875)

Establishment				
Name	Address	ID/FEI	Business Operations	
Amneal Pharmaceuticals, LLC		079389286	analysis(69238-1733) , label(69238-1733) , manufacture(69238-1733) , pack(69238-1733)	

Revised: 8/2023

Amneal Pharmaceuticals NY LLC