# CORICIDIN HBP MAXIMUM STRENGTH COLD, COUGH AND FLUacetaminophen, dextromethorphan hydrobromide liquid Bayer HealthCare LLC.

-----

# Coricidin HBP Maximum Strength Cold , Cough & Flu Liquid

# **Drug Facts**

Active ingredients (in each 30 mL)	Purposes
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan hydrobromide 20 mg	Cough suppressant

### Uses

- temporarily relieves these symptoms due to a cold or flu:
  - minor aches and pains
  - headache
  - cough
  - sore throat
- temporarily reduces fever

## Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

 with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

## Ask a doctor before use if you have

- liver disease
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that is accompanied with excessive phlegm (mucus)
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

# Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs
  of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## **Directions**


- do not take more than recommended dose
- adults and children 12 years and over: take 30 mL in the dosing cup provided every 4 hours. Do not exceed 150 mL (5 doses) in 24 hours or as directed by a doctor.
- children under 12 years of age: do not use

### Other information


- each 30 mL contains: sodium 38 mg
- store between 20° to 25°C (68° to 77°F)

**Inactive ingredient**s anhydrous citric acid, FD&C blue #1, FD&C red # 40, flavor, glycerin, maltitol solution, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

**Questions or comments? 1-800-317-2165** (Mon – Fri 9AM – 5PM EST)







Approved by eoknj on 07-Feb-2022 15:09

New Formula!

Coricidin®

**HBP** 

COLD RELIEF FOR PEOPLE WITH

HIGH BLOOD PRESSURE†

MAXIMUM STRENGTH

**NIGHTTIME** 

**COLD** 

& FLU

Relieves

**Body Aches & Pains** 

Headache, Fever,

Cough, Sore Throat,

**SUGAR FREE** 

Alcohol Free

Cherry

†Decongestant Free

# **Acetaminophen -**

Pain Reliever/Fever Reducer

Dextromethorphan HBr-

Cough Suppressant

12 FL OZ (355 mL)

# CORICIDIN HBP MAXIMUM STRENGTH COLD, COUGH AND FLU

acetaminophen, dextromethorphan hydrobromide liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11523-0093
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	650 mg in 30 mL	
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL	

Inactive Ingredients		
Ingredient Name	Strength	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
WATER (UNII: 059QF0KO0R)		
SACCHARIN SODIUM (UNII: SB8ZUX40TY)		
SODIUM CITRATE (UNII: 1Q73Q2JULR)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		

ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
MALTITOL (UNII: D65DG142WK)	

l	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:11523- 0093-1	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	04/27/2022	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/27/2022	

Labeler - Bayer HealthCare LLC. (112117283)

Revised: 11/2023 Bayer HealthCare LLC.