


HYDROCORTISONE ACETATE - hydrocortisone acetate suppository
Rising Pharma Holdings, Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Hydrocortisone Acetate Suppositories, 25 mg
For Rectal Administration
Rx only

DESCRIPTION

Hydrocortisone acetate is a corticosteroid designed chemically as pregn-4-ene-3, 20-dione, 21-(acetyloxy)-11, 17-dihydroxy-(11 β) with the following structural formula:



Each suppository for rectal administration contains hydrocortisone acetate, USP 25 mg in a specially blended hydrogenated vegetable base.

CLINICAL PHARMACOLOGY

In normal subjects, about 26% of hydrocortisone acetate is absorbed when the hydrocortisone acetate suppository is applied to the rectum. Absorption of hydrocortisone acetate may vary across abraded or inflamed surfaces.

Topical steroids are primarily effective because of their anti-inflammatory, anti-pruritic and vasoconstrictive action.

INDICATIONS AND USAGE

Hydrocortisone acetate suppositories are indicated for use in inflamed hemorrhoids, post-irradiation (factitial) proctitis, as an adjunct in the treatment of chronic ulcerative colitis, cryptitis, other inflammatory conditions of anorectum, and pruritus ani.

CONTRAINDICATIONS

Hydrocortisone acetate suppositories are contraindicated in those patients having a history of hypersensitivity to hydrocortisone acetate or any of the components.

PRECAUTIONS

Do not use hydrocortisone acetate suppositories unless adequate proctologic examination is made.

If irritation develops, the product should be discontinued and appropriate therapy instituted.

In the presence of an infection, the use of an appropriate antifungal or antibacterial agent should be instituted. If a favorable response does not occur promptly, hydrocortisone acetate should be discontinued until the infection has been adequately controlled.

Carcinogenesis

No long-term studies in animals have been performed to evaluate the carcinogenic potential of corticosteroid suppositories.

INFORMATION FOR PATIENTS

Staining of fabric may occur with use of the suppository. Precautionary measures are recommended.

PREGNANCY CATEGORY C

In laboratory animals, topical steroids have been associated with an increase in the incidence of fetal abnormalities when gestating females have been exposed to rather low dosage levels. There are no adequate and well-controlled studies in pregnant women.

Hydrocortisone acetate suppositories should only be used during pregnancy if the potential benefit justifies the risk of the fetus. Drugs of this class should not be used extensively on pregnant patients, in large amounts, or for prolonged periods of time.

It is not known whether this drug is excreted in human milk, and because many drugs are excreted in human milk and because of the potential for serious adverse reactions in

nursing infants from hydrocortisone acetate suppositories, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

ADVERSE REACTIONS

The following local adverse reactions have been reported with hydrocortisone acetate suppositories; burning, itching, irritation, dryness, folliculitis, hypopigmentation, allergic contact dermatitis, secondary infection.

To report an adverse event, please contact Rising Pharma Holdings, Inc. at 1-844-874-7464, or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG ABUSE AND DEPENDENCE

Drug abuse and dependence have not been reported in patients treated with hydrocortisone acetate suppositories.

OVERDOSAGE

If signs and symptoms of systemic overdose occur, discontinue use.

DOSAGE AND ADMINISTRATION

For rectal administration. Detach one suppository from strip of suppositories. Hold suppository upright. Separate tabs at top opening and pull downward from the pointed end. Continue pulling downward to almost the full length of the suppository. Carefully remove the suppository from the pocket. Avoid excessive handling of the suppository which is designed to melt at body temperature. Insert suppository into the rectum with gentle pressure, pointed end first. Insert one suppository in the rectum twice daily, morning and night for two weeks, in nonspecific proctitis. In more severe cases, one suppository three times a day or two suppositories twice daily. In factitial proctitis, the recommended duration of therapy is six to eight weeks or less, according to the response of the individual case.

HOW SUPPLIED

Hydrocortisone acetate suppositories 25 mg are white to off-white, smooth surfaced and bullet shaped with one pointed end.

Box of 12 suppositories, NDC 16571-676-21

Box of 24 suppositories, NDC 16571-676-42

STORAGE

Store at 20°-25°C (68°-77°F) [See USP Controlled Room Temperature]. Excursions permitted to 15°-30°C (59°-86°F). Store away from heat. Protect from freezing. Avoid contact with eyes.

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

Rx Only

Distributed by:

Rising Pharma Holdings, Inc.
East Brunswick, NJ 08816

Manufactured by:

Quagen Pharmaceuticals LLC
West Caldwell, NJ 07006

52030
Rev.01/21

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

Rising[®]

NDC 16571-676-21

**Hydrocortisone Acetate
Suppositories
25 mg**

For Rectal Administration

12 Adult Suppositories Rx only

No Varnish

Each suppository contains: Hydrocortisone Acetate 25 mg in a specially blended hydrogenated vegetable oil base. FOR RECTAL ADMINISTRATION.

Dosage and Administration: Read package insert for complete information before use.

Directions for Use: Detach one suppository from strip of suppositories. Hold suppository upright. Separate tabs at top opening and pull downward from the pointed end. Continue pulling downward to almost the full length of the suppository. Carefully remove the suppository from the pocket. Avoid excessive handling of the suppository, which is designed to melt at body temperature. Insert suppository into rectum with gentle pressure, pointed end first.

Store at 20-25°C (68-77°F) [See USP Controlled Room Temperature]. Store away from heat. Protect from freezing.

Note: Staining of fabric may occur with the use of the suppository. Precautionary measures as recommended.

Warning: Keep this and all drugs out of the reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

Distributed by:
Rising Pharma Holdings, Inc.
East Brunswick, NJ 08816

Manufactured by:
Quagen Pharmaceuticals LLC
West Caldwell, NJ 07006

Rising®
PHARMACEUTICALS

NDC 16571-676-21

**Hydrocortisone Acetate
Suppositories**

25 mg

For Rectal Administration

12 Adult Suppositories

Rx only

Rising®
Hydrocortisone Acetate
Suppositories
25 mg
For Rectal Administration
12 Adult Suppositories

NDC 16571-676-21

Rx only

51027
Rev. 12/2020



1.5" X 0.9375"
NON VARNISH
AREA

Rising®

NDC 16571-676-42

**Hydrocortisone Acetate
Suppositories
25 mg**

For Rectal Administration

24 Adult Suppositories

Rx only

No Varnish

Each suppository contains: Hydrocortisone Acetate 25 mg in a specially blended hydrogenated vegetable oil base, FOR RECTAL ADMINISTRATION.

Dosage and Administration: Read package insert for complete information before use.

Directions for Use: Detach one suppository from strip of suppositories. Hold suppository upright. Separate tabs at top opening and pull downward from the pointed end. Continue pulling downward to almost the full length of the suppository. Carefully remove the suppository from the pocket. Avoid excessive handling of the suppository, which is designed to melt at body temperature. Insert suppository into rectum with gentle pressure, pointed end first.

Store at 20-25°C (68-77°F) (See USP Controlled Room Temperature). Store away from heat. Protect from freezing.

Note: Staining of fabric may occur with the use of the suppository. Precautionary measures as recommended.

Warning: Keep this and all drugs out of the reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

Distributed by:
Rising Pharma Holdings, Inc.
East Brunswick, NJ 08816

Manufactured by:
Quagen Pharmaceuticals LLC
West Caldwell, NJ 07006

24 Adult Suppositories
For Rectal Administration
25 mg
Rising
PHARMACEUTICALS
Hydrocortisone Acetate
Suppositories
NDC 16571-676-42
Rx only

Rising
PHARMACEUTICALS

NDC 16571-676-42

Hydrocortisone Acetate
Suppositories

25 mg

For Rectal Administration

24 Adult Suppositories

Rx only

51028
Rev. 12/2020



1.5" X1.5"
NON VARNISH
AREA

HYDROCORTISONE ACETATE

hydrocortisone acetate suppository

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:16571-676
Route of Administration	RECTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROCORTISONE ACETATE (UNII: 3X7931PO74) (HYDROCORTISONE - UNII:W4X0X7BPJ)	HYDROCORTISONE ACETATE	25 mg

Inactive Ingredients

Ingredient Name	Strength
HYDROGENATED PALM KERNEL OIL (UNII: FM8D1RE2VP)	

Product Characteristics

Color	WHITE	Score	
Shape	BULLET	Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:16571-676-21	12 in 1 CARTON; Type 0: Not a Combination Product	02/01/2021	
2	NDC:16571-676-42	24 in 1 CARTON; Type 0: Not a Combination Product	02/01/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
UNAPPROVED DRUG OTHER		02/01/2021	

Labeler - Rising Pharma Holdings, Inc. (835513529)

Establishment

Name	Address	ID/FEI	Business Operations
QUAGEN PHARMACEUTICALS LLC		080281331	MANUFACTURE(16571-676) , PACK(16571-676)

Revised: 1/2021

Rising Pharma Holdings, Inc.