CLOTRIMAZOLE- clotrimazole cream Strategic Sourcing Services LLC

Clotrimazole Vaginal Cream, USP

Drug Facts

Active ingredients

Clotrimazole, USP 1% (50 mg in each applicatorful)

Active ingredients

Clotrimazole, USP 1% (external cream)

Purpose

Vaginal antifungal

Purpose

Vaginal antifungal

Uses

- treats vaginal yeast infections
- relieves external itching and irritation due to a vaginal yeast infection

Warnings

For vaginal use only

Do not use

if you have never had a vaginal yeast infection diagnosed by a doctor

Ask a doctor before use if you have

- vaginal itching and discomfort for the first time
- lower abdominal, back or shoulder pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge. You may have a more serious condition.
- vaginal yeast infections often (such as once a month or 3 in 6 months). You could be pregnant, or have a serious underlying medical cause for your symptoms, including diabetes or a weakened immune system.
- been exposed to the human immunodeficiency virus (HIV) that causes AIDS

When using this product

- do not use tampons, douches, spermicides, or other vaginal products. Condoms and diaphragms may be damaged and fail to prevent pregnancy or sexually transmitted diseases (STDs).
- do not have vaginal intercourse
- mild increase in vaginal burning, itching or irritation may occur
- if you do not get complete relief, ask a doctor before using another product

Stop use and ask a doctor if

- symptoms do not get better in 3 days
- symptoms last more than 7 days.
- you get a rash or hives, abdominal pain, fever, chills, nausea, vomiting, or a foul-smelling vaginal discharge

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- before using this product, read the enclosed educational brochure for complete directions and information
- adults and children 12 years of age and over:
 - **vaginal cream:** insert one applicatorful of cream into the vagina at bedtime for 7 days in a row. Wash applicator after each use.
 - external cream: use the same tube of cream if you have itching and irritation on the skin outside the vagina. Squeeze a small amount of cream onto your fingertip. Apply to itchy, irritated skin outside the vagina. Use 2 times daily for up to 7 days as needed.
- children under 12 years of age: ask a doctor

Other information

- to open tube: unscrew cap, lift tab, and pull to remove foil seal prior to use
- do not use if foil seal on tube opening is broken or missing
- do not purchase if carton is open
- store at room temperature 20° to 25°C(68° to 77°F)
- before using any medication, read all label directions, Keep carton, it contains important information.

Inactive ingredients

benzyl alcohol, cetostearyl alcohol, cetyl esters wax, 2-octyldodecanol, polysorbate 60, purified water, sodium phosphate monobasic, sorbitan monostearate

Questions?

Call 1-833-358-6431 Monday to Friday 9am to 7pm EST

Principal display panel

clotrimazole

Vaginal Cream, USP 1% • Vaginal Antifungal (with 1 Reusable Applicator)

Cures most vaginal yeast infections

7-day therapy / 7- day treatment

One 1.59 OZ (45g) TUBE OF VAGINAL CREAM (1%)

AND 1 REUSABLE PLASTIC APPLICATOR (7-DAY THERAPY

Distributed by McKesson Corp.,

via Strategic Sourcing Services LLC,

Memphis, TN 38141

www.sunmarkbrand.com

Package label

egnerhaeld isnigev gnillame-luot

Stop use and sak a doctor if III waymphome do not get better in 3 days — waymphome last more than 7 days — wayou get a reah or hivse, abdominal pain, fever, chills, neuses, vomiting, or a feet and a prouget a read or his section.

- # If you do not get complete relief, ask a doctor before using another product
- deesses (570s). • do not have veginal infercourse • mild increase in veginal burning, liching or inflation may occur

When using this product m do not use tempons, doubles, epermicides, or other vaginal products. Condoms and displicagins may be damaged and field to prevent pregnancy or sexually transmitted

- In Section (VIH) turn of the human immunodehology vinus (VIH) that causes AUCS
- s seçons nuquivub usequej quote sense jo: Xon. akubpuas' judnqua' qapeçes ou a xeziseueq juumne m xabiusi, keest juqecpous oyeu (sncy as ouce a moutu ou 3 ju @ moutps)' yon coniq pe buedusur' ou yawa qaseysadar yawa sense a mous sentons countgour
- Ask a doctor before use if you have m vaginal liching and discomforf for the first time m lower abdominal, back or alroubler pain, fever, chills, nausea, vomiting, or foul-amelling vaginal

Do not use it you have never had a vaginal yeast infection diagnosed by a doctor.

Warnings For vapinal use only.

USCS III treats vaginal yeast intections on the treats washing and intections are vaginal yeast intection

Active ingredients Vegme in each applicatorful, Vegma (1997 e.g. Vegma anthugal Cotimazole, USP 1% (50 mg in each applicatorful) Vegma (1997 e.g. Vegma anthugal Cotimazole, USP 1% (external cream)

Drug Facts

Questions? Cal 833-358-6431 Monday to Friday 9:00am to 7:00pm EST

Inactive ingredients becay storbe, estabasey storbe, ostyl esters way. 2-octyloodecanol, polysorteté EÚ, purited water, sodum phosphate monobasio, sorbiten monoabariate

- store at noom temperature 20° to 25°C (66° to 77°F) ■ before using any medication, read all label directors. Keep carton, it contains important information.
 - do not purchase if carton is open
 - m do not use if foil seal no fast and is some opening is broken or missing on ob m
 - m to open tube: unscrew cap, lift tab, and pull to remove foil seal prior to use

подъптотпі тэпо

m children under 12 years of age; ask a doctor

vagine. Use 2 times daily for up to 7 days as needed.

- regimer ¿dinectos o ameig amonunt or cueam outo Aom, protector y objecto, propor, impresso esqui contarge pue exclaiment cueamic nees pue escue prote or cueam qui Aom prave gicturos and insignipou ou pue aqui contarge pue esto prespor agrac escru nee:
- orecoons and interests here of age and over: • auths and children 12 years of age and over: • audits and children 12 years of age and over:


Divections - before using this product, read the enclosed educational brochure for complete

(1-800-222-1223) right away.

If pregnant or breast-feeding, sak a health professional before use.

The begond in contact a Poison Control Center of the poison contact a Poison Control Center of the professional cent

Drug Facts (continued)



NDC 49348-793-76

sun mark®

clotrimazole

Vaginal Cream, USP 1% - Vaginal Mallange Malla I Record to Amelicator

Cures most vaginal yeast infections 7-day therapy / 7-day treatment EDUCATIONAL PAMPHLET ENCLOSED

ONE 1.59 OZ (45 g) TUBE OF VAGINAL CREAM (1%) AND 1 REUSABLE PLASTIC APPLICATOR (7-DAY THERAPY) Product of Italy Rev. 06/2022

Code: GO/DRUGS/361 AAE3A/03



SUNMARK Vaginal Cream with Reusable Applicator

CLOTRIMAZOLE

clotrimazole cream

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49348-793
Route of Administration	VAGINAL		

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
CLOTRIMAZOLE (UNII: G07GZ97H65) (CLOTRIMAZOLE - UNII:G07GZ97H65)	CLOTRIMAZOLE	10 mg in 1 g		

Inactive Ingredients		
Ingredient Name	Strength	
BENZYL ALCOHOL (UNII: LKG8494WBH)		
CETOSTEARYL ALCOHOL (UNII: 2DMT128M1S)		
CETYL ESTERS WAX (UNII: D072FFP9GU)		
OCTYLDODECYL STEARATE (UNII: K6F16QGO28)		
POLYSORBATE 60 (UNII: CAL22UVI4M)		
WATER (UNII: 059QF0KO0R)		
SODIUM PHOSPHATE, MONOBASIC, MONOHYDRATE (UNII: 593YOG76RN)		
SORBITAN MONOSTEARATE (UNII: NVZ4I0H58X)		

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49348-793- 76	1 in 1 CARTON	07/19/1993	
1		45 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information			
Marketing	Application Number or Monograph	Marketing Start	Marketing End

Category	Citation	Date	Date
ANDA	ANDA074165	07/19/1993	

Labeler - Strategic Sourcing Services LLC (116956644)

Revised: 10/2022 Strategic Sourcing Services LLC