

**GRAVOL CHEWABLES MOTION SICKNESS DIMENHYDRINATE-**  
**dimenhydrinate tablet, chewable**  
**Church & Dwight Co., Inc.**

-----  
**Gravol™ Chewables Motion Sickness Dimenhydrinate**

***Drug Facts***

**Active ingredient**

Dimenhydrinate 50 mg

**Purpose**

antiemetic

**Uses**

For the prevention and treatment of the nausea, vomiting and dizziness associated with motion sickness

**Warnings**

**Do not use** for children under 2 years of age unless directed by a doctor

**Ask a doctor before use if you or your child has**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

**Ask a doctor before use if you or your child is** taking a sedative or tranquilizer

**When using this product**

- may cause marked drowsiness
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase the drowsiness effect
- use caution when driving a motor vehicle or operating machinery

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison control Center (1-800-222-1222) right away.

**Directions**

To prevent motion sickness, the first dose should be taken ½ to 1 hour before starting activity

- Adults and children 12 years of age and older: take 1 to 2 tablets every 4 to 6 hours. Do not exceed 8 tablets in 24 hours, or as directed by a doctor
- Children 6 to under 12 years of age: take ½ to 1 tablet every 6 to 8 hours. Do not exceed 3 tablets in 24 hours, or as directed by a doctor
- Children 2 to under 6 years of age: give ½ tablet every 6-8 hours. Do not give more than 1 ½ tablets in 24 hours, or as directed by a doctor

### **Other information**

- **Phenylketonurics: contains phenylalanine 15 mg per tablet**
- Store between 59° - 77°F

### **Inactive ingredients**

acetylated monoglycendes, amino methacrylate copolymer, aspartame, citric acid, com starch, ethylcellulose, FD&C yellow 6 aluminum lake, magnesium stearate, mannitol, microcrystalline cellulose, orange flavor, polyethylene glycol

### **Questions or comments?**

Call us at **1-888-334--5387** M-F 9am-5pm ET

### **PRINCIPAL DISPLAY PANEL - 50 mg Blister Pack Carton**

Prevents and Treats:  
Nausea, Vomiting  
& Dizziness\*

\*Prevention and treatment of nausea,  
vomiting and dizziness due to motion sickness

Gravol™ CHEWABLES

Gravol™  
MOTION SICKNESS  
Dimenhydrinate tablets 50 mg

8  
TABLETS

ORANGE  
FLAVOR

Prevents and Treats:  
**Nausea, Vomiting  
 & Dizziness\***

\*Prevention and treatment of nausea,  
 vomiting and dizziness due to motion sickness

**Gravol™ CHEWABLES**

**Gravol™**

**MOTION SICKNESS**  
 Dimenhydrinate tablets 50 mg

**8**  
 TABLETS



ORANGE  
 FLAVOR

**Gravol™ CHEWABLES**

<b>Drug Facts</b>	
<b>Active ingredient</b>	Dimenhydrinate 50 mg
<b>Purpose</b>	antiemetic
<b>Uses</b>	For the prevention and treatment of the nausea, vomiting and dizziness associated with motion sickness
<b>Warnings</b>	Do not use for children under 2 years of age unless directed by a doctor Ask a doctor before use if you or your child has • a breathing problem such as emphysema or chronic bronchitis • glaucoma • difficulty in urination due to enlargement of the prostate gland Ask a doctor before use if you or your child is taking a sedative or tranquilizer <b>When using this product</b> • alcohol, sedatives and tranquilizers may increase the drowsiness effect • may cause marked drowsiness • avoid alcoholic beverages • use caution when driving a motor vehicle or operating machinery <b>If pregnant or breast-feeding</b> , ask a health professional before use. <b>Keep out of reach of children.</b> In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.
<b>Directions</b>	To prevent motion sickness, the first dose should be taken ½ to 1 hour before starting activity • Adults and children 12 years of age and older: take 1 to 2 tablets every 4 to 6 hours. Do not exceed 8 tablets in 24 hours, or as directed by a doctor • Children 6 to under 12 years of age: take ½ to 1 tablet every 6 to 8 hours. Do not exceed 3 tablets in 24 hours, or as directed by a doctor • Children 2 to under 6 years of age: give ½ tablet every 6-8 hours. Do not give more than 1 ½ tablets in 24 hours, or as directed by a doctor
<b>Other information</b>	Phenylethanolamine 15 mg per tablet • Store between 59° - 77°F <b>Inactive ingredients</b> acetylated monoglycerols, amino methacrylate copolymer, aspartame, citric acid, corn starch, ethylcellulose, FD&C yellow 6 aluminum lake, magnesium stearate, mannitol, microcrystalline cellulose, orange flavor, polyethylene glycol 98m-5pm ET
<b>Questions or comments?</b>	Call us at 1-888-334-5387 M-F

0 22600 00283 3  
 Church & Dwight Co., Inc., Ewing, NJ 08628  
 Gravol is a trademark of Church & Dwight Canada Corp.  
 Made in Canada

**Gravol™**



GRFC-00283-01 72028257



# GRAVOL CHEWABLES MOTION SICKNESS DIMENHYDRINATE

dimenhydrinate tablet, chewable

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:10237-943
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DIMENHYDRINATE</b> (UNII: JB937PER5C) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIMENHYDRINATE	50 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>DIACETYLATED MONOGLYCERIDES</b> (UNII: 5Z17386USF)	
<b>DIMETHYLAMINOETHYL METHACRYLATE - BUTYL METHACRYLATE - METHYL METHACRYLATE COPOLYMER</b> (UNII: 905HNO1SIH)	
<b>ASPARTAME</b> (UNII: Z0H242BBR1)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>ETHYLCELLULOSE (10 MPA.S)</b> (UNII: 3DYK7UYZ62)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	

## Product Characteristics

<b>Color</b>	ORANGE	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	14mm
<b>Flavor</b>	ORANGE	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10237-943-08	1 in 1 CARTON	04/01/2024	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M009	04/01/2024	

**Labeler** - Church & Dwight Co., Inc. (001211952)

## Establishment

Name	Address	ID/FEI	Business Operations
Church & Dwight Canada Corp		253933600	MANUFACTURE(10237-943)

Revised: 3/2024

Church & Dwight Co., Inc.