

**MENSTRUAL COMPLETE- acetaminophen, caffeine, pyrilamine maleate tablet, film coated**

**Rite Aid Corporation**

-----

**Rite Aid 44-390**

***Active ingredients (in each caplet)***

Acetaminophen 500 mg

Caffeine 60 mg

Pyrilamine maleate 15 mg

***Purpose***

Pain reliever

Diuretic

Antihistamine

***Uses***

for the temporary relief of these symptoms associated with menstrual periods:

- bloating
- headache
- water-weight gain
- cramps
- backache
- fatigue
- muscle aches

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- difficulty in urination due to enlargement of the prostate gland
- liver disease
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- excitability may occur, especially in children
- use caution when driving a motor vehicle or operating machinery
- limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart rate. The recommended dose of this product contains about as much caffeine as a cup of coffee.

### **Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### ***Directions***

- **do not take more than directed**
- adults and children 12 years and over:
  - take 2 caplets with water
  - repeat every 6 hours, as needed

- do not exceed 6 caplets per day
- children under 12 years: ask a doctor

### ***Other information***

- see end flap for expiration date and lot number
- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)

### ***Inactive ingredients***

corn starch, croscarmellose sodium, crospovidone, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, silicon dioxide, stearic acid, titanium dioxide, triacetin

### ***Questions or comments?***

**1-800-426-9391**

### **Principal Display Panel**

NDC 11822-3900-8

Compare to active ingredients in **Midol® Complete\***

#### **MENSTRUAL COMPLETE**

**ACETAMINOPHEN, CAFFEINE, PYRILAMINE MALEATE**  
pain reliever • diuretic • antihistamine

#### **MULTI-SYMPTOM RELIEF**

of cramps, bloating, fatigue,  
headache & backache

ACTUAL SIZE

**24**

CAPLETS

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

\*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Midol® Complete.

50844 ORG062339008

**DISTRIBUTED BY: RITE AID,  
200 NEWBERRY COMMONS  
ETTERS, PA 17319 [www.riteaid.com](http://www.riteaid.com)**

### **SATISFACTION GUARANTEE**

If you're not satisfied, we'll

happily refund your money.

NDC 11822-3900-8      Compare to active ingredients in Midol® Complete\*

# MENSTRUAL COMPLETE

ACETAMINOPHEN, CAFFEINE, PYRILAMINE MALEATE  
pain reliever • diuretic • antihistamine

**MULTI-SYMPTOM RELIEF**

of cramps, bloating, fatigue,  
headache & backache

**24 CAPLETS**

ACTUAL SIZE

no print / no varnish area  
lot no. & exp. date

0 11822 02141 8

**TAMPER EVIDENT. DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

\*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Midol® Complete.  
50844    ORG062339008

**DISTRIBUTED BY: RITE AID**  
200 NEUBERRY COMMONS  
ETTERS, PA 17319 [www.riteaid.com](http://www.riteaid.com)

**SATISFACTION GUARANTEE**  
If you're not satisfied, we'll happily refund your money.

**Drug Facts (continued)**

■ excitability may occur, especially in children  
■ use caution when driving a motor vehicle or operating machinery

■ limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart rate. The recommended dose of this product contains about as much caffeine as a cup of coffee.

**Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison

**Drug Facts (continued)**

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- difficulty in urination due to enlargement of the prostate gland
- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers
- taking sedatives or tranquilizers

**When using this product**

- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Drug Facts (continued)**

**Control Center** night away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed
- adults and children 12 years and over: take 2 caplets with water
- do not exceed 6 caplets per day
- children under 12 years: ask a doctor

**Other information**

- store at 25°C (77°F); excursions permitted

**Drug Facts (continued)**

between 15°-30°C (59°-86°F) ■ see end flap for expiration date and lot number

**Inactive ingredients**

corn starch, croscarmellose sodium, croscopolidone, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, silicon dioxide, stearic acid, titanium dioxide, trace b1

**Questions or comments?**

1-800-426-9391

**Drug Facts (continued)**

**Warnings**

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Alergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ rash ■ skin reddening ■ blisters ■ muscle aches

**Uses**

for the temporary relief of these symptoms associated with menstrual periods: ■ bloating ■ headache ■ muscle aches ■ cramps ■ backache ■ fatigue ■ water-weight gain

**Drug Facts (continued)**

**Active Ingredients (in each caplet)**

Acetaminophen 500 mg ..... Pain reliever  
Caffeine 60 mg ..... Diuretic  
Pyrilamine maleate 15 mg ..... Antihistamine

**Purpose**

FOR COMPLETE PRODUCT INFORMATION

**KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

**Drug Facts (continued)**

help night away.

If a skin reaction occurs, stop use and seek medical

**Drug Facts (continued)**

**Warnings**

Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Alergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ rash ■ skin reddening ■ blisters ■ muscle aches

**Uses**

for the temporary relief of these symptoms associated with menstrual periods: ■ bloating ■ headache ■ muscle aches ■ cramps ■ backache ■ fatigue ■ water-weight gain

**Drug Facts (continued)**

**Active Ingredients (in each caplet)**

Acetaminophen 500 mg ..... Pain reliever  
Caffeine 60 mg ..... Diuretic  
Pyrilamine maleate 15 mg ..... Antihistamine

**Purpose**

FOR COMPLETE PRODUCT INFORMATION

**KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

Rite Aid 44-390

# MENSTRUAL COMPLETE

acetaminophen, caffeine, pyrilamine maleate tablet, film coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-3900
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	60 mg
<b>PYRILAMINE MALEATE</b> (UNII: R35D29L3ZA) (PYRILAMINE - UNII:HPE317O9TL)	PYRILAMINE MALEATE	15 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	44;390
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-3900-8	3 in 1 BOX	06/13/2024	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:11822-3900-1	1 in 1 CARTON	06/19/2024	
		40 in 1 BOTTLE PLASTIC Type 0 N/A		

2

40 IN 1 BOTTLE, PLASTIC; Type U: Not a Combination Product

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	06/13/2024	

**Labeler** - Rite Aid Corporation (014578892)**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(11822-3900)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(11822-3900) , pack(11822-3900)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(11822-3900)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(11822-3900)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(11822-3900)

Revised: 6/2024

Rite Aid Corporation