COLD RELIEF MULTI-SYMPTOM- acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, and phenylephrine hydrochloride TopCare

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

1150 - TCR - 2022-1208

COLD MAX DAY

Drug Facts

Active ingredients (in each day caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these common cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - cough
 - sinus congestion and pressure
- helps clear nasal passages
- promotes nasal and sinus drainage
- temporarily reduces fever

COLD MAX NIGHT

Drug Facts

Active ingredients (in each night caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg	Antihistamine
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these common cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - runny nose and sneezing
 - cough
 - sinus congestion and pressure
- helps clear nasal passages
- relieves cough to help you sleep
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 12 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease

- high blood pressure
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

do not exceed recommended dosage

In addition, when using Cold Multi-Symptom Night:

- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- do not take Day and Night caplets at the same time
- do not take more than a total of 12 caplets in 24 hours

- take 2 caplets every 4 hours
- swallow whole do not crush,

12 years and	chew, or dissolve
over	
children under 12 years	ask a doctor

Other information

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information

Inactive ingredients

Cold Multi-Symptom Day

acesulfame potassium, colloidal silicon dioxide, croscarmellose sodium, crospovidone, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

Cold Multi-Symptom Night

acesulfame potassium, colloidal silicon dioxide, croscarmellose sodium, FD&C blue #1 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

PRINCIPAL DISPLAY PANEL

TopCare® Health

36800-250-56

Cold Max

Day + Night

For Adults

Our Pharmacists Recommend

Daytime

Pain Reliever/Fever Reducer - Acetaminophen

Cough Suppressant – Dextromethorphan HBr

Nasal Decongestant - Phenylephrine HCl

Day Caplet Relieves:

- Head + Body Aches
- Fever + Sore Throat
- Cough

Nasal Congestion

12 Cool Taste Caplets

Nighttime


Pain Reliever/Fever Reducer - Acetaminophen

Nasal Decongestant - Phenylephrine HCl

Antihistamine - Chlorpheniramine Maleate*

Night Caplet Relieves:

- Head + Body Aches
- Fever + Sore Throat
- Cough
- Nasal Congestion
- Runny Nose*
- *Antihistamine in Nighttime Only
- 8 Cool Taste Caplets



Cold Max Day + Night

+TopCare * health

NDC 36800-250-56

For Adults

Cold Max Day + Night



PAIN RELIEVER/FEVER REDUCER - ACETAMINOPHEN

COUGH SUPPRESSANT - DEXTROMETHORPHAN HBr NASAL DECONGESTANT - PHENYLEPHRINE HCI

DAY CAPLET RELIEVES:

- · Head + Body Aches
- Fever + Sore Throat
- Cough
- Nasal Congestion



actual size

NIGHTTIME

PAIN RELIEVER/FEVER REDUCER -

ACETAMINOPHEN

COUGH SUPPRESSANT - DEXTROMETHORPHAN HBr NASAL DECONGESTANT - PHENYLEPHRINE HCI ANTIHISTAMINE* - CHLORPHENIRAMINE MALEATE*

NIGHT CAPLET RELIEVES:

- Head + Body Aches
- Fever + Sore Throat
- Cough
- Nasal
 - Congestion
- Runny Nose*

*Antihistamine in Nighttime Only



actual size

12 COOL TASTE

8 COOL TASTE

Inactive ingredients

Cold Max Day sessultame potassium, colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, propylene glycol, stearic scid, talc, titanium dioxide crospovidone, pregelatinized starch, propylene glycol, stearic scid, talc, titanium dioxide, corn starch, croscarmellose sodium, crospovidone, FD&C blue #1 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

Drug Facts (continued)

DISTRIBUTED BY
TOPCO ASSOCIATES LLC
ELK GROVE VILLAGE, IL 60007
©TOPCO AAPA0922
QUESTIONS? 1-888-423-0139
topcare@topco.com
www.topcarebrand.com

Scan here for information



Other information a store between 20-25 °C (68-77 °F) in a dry place retain carton for complete product information and warnings

sdults and children

adults and over

swallow whole; do not crush, chew, or dissolve children under 12 years

ask a doctor

do not take Day and Might caplets at the same time
 do not take more than a total of 10 caplets in 24 hours

DIRECTIONS = do not take more than directed (see overdose warning)

any signs or symptoms.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away

(1-800-222-1222), Quick medical attention is critical for adults as well as for children even if you do not notice

condu course pack or occurs with rash or headache that lasts

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

These could be signs of a serious condition.

uew symptoms occur
 iever gets worse or issus

■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present

S



Stop use and ask a doctor if a nervousness, dizziness, or sleeplessness occur be careful when driving a motor vehicle or operating machinery svoid alcoholic drinks sicohol, sedatives, and tranquilizers may increase drowsiness marked drowsiness may occur excitability may occur, especially in children In addition, when using Cold Max Night: do not exceed recommended dosage When using this product taking the blood thinning drug warfarin taking sedatives or tranquilizers Ask a doctor or pharmacist before use if you are s preathing problem such as emphysema or chronic bronchius bersistent or chronic cough such as occurs with smoking, asthma, or emphysema condu that occurs with too much phiegm (mucus) ■ trouble uninating due to an enlarged prostate gland ■ liver disease ■ heart disease ■ thyroid disease ■ high blood pressure ■ diabetes ■ glaucoma Ask a doctor before use if you have if you have ever had an allergic reaction to this product or any of its ingredients F115002TCR do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, **Drug Facts** (continued)

2

COLD RELIEF MULTI-SYMPTOM

acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, and phenylephrine hydrochloride kit

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:36800-250

ı	Packaging			
4	tem Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:36800-250-56	1 in 1 CARTON	10/01/2011	

Quantity of Parts			
Part #	Package Quantity	Total Product Quantity	
Part 1	1 BLISTER PACK	12	
Part 2	1 BLISTER PACK	8	

Part 1 of 2

ACETAMINOPHEN, DEXTROMETHORPHAN HYDROBROMIDE, PHENYLEPHRINE HYDROCHLORIDE

acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride tablet, coated

Product Information

ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Inactive Ingredients	
Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)	
CROSPOVIDONE (UNII: 2S7830E561)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
POVIDONE (UNII: FZ 989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	white	Score	no score
Shape	OVAL (capsule-shaped)	Size	17mm
Flavor	MINT	Imprint Code	AAA;1138
Contains			

l	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341		

Part 2 of 2

ACETAMINOPHEN, CHLORPHENIRAMINE MALEATE, DEXTROMETHORPHAN HYDROBROMIDE, PHENYLEPHRINE HYDROCHLORIDE

acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, and phenylephrine hydrochloride tablet, coated

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg	
CHLORPHENIRAMINE MALEATE (UNII: V1Q0090J9Z) (CHLORPHENIRAMINE - UNII: 3U6I01965U)	CHLORPHENIRAMINE MALEATE	2 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Inactive Ingredients	
Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics				
Color	blue	Score	no score	
Shape	OVAL (capsule-shaped)	Size	17mm	
Flavor	MINT	Imprint Code	AAA;1139	
Contains				

Packaging				
#	ltem Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information					
	Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
	OTC monograph final	part341			

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC monograph final	part341	10/01/2011			

Labeler - TopCare (006935977)

Revised: 12/2022 TopCare