

**CORICIDIN HBP MAXIMUM STRENGTH COLD FLU AND CHEST CONGESTION-
acetaminophen, guaifenesin capsule, gelatin coated
Bayer HealthCare LLC.**

Coricidin[®] HBP Maximum Strength Cold, Flu & Chest Congestion Liquid Gels
Drug Facts

Active ingredients

(in each tablet)	Purpose
Acetaminophen 325 mg	Pain reliever/ fever reducer
Guaifenesin 200 mg	Expectorant

Uses

- temporarily relieves these symptoms due to a cold or flu:
 - minor aches and pains
 - headache
 - sore throat
 - helps loosen phlegm(mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 4,000 mg of acetaminophen in 24 hours • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include: • skin reddening • blisters • rash • hives • facial swelling • asthma (wheezing) • shock If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

- in children under 12 years of age

Ask a doctor before use if you have ● liver disease ● persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ● cough that is accompanied by excessive phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if • pain or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.
- children under 12 years of age: do not use

Other information

- store at 15°-25°C (59°-77°F)

Inactive ingredients

FD&C yellow # 6, gelatin, glycerin, iron oxide black, lecithin, medium-chain triglycerides, mica, polyethylene glycol, potassium aluminum silicate, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

?

Questions or comments? 1-800-317-2165 (Mon-Fri 9AM -5PM EST)

PRINCIPAL DISPLAY PANEL - 24 Liquid Gels


Coricidin ®

HIGH BLOOD PRESSURE

Acetaminophen - Pain Reliever/Fever Reducer

Guaifenesin - Expectorant

24 Liquid Gels



CORICIDIN HBP MAXIMUM STRENGTH COLD FLU AND CHEST CONGESTION

acetaminophen, guaifenesin capsule, gelatin coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11523-0094
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
MICA (UNII: V8A1AW0880)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
SHELLAC (UNII: 46N107B71O)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
SORBITAN (UNII: 6O92ICV9RU)	
POTASSIUM ALUMINUM DISILICATE (UNII: SRB14JRX6C)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL (Oblong)	Size	18mm
Flavor		Imprint Code	CHBPD
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11523-0094-1	2 in 1 CARTON	05/02/2022	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/01/2022	

Labeler - Bayer HealthCare LLC. (112117283)