AXIM ALLERGY- diphenhydramine hydrochloride capsule, liquid filled VIVUNT LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Axim Allergy

Drug Facts

Active Ingredient (in each softgel)	Purpose
Diphenhydramine HCl 25 mg	Antihistamine

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- temporarily relieves these symptoms due to the common cold:
- runny nose
- sneezing

Warnings

Do not use

- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Keep out of reach of children.

If pregnant or breast-feeding

ask a health professional before use.

Overdose Warning

In case of overdose, get medical help or contact a poison control center immediately.

Directions

- Take every 4 to 6 hours, or as directed by a doctor
- Do not take more than 12 Softgels in 24 hours for Adults and do not take more than 6 Softgels for children 6 to 12 years of age in 24 hours.

adults and children 12 years and over	1 or 2 Softgels
children 6 to under 12 years	1 Softgel
children under 6 years	do not use

Other information

- Store at 20° 25 °C (68 °- 77 °F)
- Read all product information before using
- Tamper Evident: Do not use if the foil printed on the blister is torn or ripped.

Inactive ingredients

Gelatin, Glycerin, Methylparaben, Polyethylene Glycol 400, Propylparaben, Purified Water, Sorbitol, Titanium Dioxide

PRINCIPAL DISPLAY PANEL 24

Compare to Benadryl® Allergy Liqui-Gels® active ingredients*

NDC 82706-005-01


Antihistamine

- Runny Nose
- Sneezing
- Itchy, Watery Eyes
- Itchy Throat

DYE-FREE

Diphenhydramine HCl

24 SOFTGELS** **Liquid-filled capsules





FTARM LLC V 74th Court, Suite 1901 Florida Z.C. 33156-3178



PRINCIPAL DISPLAY PANEL 6

Compare to Benadryl® Allergy Liqui-Gels®

active ingredients*

NDC 82706-005-02

Antihistamine

Runny Nose Sneezing Itchy, Watery Eyes Itchy Throat

DYE-FREE

Diphenhydramine HCl

6 SOFTGELS** **Liquid-filled capsules





Allergy Relief

ANTIHISTAMINE



Diphenhydramine HCI

6 SOFTGELS**

**Liquid-filled capsules



Sorbitol, Titanium Dioxide.

Gelatin, Glycerin, Methylparaben, Polyethylene Glycol 400, Propylparaben, Purified Water,

Inactive ingredients

- Tamper evident: Do not use if carton or pouch is open.
 - Read all product information before using
 - Store at 20°-25°C (68°-77°F)

Other information

- ■children under 6 years: do not use
- children 6 to under 12 years: 1 softgel
- ■adults and children over 12 years: 1 or 2 softgels
 - for children 6 to 12 years of age in 24 hours.
- do not take more than 12 softgels in 24 hours for adults and do not take more than 6 softgels
 - take every 4 to 6 hours, or as directed by a doctor

Directions

immediately.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center It pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

- excitability may occur, especially in children.
- may increase drowsiness be careful when driving a motor vehicle or operating machinery ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers When using this product do not exceed recommended dose

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

bronchitis

glaucoma

trouble urinating due to an enlarged prostate gland. Ask a doctor before use if you have a a breathing problem such as emphysema or chronic

■ to make a child sleep ■ with any other product containing diphenhydramine, even one used on Do not use

Warnings

relieves these symptoms due to the common cold: ■ runny nose ■ sneezing ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat ■ temporarily


Uses Temporarily relieves these symptoms due to hay fever or other upper respiratory

Antihistamine Purpose

Diphenhydramine HCI 25 mg Active ingredient (in each softgel)

ยาวธา ยูมาน

Barcode



AXIM ALLERGY

diphenhydramine hydrochloride capsule, liquid filled

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:82706-005
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg	

Inactive Ingredients		
Ingredient Name	Strength	
PROPYLPARABEN (UNII: Z8IX2SC10H)		
GELATIN (UNII: 2G86QN327L)		
METHYLPARABEN (UNII: A218C7HI9T)		
WATER (UNII: 059QF0KO0R)		
GLYCERIN (UNII: PDC6A3C0OX)		
SORBITOL (UNII: 506T60A25R)		
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)		
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)		

Product Characteristics				
Color white (Transparent) Score no score				
Shape	OVAL	Size	22mm	
Flavor		Imprint Code	axim	
Contains				

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82706- 005-01	2 in 1 CARTON	05/09/2022	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:82706- 005-02	3 in 1 CARTON	09/20/2023	

2	in 1 POUCH; Type 0: Not a Combination Product		
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	05/09/2022	

Labeler - VIVUNT LLC (045829437)

Revised: 9/2023 VIVUNT LLC