

**MILKHOUSE BRAND IODINE TEAT DIP CONTROLLED- iodine liquid**  
**Stearns Packaging Corporation**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----

**Controlled Iodine Teat Dip**

**TEAT DIPPING is an important part of a  
MASTITIS CONTROL PLAN:**

1. Dip teats immediately after milking.
2. Use good milking and environmental management procedures.
3. Use properly functioning milking equipment.
4. Treat every quarter of every cow at dry off.
5. Identify and treat or manage clinical cases promptly.
6. Cull chronically infected cows.

**TEAT DIP**

**GREAT FOR FOAMING**

**TEAT DIPPING IN COLD WEATHER:**

When the temperature falls below 10 degrees F, or when wind chill is significant, special precautions should be taken to avoid chapped and frozen teats.

1. Teats should be dry before turning cows out into cold weather.
2. When teats are dipped after milking, allow 30 seconds contact, and blot off any excess teat dip with a single service towel.
3. Warm the teat dip to reduce drying time.
4. Provide windbreaks in outside areas for cows.
5. Monitor fresh cows with swollen udders and teats since they are more susceptible to chapped and frozen teats.

**DIRECTIONS FOR USE:**

- DO NOT DILUTE.
- Dip teats immediately after milking as a routine part of milking procedures.
- Dip to THOROUGHLY COVER at least the lower one-third of the teat (preferably to the base of the udder), IMMEDIATELY after every milking.

**TAKE TIME**

**OBSERVE**

**LABEL DIRECTIONS**

- Empty and clean dip cups after each milking, or during milking, if they become contaminated.
- Do not pour used dip back into original container.
- KEEP FROM FREEZING
- Keep container closed to prevent contamination.
- Store container in an upright position.

**FIRST-AID**

**EYES:** If contact with eyes occurs, flush with plenty of cool water for 15 minutes. Consult a physician.

**INTERNAL:** May be harmful if swallowed. If ingested, drink large amounts of water. DO NOT induce vomiting. Get medical attention immediately.

**24 HOUR EMERGENCY CONTACT:  
1-800-255-3924**

**MILKHOUSE BRAND**

**Controlled  
Iodine Teat Dip**

**A SANITIZING TEAT DIP  
An Aid in Reducing the Spread of  
Organisms Which May Cause Mastitis**

**ACTIVE INGREDIENTS**

Concentrated Iodophor, containing Nonylphenol, ethoxylated iodine complex.  
(provides 1/2% titratable iodine, equivalent to 5,000 ppm titratable iodine)

**EMOLLIENT SYSTEM**

Contains 5% glycerin

**KEEP OUT OF THE REACH OF CHILDREN**

See side panel for Precautionary Statements

**NET CONTENTS: 1 U.S. FL. GALLON (3.78 LITERS)**

**STEARNS PACKAGING CORPORATION**

4200 Sycamore Ave. (53714), PO Box 3216, Madison, WI 53704  
(800) 655-5008

**DIRECTIONS FOR USE:**

- DO NOT DILUTE.
- Dip teats immediately after milking as a routine part of milking procedures.
- Dip to THOROUGHLY COVER at least the lower one-third of the teat (preferably to the base of the udder), IMMEDIATELY after every milking.
- Empty and clean dip cups after each milking, or during milking, if they become contaminated.
- Do not pour used dip back into original container.
- KEEP FROM FREEZING
- Keep container closed to prevent contamination.
- Store container in an upright position.

**TEAT DIPPING is an important part of a**

**MASTITIS CONTROL PLAN:**

1. Dip teats immediately after milking.
2. Use good milking and environmental management procedures.
3. Use properly functioning milking equipment.
4. Treat every quarter of every cow at dry off.
5. Identify and treat or manage clinical cases promptly.
6. Cull chronically infected cows.

**TEAT DIPPING IN COLD WEATHER:**

- When the temperature falls below 10°F, or when wind chill is significant, special precautions should be taken to avoid chapped and frozen teats.
1. Teats should be dry before turning cows out into cold weather.
  2. When teats are dipped after milking, allow 30 seconds contact, and blot off any excess teat dip with a single service towel.
  3. Warm the teat dip to reduce drying time.
  4. Provide windbreaks in outside areas for cows.
  5. Monitor fresh cows with swollen udders and teats since they are more susceptible to chapped and frozen teats.



**GREAT FOR FOAMING!**



**A SANITIZING TEAT DIP  
An Aid in Reducing the Spread of  
Organisms Which May Cause Mastitis**

**ACTIVE INGREDIENTS:**  
Concentrated Iodophor, containing Nonylphenol, ethoxylated iodine complex.  
(provides 1/2% titratable iodine, equivalent to 5,000 ppm titratable iodine)

**EMOLLIENT SYSTEM:**  
Contains 5% glycerin.

**KEEP OUT OF THE REACH OF CHILDREN**  
See side panel for Precautionary Statements.

**NET CONTENTS: 1 U.S. FL. GALLON (3.78 LITERS)**

**STEARNS PACKAGING CORPORATION**  
4200 Sycamore Ave. (53714), PO Box 3216, Madison, WI 53704  
(800) 655-5008

**FIRST-AID:**

**EYES:** If contact with eyes occurs, flush with plenty of cool water for 15 minutes. Consult a physician.  
**INTERNAL:** May be harmful if swallowed. If ingested, drink large amounts of water. DO NOT induce vomiting. Get medical attention immediately.

**PRIMEROS AUXILIOS:**

**OJOS:** Si ocurre contacto con los ojos, lavar con abundante agua fría durante 15 minutos. Consulte a un médico.  
**INTERNO:** Puede ser nocivo en caso de ingestión. Beba grandes cantidades de agua si se ingiere este producto. NO induzca el vómito. Obtenga atención médica inmediatamente.

**24 HOUR EMERGENCY CONTACT**

24 horas servicio telefónico de respuesta a emergencias:  
1-800-255-3924

**ST-200**



LOT NO.:

EXP. DATE:

**INSTRUCCIONES POR EL USO:**

- NO lo diluir.
- Sumergir los pezones inmediatamente después del ordeño como parte rutinaria de procedimientos de ordeño.
- Sumergir para cubrir a fondo al menos el tercio inferior de la tetina (preferiblemente a la base de la ubre), inmediatamente después de cada ordeño.
- Vaciar y limpiar las tetas sumergiendo después de cada ordeño, o durante el ordeño, si contaminan.
- NO DEJE QUE SE CONGELÉ.
- Mantenga el recipiente bien cerrado para evitar la contaminación.
- Mantenga el recipiente en una posición vertical.

**BAÑO DE PEZONES ES UNA PARTE IMPORTANTE DE UN PLAN PARA CONTROL**

**DE LA MASTITIS:**

1. Sumerja los pezones inmediatamente después del ordeño.
2. Utilice buenas prácticas de gestión para el ordeño y medio ambiente.
3. Utilice equipo de ordeño que funcione correctamente.
4. Trate a cada cuarto de la ubre de cada vaca el dentro de fase de secado.
5. Identifique y trate o maneje casos clínicos con prontitud.
6. Entienda las vacas que están infectadas crónicamente.

**BAÑO DE PEZONES EN TIEMPO FRÍO:**

- Cuando la temperatura cae por debajo de 10°F (-12°C), o cuando el factor de enfriamiento del viento es importante, tome precauciones especiales para evitar los pezones agitados y congelados.
1. Los pezones deben estar secos antes de zollar las vacas en el frío.
  2. Cuando los pezones se sumergen después del ordeño, espere 30 segundos de contacto, y posteriormente secalos el exceso de baño de pezones con una toalla desechable.
  3. Caliente el baño de pezones para reducir el tiempo de secado.
  4. Proporcione protección contra el viento en las áreas fuera para las vacas.
  5. Vigile las vacas frescas con ubres hinchadas y pezones ya que son más susceptibles a los pezones agitados y congelados.



<b>MILKHOUSE BRAND IODINE TEAT DIP CONTROLLED</b>			
iodine liquid			
<b>Product Information</b>			
<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:60282-1200
<b>Route of Administration</b>	TOPICAL		
<b>Active Ingredient/Active Moiety</b>			
<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>	
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	0.5 mg in 100 L	
<b>Inactive Ingredients</b>			
<b>Ingredient Name</b>	<b>Strength</b>		
GLYCERIN (UNII: PDC6A3C0OX)	5 mg in 100 L		

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:60282-1200-1	3.78 L in 1 BOTTLE, PLASTIC		
2	NDC:60282-1200-2	18.93 L in 1 PAIL		
3	NDC:60282-1200-3	56.78 L in 1 DRUM		
4	NDC:60282-1200-4	208.2 L in 1 DRUM		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		01/02/1982	

**Labeler** - Stearns Packaging Corporation (006069256)

**Registrant** - Stearns Packaging Corporation (006069256)

## Establishment

Name	Address	ID/FEI	Business Operations
Stearns Packaging Corporation		006069256	manufacture, api manufacture

Revised: 2/2019

Stearns Packaging Corporation