# IBUPROFEN PM- diphenhydramine citrate, ibuprofen tablet, coated Family Dollar (FAMILY WELLNESS)

-----

# **Drug Facts**

# Active ingredient (in each caplet)

Diphenhydramine citrate 38 mg ibuprofen 200 mg(NSAID)\*

\*nonsteroidal anti-inflammatory drug

### **Purpose**

Nighttime sleep-aid

Pain Reliever

#### Uses

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

# Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart

attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed

#### Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- unless you have time for a full night's sleep
- in children under 12 years of age
- right before or after surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

#### Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have problem or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma or had a stroke
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

# Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any continuing medical illness
- taking any other antihistamines
- taking aspirin for heart attack or stroke, because ibuprofen may decease this benefit of aspirin
- taking any other drug

# When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- · take with food or milk if stomach upset occurs

# Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problem or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body

- slurred speech
- leg swelling
- pain gets worse or last more than 10 days
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness
- redness or swelling is present in the painful area
- any new symptoms appear

### If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### **Directions**

- do not take more than directed
- adults and childlren 12 years and over: take 2 caplets at bedtime
- do not take more than 2 caplets in 24 hours

#### Other information

- read all warnings and directions before use. Keep carton.
- store between 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

# Inactive ingredients

carnauba wax, corn starch, croscarmellose sodium,FD&C blue #1, hypromellose, microcrystalline cellulose, polydestrose, polyethylene glycol 400, silicon dioxide, stearic acid, titanium dioxide

### **Questions or comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

# **Principal Display Panel**

COMPARE TO THE ACTIVE INGREDIENTS IN ADVIL® PM†

IBUPROFEN PM

Ibuprofen 200 mg

Pain reliever (NSAID)

Diphenhydramine Citrate 38 mg

Nighttime sleep-aid

COATED CAPLET\*\*

(\*\*Capsules-Shaped Tablets)

†This product is not manufactured or distributed by Heleon Group of Companies, distributor Advil® PM.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY:
MIDWOOD BRAND LLC
500 VOLVO PHWY,
CHESAPEAKE, VA 23320

**Product Packaging** 



### **FAMILY WELLNESS Ibuprofen PM Caplets**

Drug Facts (continued)

Questions or comments?

Inactive ingredients carnauba wax, corn starch

microcrystalline cellulose, polydextrose, polyethylene glycol 400, silicon dioxide, stearic acid, titanium dioxide

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

This product is not manufactured or distributed by Haleon Group of Companies, distributor of Advil® PM

croscarmellose sodium, FD&C blue #2, hypromellose,

### **IBUPROFEN PM**

diphenhydramine citrate, ibuprofen tablet, coated

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-197	
Route of Administration	ORAL			

Active Ingredient/Active Moiety			
Ingredient Name	<b>Basis of Strength</b>	Strength	
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg	
<b>DIPHENHYDRAMINE CITRATE</b> (UNII: 40D433S209) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE CITRATE	38 mg	

Inactive Ingredients	
Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics				
Color	blue	Score	no score	
Shape	OVAL	Size	15mm	
Flavor		Imprint Code	PL72	
Contains				

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:55319- 197-20	1 in 1 BOX	03/01/2024			
1		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product				
2	NDC:55319- 197-40	1 in 1 BOX	03/01/2024			
2		40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product				

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA211404	03/01/2024	