

**MUCUS RELIEF COLD FLU AND SORE THROAT MAX MAXIMUM STRENGTH-  
acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine  
hcl tablet, film coated  
Rite Aid Corporation**

-----  
**Rite Aid 44-616-Delisted**

***Active ingredients (in each caplet)***

Acetaminophen 325 mg  
Dextromethorphan HBr 10 mg  
Guaifenesin 200 mg  
Phenylephrine HCl 5 mg

***Purpose***

Pain reliever/fever reducer  
Cough suppressant  
Expectorant  
Nasal decongestant

***Uses***

- temporarily relieves these common cold and flu symptoms:
  - nasal congestion
  - sore throat
  - headache
  - minor aches and pains
  - cough
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- liver disease
- diabetes
- heart disease
- difficulty in urination due to enlargement of the prostate gland
- thyroid disease
- high blood pressure
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

**When using this product**

**do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- new symptoms occur
- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do

not notice any signs or symptoms.

### ***Directions***

- **do not take more than directed**
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years and over: take 2 caplets every 4 hours
- children under 12 years: do not use

### ***Other information***

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- use by expiration date on package

### ***Inactive ingredients***

corn starch, crospovidone, FD&C blue #1 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

### ***Questions?***

**1-800-426-9391**

### ***Principal Display Panel***

NDC 11822-6160-9

**MULTI-SYMPTOM RELIEF**

**MAXIMUM STRENGTH**

**MUCUS RELIEF**

**COLD, FLU, &  
SORE THROAT MAX**

**ACETAMINOPHEN**

DEXTROMETHORPHAN HBr

GUAIFENESIN

PHENYLEPHRINE HCl

PAIN RELIEVER / FEVER REDUCER

COUGH SUPPRESSANT • EXPECTORANT

NASAL DECONGESTANT

Relieves headache, fever & sore throat

Controls cough

Relieves nasal & chest congestion

Thins & loosens mucus

For ages 12 & over

ACTUAL SIZE

**20**  
CAPLETS

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

50844

REV0419C61609

**DISTRIBUTED BY:**

RITE AID, 30 HUNTER LANE,  
CAMP HILL, PA 17011

**www.riteaid.com**

**SATISFACTION  
GUARANTEE**

**If you're not satisfied, we'll  
happily refund your money.**



Rite Aid 44-616



**MUCUS RELIEF COLD FLU AND SORE THROAT MAX MAXIMUM STRENGTH**

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, film coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-6160
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSPVIDONE</b> (UNII: 2S7830E561)	
<b>FD&amp;C BLUE NO. 1 ALUMINUM LAKE</b> (UNII: J9EQA3S2JM)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6130)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	44;616
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:11822-6160-9	2 in 1 PACKAGE	04/07/2013	01/10/2025
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/07/2013	01/10/2025

**Labeler** - Rite Aid Corporation (014578892)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(11822-6160) , pack(11822-6160)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(11822-6160)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(11822-6160)

Revised: 10/2023

Rite Aid Corporation