

ACTIVCIN- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled
OPMX LLC.

Activcin Day-Time Relief

Drug Facts

Active ingredients (in each LiquiCap)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

Temporarily relieves common cold/flu symptoms:

- Nasal congestion
- Cough due to minor throat & bronchial irritation
- Sore throat
- Headache
- Minor aches & pains
- Fever

Warnings

Liver warning

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include.

Skin reddening blisters • rash • If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription).

- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug.
- If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- Liver disease
- Heart disease
- Diabetes
- High blood pressure
- Trouble urinating due to enlarged prostate gland
- Cough that occurs with too much phlegm (mucus)
- Persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis or emphysema.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product,

do not use more than directed.

Stop use and ask a doctor if

- You get nervous, dizzy or sleepless.
- Symptoms get worse or last more than 5 days (children) or 7 days (adults).
- Fever gets worse or lasts more than 3 days.
- Redness or swelling is present.
- New symptoms occur.
- Cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well for children even if you do not notice any signs or symptoms.

Directions

- Take only as directed.
 - Do not exceed 4 doses per 24 hrs.
-

Adults & children 12 yrs & over	2 softgels with water every 4 hrs
Children 4 to under 12 yrs	Ask a doctor
Children under 4 yrs	Do not use

Other information

Store at room temperature

Inactive ingredients

FD&C Red No. 40, FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol special, titanium dioxide.

Questions?

Call to free 619-600-5632 Monday through Friday 9AM - 5PM EST

Made for:

RRX

Exclusively distributed by:

OPMX

Chula Vista, CA 91910

Phone: 619-600-5632

PRINCIPAL DISPLAY PANEL

NDC 69729-008-10

COUGH & FLU / TOS Y GRIPE

Activsin

Day-TimeRelief

Non-Drowsy Multi-Symptom Relief

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl

- Body Aches / Dolor de cuerpo
- Fever / Fiebre
- Sore throat / Dolor de cabeza
- Nasal Congestion / Congestion Nasal
- Cough / Tos

10 Softgels / 10 Cápsulas blandas de gel



Drug Facts (continued)

Active ingredients (in each Softgel) **Purpose**
 Acetaminophen 325 mg Pain reliever/fever reducer
 Dextromethorphan HBr 10 mg Cough suppressant
 Phenylephrine HCl 5 mg Nasal decongestant

Uses
 Temporary relief of symptoms of the common cold/flu such as:
 • Cough due to minor throat & bronchial irritation
 • Sore throat
 • Headache
 • Minor aches and pains
 • Fever

Warnings
 This product contains acetaminophen. Severe liver damage may occur if you take:
 • More than 4 doses in 24 hrs, which is the maximum daily amount for this product.
 • With other drugs containing acetaminophen.
 • 3 or more alcoholic drinks daily while using this product.

Warnings
 Symptoms may include:
 • Skin redness • Blisters • Rash
 • If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
 • With any other drug containing acetaminophen (prescription or nonprescription).
 • If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 • If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
 • Liver disease • Heart disease • High blood pressure
 • Trouble urinating due to enlarged prostate gland • Cough that

Questions?
 Call toll free 619-600-5632 Monday through Friday, 9AM - 5PM EST

Other information
 • Store at room temperature.
 Children under 4 yrs Do not use
 Children 4 to under 12 yrs Ask a doctor
 Adults & children 12 yrs & over 2 softgels with water every 4 hrs.
 • Take only as directed. • Do not exceed 4 doses per 24 hours.

Directions
 • Do not exceed 4 doses per 24 hours.
 • Take only as directed. • Do not exceed 4 doses per 24 hours.

Overdose warning: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Keep out of reach of children.
 If pregnant or breast-feeding, ask a health professional before use.

When using this product, do not use more than directed.
 Taking the blood thinning drug warfarin

Stop use and ask a doctor if
 • You get nervous, dizzy or sleepless. • Symptoms get worse or last more than 5 days (children) or 7 days (adults). • Fever gets worse or lasts more than 3 days. • Redness or swelling is present.
 • New symptoms occur. • Cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

Ask a doctor or pharmacist before use if you are:
 • Taking the blood thinning drug warfarin

Drug Facts (continued)

TAMPER EVIDENT:
 Do not use if package is opened or if blister unit is bent, broken or shows any sign of tampering.

Exclusively distributed by
OPIMIX
 Chula Vista, CA 91910
 Phone: 619-600-5632

Made for:
RRX

Activcin
 Day-Time Relief

COUGH & FLU TOS Y GRIPE

Activcin
 Day-Time Relief

Non-Drowsy Multi-Symptom Relief
 Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl

- Body Aches / Dolor de Cuerpo
- Fever / Fiebre
- Sore Throat / Dolor de Garganta
- Nasal Congestion / Congestión Nasal
- Cough / Tos

**10 Softgels *
 10 Cápsulas Blandas de Gel ***

*Capsules, Liquid-Filled / *Cápsulas rellenas de líquido

Lot:
 Exp:

QR Code

8 10053 18044 0
 REV 06/2022 ITEM # RRX1109

ACTIVCIN
 acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69729-008
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	AP01
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69729-008-10	1 in 1 CARTON	05/23/2022	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/23/2022	

Labeler - OPMX LLC. (029918743)

Establishment

Name	Address	ID/FEI	Business Operations
SOFTECH PHARMA PRIVATE LIMITED		677111277	label(69729-008) , manufacture(69729-008) , pack(69729-008)

Revised: 4/2024

OPMX LLC.