# XRAY DOL- camphor, menthol, and methyl salicylate cream Genomma Lab USA, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

### X Ray Pain Relief Cream 2oz 6454 ZDP

### **Active ingredients Purposes**

Camphor 4%......Topical analgesic

Menthol 10%......Topical analgesic

Methyl salicylate 30%.....Topical analgesic

### Uses

- temporarily relieves the minor aches and pains of muscles and joints associated with
- sprains
- simple backache
- arthritis
- strains
- bruises

### Warnings

### For external use only

**Allergy alert:** If prone to allergic reaction from aspirin or salicylates, consult a doctor before use.

### Do not use

- on wounds or damaged skin
- with a heating pad
- on a child under 12 years of age with arthritis-like conditions

Ask a doctor before use if you have redness over the affected area.

## When using this product

- do not use otherwise than as directed
- avoid contain with the eyes, mucous membranes or rashes
- do not bandage tightly

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away

#### **Directions**

- use only as directed
- adults and children 12 years of age and older: apply to affected area not more than 3 to 4 times daily
- children under 12 years of age: ask a doctor

### Other information

store at room temperature 20-25°C (68-77°F)

### **Inactive ingredients**

ethylparaben, glyceryl monostearate, polysorbate 80, purified water, sorbitan monooleate, stearic acid

### Distributed by

Genomma Lab USA, Inc.

Made in China







#### Drug Facts

Active ingredients Purposes Camphor 4%..... Menthol 10%..... Methyl salicylate 30% .. Topical analgesic .Topical analgesic

Uses ■ temporarily relieves the minor aches and ains of muscles and joints associated with sprains ■simple backache ■arthritis ■strains ■bruises

#### Warnings

Allergy alert: If prone to allergic reaction from aspirin or salicylates, consult a doctor before use.

Do not use ■ on wounds or damaged skin ■ with a heating pad ■ on a child under 12 years of age with arthritis-like conditions

Ask a doctor before use if you have redness over the affected area.

When using this product ■ do not use otherwise than as directed avoid contact with eyes, mucous membranes or rashes do not bandage tightly

Stop use and ask a doctor if redness is present stup use and ask a ductor in a lettiless is present or irritation or rash develops a condition worsens or symptoms persist for more than 7 days a symptoms clear up and occur again within a few days if pregnant or breast-feeding, ask a health

professional before use.

Keep out of reach of children. If swallowed, get
medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**■ use only as directed ■ adults and children 12 years of age and older: apply to affected area not more than 3 to 4 times daily ■ children under 12 years of age: ask a doctor

### Other information

tore at room temperature 20-25°C (68-77°F)

### Inactive ingredients

ethylparaben, glyceryl monostearate, polysorbate 80. purified water, sorbitan monooleate, stearic acid

Questions or comments: Call toll free 1-877-994-3666 Monday to Friday, 8:00 am - 6:00 pm CST

Made in / Hecho en China

(G)



FOR **DAILY TREATMENT** OF ARTHRITIS AND JOINT PAIN\*



**Topical Analgesic** 

\*See Drug Facts for more information

Menthol 10% Methyl salicylate 30% NET WT. 2 OZ (56.7 G)

#### Información del Medicamento

Ingredientes activos Propósitos licilato de metilo 30% ..

Usos ■ alivia temporalmente los dolores y molestias menores de músculos y articulaciones asociados con: ■ esguinces ■ dolor de espalda ■ artritis ■ distensiones ■ contusiones

olo para uso externo

médico antes de usar. No lo use ■ en heridas o piel dañada ■ con una

almohadilla térmica ■ en niños menores de 12 años

con condiciones similares a la artritis

Consulte a un médico antes de usar si tiene projecimiento en el área afectada.

Cuando use este producto ■ no lo use de otra manera que no sea como se indica ■ evite el contacto con los ojos, las membranas mucosas o las erupciones ■ no use vendajes apretados

Deje de usarlo y consulte a un médico si ■ hay enrojecimiento, irritación o sarpullido ■ la condición empeora o los síntomas persisten durante más de 7 días ■ los síntomas desaparecen y vuelven a aparecer en unos pocos días

Si está embarazada o amamantando, consulte a un profesional de la salud antes de usar. Mantener fuera del alcance de los niños. Si se ingiere, obtenga ayuda médica o comuniquese con un Centro de control de envenenamiento (1-800-222-1222) de inmediato.

Instrucciones ■ use solo como se indica ■ adultos y niños mayores de 12 años: aplique en el área afectada no más de 3 a 4 veces al día ■ niños nores de 12 años: consulte a un médico

**Otra información** almacene a temperatura ambiente 20-25°C (68-77°F)

#### Ingredientes inactivos

etilparabeno, monoestearato de glicerilo, polisorbato 80, agua purificada, monooleato de sorbitán, ácido

¿Preguntas o comentarios? Llame lada sin costo **1-877-994-3666** Lunes a viernes, 8:00 am – 6:00 pm CST

2000011966



Mentol 10%

Salicilato de metilo 30%

### **XRAY DOL**

camphor, menthol, and methyl salicylate cream

\*Lea la Información de Medicamento

NET WT. 2 OZ (56.7 G)

#### **Product Information**

**Product Type** HUMAN OTC DRUG Item Code (Source) NDC:50066-454

Route			

TOPICAL

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
<b>CAMPHOR (SYNTHETIC)</b> (UNII: 5TJD82A1ET) (CAMPHOR (SYNTHETIC) - UNII:5TJD82A1ET)	CAMPHOR (SYNTHETIC)	40 mg in 1 g		
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	100 mg in 1 g		

Inactive Ingredients			
Ingredient Name	Strength		
METHYL SALICYLATE (UNII: LAV5U5022Y)	300 mg in 1 g		
ETHYLPARABEN (UNII: 14255EXE39)			
GLYCERYL MONOSTEARATE (UNII: 2300U9XXE4)			
POLYSORBATE 80 (UNII: 60ZP39ZG8H)			
WATER (UNII: 059QF0KO0R)			
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)			
STEARIC ACID (UNII: 4ELV7Z65AP)			

ı	Packaging						
7	# Item Code	Package Description	Marketing Start Date	Marketing End Date			
:	NDC:50066-454- 56	1 in 1 CARTON	06/02/2022				
:	L	56.7 g in 1 TUBE; Type 0: Not a Combination Product					

Marketing Information					
Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
art348	06/02/2022				
	Application Number or Monograph Citation	Application Number or Monograph Marketing Start Citation Date			

## Labeler - Genomma Lab USA, Inc. (832323534)

Revised: 6/2022 Genomma Lab USA, Inc.