

**EPHED PLUS LIQUID- ephedrine guaifenesin liquid**  
**Dickey Consumer Products DBA DMD**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

Warnings

Purpose: Bronchodilator, Expectorant

Uses for temporary relief of mild symptoms of intermittent asthma: wheezing tightness of chest, shortness of breath, helps loosen phlegm (mucus) and thin bronchial secretions to rid bronchial passageways of bothersome mucus, and to make coughs more productive

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

If pregnant or breast-feeding, ask a health professional before use

Directions do not take more than directed

adults and children 12 years of age and over: take 1

bottle every 4 hours as needed. Do not take more than

6 bottles in 24 hours children under 12 years of age:

do not use

Inactive ingredients Blue #1, Citric Acid, Filtered Water, Malic Acid, Natural and Artificial Flavors, Potassium Sorbate, Propylene Glycol, Sodium

Ephedrine HCl 25 mg, Bronchodilator  
Guaifenesin 400 mg, Expectorant

TAMPER EVIDENT: Do not use  
if bottle sleeve is torn or cut  
TWIST OFF



**Drug Facts**

**Active ingredients (in each bottle)** . . . . . **Purpose**  
 Ephedrine HCl, 25 mg . . . . . Bronchodilator  
 Guaifenesin, USP 400 mg . . . . . Expectorant

**Uses** ■ for temporary relief of mild symptoms of intermittent asthma: ■ wheezing  
 ■ tightness of chest ■ shortness of breath ■ helps loosen phlegm (mucus) and thin  
 bronchial secretions to rid bronchial passageways of bothersome mucus, and to make  
 coughs more productive

**Warnings**  
**Asthma alert:** Because asthma may be life threatening, see a doctor if you are not  
 better in 60 minutes ■ get worse ■ need more than 6 bottles in 24 hours ■ use more  
 than 4 bottles in 24 hours for 3 or more days a week ■ have more than 2 asthma attacks  
 in a week. These may be signs that your asthma is getting worse ■ this product will not  
 give you asthma relief as quickly as an inhaled bronchodilator

**Do not use** ■ unless a doctor said you have asthma ■ if you are now taking a  
 prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression,  
 psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after  
 stopping the MAOI drug. If you do not know if your prescription drug contains an  
 MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have** ■ ever been hospitalized for asthma ■ heart  
 disease ■ high blood pressure ■ diabetes ■ thyroid disease ■ seizures ■ narrow  
 angle glaucoma ■ a psychiatric or emotional condition ■ trouble urinating due to an  
 enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ cough that  
 lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Ask a doctor or pharmacist before use if you are** taking prescription drugs for asthma,  
 obesity, weight control, depression, or psychiatric or emotional conditions ■ taking any  
 drug that contains phenylephrine, pseudoephedrine, ephedrine or caffeine (such as for  
 allergy, cough-cold or pain)

**NON-DROWSY**

# Ephed Plus®

Ephedrine HCl 25 mg, Bronchodilator  
 Guaifenesin 400 mg, Expectorant

**TEMPORARY RELIEF OF MILD  
 INTERMITTENT ASTHMA SYMPTOMS**

natural and artificial  
**GRAPE FLAVOR**

**2 FL OZ  
 (60ML)**

Marketed by:  
**DMD Pharmaceuticals®**  
 A Division of Dickey Consumer Products, Inc.  
 Noblesville, IN 46060 • 800-795-2477 • dmdpharm.com

**Drug Facts (continued)**

**When using this product** ■ your blood pressure or heart rate may go up.  
 This could increase your risk of heart attack or stroke, which may cause  
 death. ■ your risk of heart attack or stroke increases if you: ■ have a  
 history of high blood pressure or heart disease ■ take this product more  
 frequently or take more than the recommended dose ■ avoid foods or  
 beverages that contain caffeine ■ avoid dietary supplements containing  
 ingredients reported or claimed to have a stimulant effect ■ intentional  
 abuse of this product can be harmful or fatal

**Stop use and ask a doctor if** ■ your asthma is getting worse (see Asthma  
 alert) ■ you have difficulty sleeping ■ you have a rapid heartbeat ■ you  
 have tremors, nervousness or seizure ■ cough lasts for more than 7 days,  
 comes back, or occurs with fever, rash, or persistent headache. These could  
 be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health  
 professional before use

**Keep out of reach of children.** In case of overdose, get  
 medical help or contact a Poison Control Center right away.

**Directions** do not take more than directed  
 ■ adults and children 12 years of age and over: take 1  
 bottle every 4 hours as needed. Do not take more than  
 6 bottles in 24 hours ■ children under 12 years of age:  
 do not use

**Other information** no refrigeration necessary.

**Inactive ingredients** Blue #1, Citric Acid,  
 Filtered Water, Malic Acid, Natural and Artificial Flavors,  
 Potassium Sorbate, Propylene Glycol, Sodium  
 Benzoate, Sucralose, Red #40.



## EPHED PLUS LIQUID

ephedrine guaifenesin liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:65193-180
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
EPHEDRINE HYDRO CHLORIDE (UNII: NLJ6390P1Z) (EPHEDRINE - UNII:GN83C131XS)	EPHEDRINE HYDROCHLORIDE	25 mg in 60 mL
guaifenesin (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	guaifenesin	400 mg in 60 mL

Inactive Ingredients	
Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)	

<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>MALIC ACID</b> (UNII: 817L1N4CKP)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	

### Product Characteristics

<b>Color</b>	purple (purple)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	grape	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65193-180-57	60 mL in 1 BOTTLE; Type 0: Not a Combination Product	11/10/2017	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	11/10/2017	

**Labeler** - Dickey Consumer Products DBA DMD (835974056)

**Registrant** - Dickey Consumer Products DBA DMD (835974056)

### Establishment

Name	Address	ID/FEI	Business Operations
Ultraseal Corporation		085752004	pack(65193-180)

### Establishment

Name	Address	ID/FEI	Business Operations
ULTRAtab Laboratories, Inc.		151051757	manufacture(65193-180)