

SPECTOGARD WATER SOLUBLE POWDER- spectinomycin powder, for solution
Bimeda, Inc.

SpectoGard® Water Soluble Powder
(spectinomycin)

FOR GROWING CHICKENS

Anti-infective

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

NOT FOR USE IN HUMANS

KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California) - Use only as directed

Approved by FDA under NADA # 038-661

INDICATIONS, DOSAGE AND ADMINISTRATION:

As an aid in the prevention or control of losses due to CRD associated with *M. gallisepticum* (PPLO). Administer two (2) grams per gallon of water as the only source of drinking water for the first three days of life and for one day following each vaccination.

As an aid in controlling infectious synovitis due to *Mycoplasma synoviae* in broiler chickens. Administer one (1) gram per gallon of water as the only source of drinking water for the first three to five days of life.

DIRECTIONS FOR USE:

To provide 2 g/gal: Dissolve contents in 250 gallons of water. For proportioners, dissolve contents of 1/2 package in 1 gallon of water. Meter at 1 oz/gal.

To provide 1 g/gal: Dissolve contents in 500 gallons of water. For proportioners, dissolve contents of 1/2 package in 2 gallons of water. Meter at 1 oz/gal.

RESIDUE WARNING: DO NOT administer this drug within 5 days of slaughter. DO NOT administer to laying chickens.

PRECAUTION:

Rarely, some individuals who handle spectinomycin develop serious reactions involving skin, nails and eyes. Individuals who have experienced rash or other evidence of allergic reaction should avoid further contact with spectinomycin.

The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report suspected adverse drug events, for technical assistance or to obtain a copy of the SDS, contact Bimeda, Inc. at 1888-524-6332. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at www.fda.gov/reportanimalae.

Store at controlled room temperature 20°C - 25°C (68°F - 77°F), with excursions permitted 15°C - 30°C (59°F - 86°F).

INDICATIONS, DOSAGE AND ADMINISTRATION:

As an aid in the prevention or control of losses due to CRD associated with *M. gallisepticum* (PPLD). Administer two (2) grams per gallon of water as the only source of drinking water for the first three days of life and for one day following each vaccination.

As an aid in controlling infectious synovitis due to *Mycoplasma synoviae* in broiler chickens. Administer one (1) gram per gallon of water as the only source of drinking water for the first three to five days of life.

DIRECTIONS FOR USE:

To provide 2 g/gal: Dissolve contents in 250 gallons of water. For proportioners, dissolve contents of 1/2 package in 1 gallon of water. Meter at 1 oz/gal.

To provide 1 g/gal: Dissolve contents in 500 gallons of water. For proportioners, dissolve contents of 1/2 package in 2 gallons of water. Meter at 1 oz/gal.

Manufactured by:
Bimeda, Inc.
Le Sueur, MN 56058
www.bimeda.com

Lot:
Exp:



SpectoGard® Water Soluble Powder (spectinomycin)

FOR GROWING CHICKENS

Anti-infective

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**NOT FOR USE IN HUMANS
KEEP OUT OF REACH OF CHILDREN**

Restricted Drug (California) - Use Only as Directed
Approved by FDA under NADA # 038-661

Each 1000 g contains: Spectinomycin dihydrochloride pentahydrate equivalent to spectinomycin activity 500.00 g

RESIDUE WARNING: DO NOT administer this drug within 5 days of slaughter. DO NOT administer to laying chickens.

PRECAUTION:

Rarely, some individuals who handle spectinomycin develop serious reactions involving skin, nails and eyes. Individuals who have experienced rash or other evidence of allergic reaction should avoid further contact with spectinomycin.

The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report suspected adverse drug events, for technical assistance or to obtain a copy of the SDS, contact Bimeda, Inc. at 1-888-524-6332. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at www.fda.gov/reportanimalae.

Store at controlled room temperature 20°C - 25°C (68°F - 77°F), with excursions permitted to 15°C - 30°C (59°F - 86°F).

SpectoGard® is a Registered Trademark of Bimeda, Inc.
N.A. Corp. Address: Bimeda, Inc., One Tower Lane, Oakbrook Terrace, IL 60181



Net Contents:
1000 g (35.27 oz)



SPECTOGARD WATER SOLUBLE POWDER

spectinomycin powder, for solution

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:61133-5485
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SPECTINOMYCIN (UNII: 93AKI1U6QF) (SPECTINOMYCIN - UNII:93AKI1U6QF)	SPECTINOMYCIN	500 g in 1000 g

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61133-5485-1	1000 g in 1 CANISTER		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA038661	06/14/2016	

Labeler - Bimeda, Inc. (060492923)

Establishment

Name	Address	ID/FEI	Business Operations
Bimeda, Inc.		060492923	manufacture

Revised: 12/2023

Bimeda, Inc.