

**HEADACHE RELIEF- acetaminophen, aspirin, caffeine tablet, film coated  
DOLGENCORP, LLC**

-----  
**Dollar General 44-334B**

***Active ingredients (in each caplet)***

Acetaminophen 250 mg

Aspirin 250 mg (NSAID\*)

Caffeine 65 mg

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

Pain reliever

Pain reliever aid

***Uses***

- temporarily relieves minor aches and pains due to:
  - headache
  - arthritis
  - a cold
  - muscular aches
  - toothache
  - premenstrual and menstrual cramps

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- shock
- facial swelling
- hives
- asthma (wheezing)

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- 3 or more alcoholic drinks every day while using this product
- with other drugs containing acetaminophen

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

#### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

#### **Ask a doctor before use if**

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

#### **Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout, or arthritis

#### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - have bloody or black stools
  - vomit blood
  - have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- redness or swelling is present

- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms appear

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours. Do not take more than 8 caplets in 24 hours.
- children under 12 years: ask a doctor

### **Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

### **Inactive ingredients**

corn starch, croscopvidone, hypromellose, microcrystalline cellulose, povidone, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

### **Questions or comments?**

**1-888-309-9030**

### **Principal display panel**

**DG®** | health

Compare to  
active ingredients  
of Excedrin® Extra  
Strength Headache\*\*

**Extra Strength  
Headache  
Relief**

**Acetaminophen,  
Aspirin (NSAID),  
and Caffeine**

Pain Reliever / Pain Reliever Aid

Actual Caplet Size

**100** Caplets

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

\*\*This product is not manufactured or distributed by Haleon CH  
SARL, owner of the registered trademark Excedrin® Extra  
Strength Headache. 50844 ORG022433412

DISTRIBUTED BY  
OLD EAST MAIN CO.  
100 MISSION RIDGE  
GOODLETSVILLE, TN 37072

**100%**  
**Satisfaction**  
**Guaranteed!**  
(888)309-9030

No Print Area  
Lot no. & Exp. Date

DISTRIBUTED BY  
OLD EAST MAIN CO.  
100 MISSION RIDGE  
GOODLETTSVILLE, TN 37072

**100%**  
Satisfaction  
Guaranteed!  
(888) 309-9030

A0545



Compare to active ingredients of Excedrin® Extra Strength Headache\*\*



# Extra Strength Headache Relief

Acetaminophen, Aspirin (NSAID), and Caffeine  
Pain Reliever / Pain Reliever Aid



100 Caplets  
Actual Caplet Size



Compare to active ingredients of Excedrin® Extra Strength Headache\*\*

# Extra Strength Headache Relief

Acetaminophen, Aspirin (NSAID) and Caffeine



100 Caplets  
Actual Caplet Size

**Drug Facts (continued)**

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

**Inactive ingredients**

corn starch, croscopolone, hypromellose, microcrystalline cellulose, povidone, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

**Questions or comments?** 1-888-309-9030

\*\*This product is not manufactured or distributed by Haleon CH SARL, owner of the registered trademark Excedrin® Extra Strength Headache. 50844 ORG022433412

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**Drug Facts (continued)**

**Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if**

- you have liver disease
- stomach bleeding/ulcers or you have ulcers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout, or arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling is present
- any new symptoms appear

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use aspirin in 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours. Do not take more than 8 caplets in 24 hours.
- children under 12 years: ask a doctor

B-0315-334B-12  
ORG022433412

**KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

**Drug Facts**

**Active ingredients (in each caplet)**

Acetaminophen 250 mg.....Pain reliever  
Aspirin 250 mg (NSAID).....Pain reliever  
Caffeine 65 mg.....Pain reliever aid  
\*nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - toothache
  - arthritis
  - a cold
  - headache
  - muscular aches
  - premenstrual and menstrual cramps

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- shock
- facial swelling
- hives
- asthma (wheezing)

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- take more or for a longer time than directed
- have 2 or more alcoholic drinks every day while using this product

Take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness.

Dollar General 44-334

## HEADACHE RELIEF

acetaminophen, aspirin, caffeine tablet, film coated

Product Information		
Product Type	HUMAN OTC DRUG	Item Code (Source)
Route of Administration	ORAL	NDC:55910-345

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

**Product Characteristics**

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	44;334
<b>Contains</b>			

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:55910-345-08	1 in 1 CARTON	05/06/2021	
1		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:55910-345-12	1 in 1 CARTON	05/06/2021	
2		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M013	05/06/2021	

**Labeler** - DOLGENCORP, LLC (068331990)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(55910-345)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(55910-345)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(55910-345)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(55910-345)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(55910-345)

Revised: 6/2024

DOLGENCORP, LLC