

LOPERAMIDE HYDROCHLORIDE AND SIMETHICONE - loperamide hydrochloride and simethicone tablet
Camber Consumer Care Inc

Loperamide Hydrochloride and Simethicone Tablets, 2 mg/125 mg (OTC)

ACTIVE INGREDIENT(S)

(in each tablet)

Loperamide Hydrochloride USP 2 mg

Simethicone USP 125 mg

PURPOSES

Anti-diarrheal

Anti-gas

USE(S)

relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

WARNINGS

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide hydrochloride

Heart alert: Taking more than directed can cause serious heart problems or death

DO NOT USE

- if you have bloody or black stool
- if you have difficulty swallowing

ASK A DOCTOR BEFORE USE IF YOU HAVE

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

ASK A DOCTOR OR PHARMACIST BEFORE USE IF

you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

WHEN USING THIS PRODUCT

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

STOP USE AND ASK A DOCTOR IF

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- you have difficulty swallowing the tablet

If pregnant or breast-feeding, ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

DIRECTIONS

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- take only on an empty stomach (1 hour before or 2 hours after a meal)
- take with a full (8 oz.) glass of water
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 tablets after the first loose stool; 1 tablet after each subsequent loose stool; but no more than 4 tablets in 24 hours
children 9-11 years (60-95 lbs)	1 tablet after the first loose stool; 1/2 tablet after each subsequent loose stool; but no more than 3 tablets in 24 hours
children 6-8 years (48-59 lbs)	1 tablet after the first loose stool; 1/2 tablet after each subsequent loose stool; but no more than 2 tablets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

OTHER INFORMATION

- each tablet contains: **calcium 640 mg, very low sodium and potassium 5.838 mg**
- store between 20 to 25°C (68 to 77°F). Protect from light.

- **do not use if carton is open or if blister unit is open or torn**

INACTIVE INGREDIENTS

acesulfame potassium, croscarmellose sodium, dibasic calcium phosphate, microcrystalline cellulose, stearic acid and vanilla flavor

QUESTIONS OR COMMENTS

call **1-888-588-1418**

Distributed by:

Camber Consumer Care, Inc.

Piscataway, NJ 08854, USA.

PRINCIPAL DISPLAY PANEL

Loperamide Hydrochloride and Simethicone Tablets 2 mg/125 mg-1x6's carton

Drug Facts (continued)

Other information

- each tablet contains: calcium 640 mg, very low sodium and potassium 5.838 mg.
- store between 20 to 25°C (68 to 77°F). Protect from light.
- do not use if carton is open or if blister unit is open or torn

Drug Facts (continued)

Inactive ingredients
acesulfame potassium, croscarmellose sodium, dibasic calcium phosphate, microcrystalline cellulose, stearic acid and vanilla flavor

Questions or comments? call 1-888-588-1418

*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Imodium® Multi-Symptom Relief.



NDC 69230-325-06
*Compare to the active ingredients in Imodium® Multi-Symptom Relief.

Loperamide Hydrochloride and Simethicone Tablets
2 mg/125 mg
Antidiarrheal/Anti-gas

Multi-Symptom Relief
Relieves symptoms of diarrhea plus

- Cramps & Pressure
- Bloating
- Gas

Actual Size

6 (1x6) Unit-Dose Tablets

Un varnish Area for Batch Coding
70(L)x 28(H) mm

Drug Facts

Active ingredients (in each tablet)	Purposes
Loperamide Hydrochloride USP 2 mg.....	Anti-diarrheal
Simethicone USP 125 mg.....	Anti-gas

Uses relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

Warnings Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide hydrochloride



Mfg. Lot. No.: 24M07S307678
Distributed by:
Camber Consumer Care, Inc.
Piscataway, NJ 08854, USA



Drug Facts (continued)

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

- if you have bloody or black stool
- if you have difficulty swallowing

Ask a doctor before use if you have

- fever
- a history of liver disease
- mucus in the stool
- a history of an abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- diarrhea lasts for more than 2 days
- you have difficulty swallowing the tablet

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
■ drink plenty of clear fluids to help prevent dehydration

Drug Facts (continued)

caused by diarrhea

- take only on an empty stomach (1 hour before or 2 hours after a meal)
- take with a full (8 oz.) glass of water
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 tablets after the first loose stool; 1 tablet after each subsequent loose stool; but no more than 4 tablets in 24 hours
children 9-11 years (60-95 lbs)	1 tablet after the first loose stool; 1/2 tablet after each subsequent loose stool; but no more than 3 tablets in 24 hours
children 6-8 years (48-59 lbs)	1 tablet after the first loose stool; 1/2 tablet after each subsequent loose stool; but no more than 2 tablets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

LOPERAMIDE HYDROCHLORIDE AND SIMETHICONE

loperamide hydrochloride and simethicone tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69230-325
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	125 mg
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CALCIUM PHOSPHATE, DIBASIC, ANHYDROUS (UNII: L11K75P92J)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	2 pieces
Shape	CAPSULE	Size	17mm
Flavor	VANILLA	Imprint Code	H;L21
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69230-325-04	1 in 1 CARTON	06/17/2021	
1	NDC:69230-325-31	4 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:69230-325-08	2 in 1 CARTON	06/17/2021	
2	NDC:69230-325-31	4 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:69230-325-06	1 in 1 CARTON	06/17/2021	
3	NDC:69230-325-32	6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
4	NDC:69230-325-12	2 in 1 CARTON	06/17/2021	
4	NDC:69230-325-32	6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
5	NDC:69230-325-18	3 in 1 CARTON	06/17/2021	
5	NDC:69230-325-32	6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
6	NDC:69230-325-24	4 in 1 CARTON	06/17/2021	
6	NDC:69230-325-32	6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA211438	06/17/2021	

Labeler - Camber Consumer Care Inc (079539968)

Establishment

Name	Address	ID/FEI	Business Operations
Annora Pharma Private Limited		650980746	analysis(69230-325) , manufacture(69230-325)

Revised: 6/2021

Camber Consumer Care Inc