

# **MOXIFLOXACIN OPHTHALMIC SOLUTION - moxifloxacin ophthalmic solution/**

**drops**

**A-S Medication Solutions**

---

## **HIGHLIGHTS OF PRESCRIBING INFORMATION**

**These highlights do not include all the information needed to use MOXIFLOXACIN OPHTHALMIC SOLUTION safely and effectively. See full prescribing information for MOXIFLOXACIN OPHTHALMIC SOLUTION.**

### **MOXIFLOXACIN ophthalmic solution, for topical ophthalmic use**

**Initial U.S. Approval: 1999**

---

#### **INDICATIONS AND USAGE**

MOXIFLOXACIN ophthalmic solution is a topical fluoroquinolone anti- infective indicated for the treatment of bacterial conjunctivitis caused by susceptible strains of the following organisms:

*Corynebacterium species\**, *Micrococcus luteus\**, *Staphylococcus aureus*,  
*Staphylococcus epidermidis*, *Staphylococcus haemolyticus*, *Staphylococcus hominis*,  
*Staphylococcus warneri\**, *Streptococcus pneumoniae*, *Streptococcus viridans group*,  
*Acinetobacter lwoffii\**, *Haemophilus influenzae*, *Haemophilus parainfluenzae\**  
*Chlamydia trachomatis*

\*Efficacy for this organism was studied in fewer than 10 infections. (1)

---

#### **DOSAGE AND ADMINISTRATION**

Instill one drop in the affected eye 3 times a day for 7 days. (2)

---

#### **DOSAGE FORMS AND STRENGTHS**

Ophthalmic solution containing moxifloxacin 0.5%. (3)

---

#### **CONTRAINDICATIONS**

MOXIFLOXACIN ophthalmic solution is contraindicated in patients with a history of hypersensitivity to MOXIFLOXACIN, to other quinolones, or to any of the components in this medication. (4)

---

#### **WARNINGS AND PRECAUTIONS**

- Hypersensitivity Reactions: Hypersensitivity and anaphylaxis have been reported with systemic use of moxifloxacin. (5.1)
- Prolonged use: May result in overgrowth of non- susceptible organisms, including fungi. If superinfection occurs, discontinue use and institute alternative therapy. (5.2)
- Avoid Contact Lens Wear: Patients should not wear contact lenses if they have signs or symptoms of bacterial conjunctivitis. (5.3)

---

#### **ADVERSE REACTIONS**

The most frequently reported ocular adverse events were conjunctivitis, decreased visual acuity, dry eye, keratitis, ocular discomfort, ocular hyperemia, ocular pain, ocular pruritus, subconjunctival hemorrhage, and tearing. These events occurred in approximately 1-6% of patients.

**To report SUSPECTED ADVERSE REACTIONS, contact Alembic Pharmaceutical, Inc. at 1-866-210-9797 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

**See 17 for PATIENT COUNSELING INFORMATION.**

**Revised: 12/2023**

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **1 INDICATIONS AND USAGE**

### **2 DOSAGE AND ADMINISTRATION**

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

## **5 WARNINGS AND PRECAUTIONS**

- 5.1 Hypersensitivity Reactions
- 5.2 Growth of Resistant Organisms with Prolonged Use
- 5.3 Avoidance of Contact Lens Wear

## **6 ADVERSE REACTIONS**

## **7 DRUG INTERACTIONS**

## **8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics
- 12.4 Microbiology

## **13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis and Mutagenesis and Impairment of Fertility

## **14 CLINICAL STUDIES**

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

## **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

Moxifloxacin ophthalmic solution is indicated for the treatment of bacterial conjunctivitis caused by susceptible strains of the following organisms:

*Corynebacterium species*\*

*Micrococcus luteus*\*

*Staphylococcus aureus*

*Staphylococcus epidermidis*

*Staphylococcus haemolyticus*

*Staphylococcus hominis*

*Staphylococcus warneri*\*

*Streptococcus pneumoniae*

*Streptococcus viridans group*

*Acinetobacter lwoffii*\*

*Haemophilus influenzae*

*Haemophilus parainfluenzae*\*

*Chlamydia trachomatis*

\*Efficacy for this organism was studied in fewer than 10 infections.

## **2 DOSAGE AND ADMINISTRATION**

Instill one drop in the affected eye 3 times a day for 7 days. Moxifloxacin ophthalmic solution is for topical ophthalmic use.

## **3 DOSAGE FORMS AND STRENGTHS**

Ophthalmic solution containing moxifloxacin 0.5%.

## **4 CONTRAINDICATIONS**

Moxifloxacin ophthalmic solution is contraindicated in patients with a history of hypersensitivity to moxifloxacin, to other quinolones, or to any of the components in this medication.

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Hypersensitivity Reactions**

In patients receiving systemically administered quinolones, including moxifloxacin, serious and occasionally fatal hypersensitivity (anaphylactic) reactions have been reported, some following the first dose. Some reactions were accompanied by cardiovascular collapse, loss of consciousness, angioedema (including laryngeal, pharyngeal or facial edema), airway obstruction, dyspnea, urticaria, and itching. If an allergic reaction to moxifloxacin occurs, discontinue use of the drug. Serious acute hypersensitivity reactions may require immediate emergency treatment. Oxygen and airway management should be administered as clinically indicated.

### **5.2 Growth of Resistant Organisms with Prolonged Use**

As with other anti-infectives, prolonged use may result in overgrowth of non-susceptible organisms, including fungi. If superinfection occurs, discontinue use and institute alternative therapy. Whenever clinical judgment dictates, the patient should be examined with the aid of magnification, such as slit-lamp biomicroscopy, and, where appropriate, fluorescein staining.

### **5.3 Avoidance of Contact Lens Wear**

Patients should be advised not to wear contact lenses if they have signs or symptoms of bacterial conjunctivitis.

## **6 ADVERSE REACTIONS**

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to the rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The most frequently reported ocular adverse events were conjunctivitis, decreased visual acuity, dry eye, keratitis, ocular discomfort, ocular hyperemia, ocular pain, ocular pruritus, subconjunctival hemorrhage, and tearing. These events occurred in approximately 1%-6% of patients.

Nonocular adverse events reported at a rate of 1%-4% were fever, increased cough, infection, otitis media, pharyngitis, rash, and rhinitis.

## **7 DRUG INTERACTIONS**

Drug-drug interaction studies have not been conducted with moxifloxacin ophthalmic solution. *In vitro* studies indicate that moxifloxacin does not inhibit CYP3A4, CYP2D6, CYP2C9, CYP2C19, or CYP1A2, indicating that moxifloxacin is unlikely to alter the pharmacokinetics of drugs metabolized by these cytochrome P450 isozymes.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### Risk Summary

There are no adequate and well-controlled studies with moxifloxacin ophthalmic solution in pregnant women to inform any drug-associated risks.

Oral administration of moxifloxacin to pregnant rats and monkeys and intravenously to pregnant rabbits during the period of organogenesis did not produce adverse maternal or fetal effects at clinically relevant doses. Oral administration of moxifloxacin to pregnant rats during late gestation through lactation did not produce adverse maternal, fetal or neonatal effects at clinically relevant doses [see Data].

#### Data

##### *Animal Data*

Embryo-fetal studies were conducted in pregnant rats administered with 20, 100 or 500 mg/kg/day moxifloxacin by oral gavage on Gestation Days 6 to 17, to target the period of organogenesis. Decreased fetal body weight and delayed skeletal development were observed at 500 mg/kg/day (277 times the human AUC at the recommended human ophthalmic dose). The No-Observed-Adverse-Effect-Level (NOAEL) for developmental toxicity was 100 mg/kg/day (30 times the human AUC at the recommended human ophthalmic dose).

Embryo-fetal studies were conducted in pregnant rabbits administered with 2, 6.5 or 20 mg/kg/day moxifloxacin by intravenous administration on Gestation Days 6 to 20, to target the period of organogenesis. Abortions, increased incidence of fetal malformations, delayed fetal skeletal ossification, and reduced placental and fetal body weights were observed at 20 mg/kg/day (1086 times the human AUC at the recommended human ophthalmic dose), a dose that produced maternal body weight loss and death. The NOAEL for developmental toxicity was 6.5 mg/kg/day (246 times the human AUC at the recommended human ophthalmic dose).

Pregnant cynomolgus monkeys were administered moxifloxacin at doses of 10, 30 or

100 mg/kg/day by intragastric intubation between Gestation Days 20 and 50, targeting the period of organogenesis. At the maternal toxic doses of  $\geq$  30 mg/kg/day, increased abortion, vomiting and diarrhea were observed. Smaller fetuses/reduced fetal body weights were observed at 100 mg/kg/day (2864 times the human AUC at the recommended human ophthalmic dose). The NOAEL for fetal toxicity was 10 mg/kg/day (174 times the human AUC at the recommended human ophthalmic dose).

In a pre and postnatal study, rats were administered moxifloxacin by oral gavage at doses of 20, 100 and 500 mg/kg/day from Gestation Day 6 until the end of lactation. Maternal death occurred during gestation at 500 mg/kg/day. Slight increases in the duration of pregnancy, reduced pup birth weight, and decreased prenatal and neonatal survival were observed at 500 mg/kg/day (estimated 277 times the human AUC at the recommended human ophthalmic dose). The NOAEL for pre- and postnatal development was 100 mg/kg/day (estimated 30 times the human AUC at the recommended human ophthalmic dose).

## **8.2 Lactation**

### Risk Summary

There is no data regarding the presence of moxifloxacin ophthalmic solution in human milk, the effects on the breastfed infants, or the effects on milk production/excretion to inform risk of moxifloxacin ophthalmic solution to an infant during lactation.

A study in lactating rats has shown transfer of moxifloxacin into milk following oral administration.

Systemic levels of moxifloxacin following topical ocular administration are low [see Clinical Pharmacology (12.3)], and it is not known whether measurable levels of moxifloxacin would be present in maternal milk following topical ocular administration. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for moxifloxacin ophthalmic solution and any potential adverse effects on the breastfed child from moxifloxacin ophthalmic solution.

## **8.4 Pediatric Use**

The safety and effectiveness of moxifloxacin ophthalmic solution 0.5% have been established in all ages. Use of moxifloxacin ophthalmic solution is supported by evidence from adequate and well controlled studies of moxifloxacin ophthalmic solution in adults, children, and neonates [see *Clinical Studies (14)*].

There is no evidence that the ophthalmic administration of moxifloxacin ophthalmic solution has any effect on weight bearing joints, even though oral administration of some quinolones has been shown to cause arthropathy in immature animals.


## **8.5 Geriatric Use**

No overall differences in safety and effectiveness have been observed between elderly and younger patients.

## **11 DESCRIPTION**

Moxifloxacin ophthalmic solution USP 0.5% is a sterile solution for topical ophthalmic

use. Moxifloxacin hydrochloride is an 8-methoxy fluoroquinolone anti-infective, with a diazabicyclononyl ring at the C7 position.



Mol Wt 437.9

Chemical Name: 1-Cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-[(4aS,7aS)-octahydro-6H-pyrrolo[3,4 b]pyridin-6-yl]-4-oxo-3-quinolinecarboxylic acid, monohydrochloride. Moxifloxacin hydrochloride is a slightly yellow to yellow powder or crystals. Each mL of moxifloxacin ophthalmic solution USP contains 5.45 mg moxifloxacin hydrochloride, equivalent to 5 mg moxifloxacin base.

**Contains:** Active: Moxifloxacin 0.5% (5 mg/mL); **Inactives:** Boric acid, sodium chloride, and water for injection. May also contain hydrochloric acid/sodium hydroxide to adjust pH to approximately 6.8.

Moxifloxacin ophthalmic solution USP is an isotonic solution with an osmolality of approximately 290 mOsm/kg.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Moxifloxacin is a member of the fluoroquinolone class of anti-infective drugs (See 12.4 *Microbiology*).

### 12.3 Pharmacokinetics

Plasma concentrations of moxifloxacin were measured in healthy adult male and female subjects who received bilateral topical ocular doses of moxifloxacin ophthalmic solution 3 times a day. The mean steady-state C<sub>max</sub> (2.7 ng/mL) and estimated daily exposure

AUC (41.9 ng•hr/mL) values were 1600 and 1100 times lower than the mean  $C_{max}$  and AUC reported after therapeutic 400 mg doses of moxifloxacin. The plasma half-life of moxifloxacin was estimated to be 13 hours.

## 12.4 Microbiology

The antibacterial action of moxifloxacin results from inhibition of the topoisomerase II (DNA gyrase) and topoisomerase IV. DNA gyrase is an essential enzyme that is involved in the replication, transcription and repair of bacterial DNA. Topoisomerase IV is an enzyme known to play a key role in the partitioning of the chromosomal DNA during bacterial cell division.

The mechanism of action for quinolones, including moxifloxacin, is different from that of macrolides, aminoglycosides, or tetracyclines. Therefore, moxifloxacin may be active against pathogens that are resistant to these antibiotics and these antibiotics may be active against pathogens that are resistant to moxifloxacin. There is no cross-resistance between moxifloxacin and the aforementioned classes of antibiotics. Cross resistance has been observed between systemic moxifloxacin and some other quinolones.

*In vitro* resistance to moxifloxacin develops via multiple-step mutations. Resistance to moxifloxacin occurs *in vitro* at a general frequency of between  $1.8 \times 10^{-9}$  to less than  $1 \times 10^{-11}$  for Gram-positive bacteria.

Moxifloxacin has been shown to be active against most strains of the following microorganisms, both *in vitro* and in clinical infections as described in the Indications and Usage section:

### **Aerobic Gram-positive microorganisms:**

*Corynebacterium species*\*

*Micrococcus luteus*\*

*Staphylococcus aureus*

*Staphylococcus epidermidis*

*Staphylococcus haemolyticus*

*Staphylococcus hominis*

*Staphylococcus warneri*\*

*Streptococcus pneumoniae*

*Streptococcus viridans group*

### **Aerobic Gram-negative microorganisms:**

*Acinetobacter lwoffii*\*

*Haemophilus influenzae*

*Haemophilus parainfluenzae*\*

### **Other microorganisms:**

*Chlamydia trachomatis*

\*Efficacy for this organism was studied in fewer than 10 infections.

The following *invitro* data are also available, **but their clinical significance in ophthalmic infections is unknown**. The safety and effectiveness of moxifloxacin ophthalmic solution in treating ophthalmological infections due to these microorganisms

have not been established in adequate and well-controlled trials.

The following organisms are considered susceptible when evaluated using systemic breakpoints. However, a correlation between the *invitro* systemic breakpoint and ophthalmological efficacy has not been established. The list of organisms is provided as guidance only in assessing the potential treatment of conjunctival infections.

Moxifloxacin exhibits *invitro* minimal inhibitory concentrations (MICs) of 2 microgram/mL or less (systemic susceptible breakpoint) against most (greater than or equal to 90%) strains of the following ocular pathogens.

**Aerobic Gram-positive microorganisms:**

*Listeria monocytogenes*  
*Staphylococcus saprophyticus*  
*Streptococcus agalactiae*  
*Streptococcus mitis*  
*Streptococcus pyogenes*  
*Streptococcus Group C, G and F*

**Aerobic Gram-negative microorganisms:**

*Acinetobacter baumannii*  
*Acinetobacter calcoaceticus*  
*Citrobacter freundii*  
*Citrobacter koseri*  
*Enterobacter aerogenes*  
*Enterobacter cloacae*  
*Escherichia coli*  
*Klebsiella oxytoca*  
*Klebsiella pneumoniae*  
*Moraxella catarrhalis*  
*Morganella morganii*  
*Neisseria gonorrhoeae*  
*Proteus mirabilis*  
*Proteus vulgaris*  
*Pseudomonas stutzeri*

**Anaerobic microorganisms:**

*Clostridium perfringens*  
*Fusobacterium species*  
*Prevotella species*  
*Propionibacterium acnes*

**Other microorganisms:**

*Chlamydia pneumoniae*  
*Legionella pneumophila*  
*Mycobacterium avium*  
*Mycobacterium marinum*  
*Mycoplasma pneumoniae*

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis and Mutagenesis and Impairment of Fertility**

### Carcinogenesis

Long-term studies in animals to determine the carcinogenic potential of moxifloxacin have not been performed. However, in an accelerated study with initiators and promoters, moxifloxacin was not carcinogenic in rats following up to 38 weeks of oral dosing at 500 mg/kg/day (3224 times the highest recommended total daily human ophthalmic dose for a 60 kg person, based on body surface area).

### Mutagenesis

Moxifloxacin was not mutagenic in four bacterial strains used in the Ames Salmonella reversion assay. As with other quinolones, the positive response observed with moxifloxacin in strain TA 102 using the same assay may be due to the inhibition of DNA gyrase. Moxifloxacin was not mutagenic in the CHO/HGPRT mammalian cell gene mutation assay. An equivocal result was obtained in the same assay when v79 cells were used. Moxifloxacin was clastogenic in the v79 chromosome aberration assay, but it did not induce unscheduled DNA synthesis in cultured rat hepatocytes. There was no evidence of genotoxicity *in vivo* in a micronucleus test or a dominant lethal test in mice.

### Impairment of Fertility

Moxifloxacin had no effect on fertility in male and female rats at oral doses as high as 500 mg/kg/day, approximately 3224 times the highest recommended total daily human ophthalmic dose, based on body surface area. At 500 mg/kg/day orally there were slight effects on sperm morphology (head-tail separation) in male rats and on the estrous cycle in female rats.

## **14 CLINICAL STUDIES**

In two randomized, double-masked, multicenter, controlled clinical trials in which patients were dosed 3 times a day for 4 days, moxifloxacin ophthalmic solution produced clinical cures on day 5-6 in 66% to 69% of patients treated for bacterial conjunctivitis.

Microbiological success rates for the eradication of baseline pathogens ranged from 84% to 94%.

In a randomized, double-masked, multicenter, parallel-group clinical trial of pediatric patients with bacterial conjunctivitis between birth and 31 days of age, patients were dosed with moxifloxacin ophthalmic solution or another anti-infective agent. Clinical outcomes for the trial demonstrated a clinical cure rate of 80% at Day 9 and a microbiological eradication success rate of 92% at Day 9.

Please note that microbiologic eradication does not always correlate with clinical outcome in anti-infective trials.

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

Product: 50090-4781

NDC: 50090-4781-0 3 mL in a BOTTLE / 1 in a CARTON

## **17 PATIENT COUNSELING INFORMATION**

### Avoid Contamination of the Product

Advise patients not to touch the dropper tip to any surface to avoid contaminating the contents.

### Avoid Contact Lens Wear

Advise patients not to wear contact lenses if they have signs and symptoms of bacterial conjunctivitis [see Warnings and Precautions (5.3)].

### Hypersensitivity Reactions

Systemically administered quinolones including moxifloxacin have been associated with hypersensitivity reactions, even following a single dose. Instruct patients to discontinue use immediately and contact their physician at the first sign of a rash or allergic reaction [see Warnings and Precautions (5.1)].

Rx Only

### **Manufactured for:**

**Alembic Pharmaceuticals, Inc.**

Bedminster, NJ 07921,  
USA


Made in India.

### **Manufactured by:**

**Gland Pharma Limited**

D.P. Pally, Dundigal Post,  
Hyderabad-500 043, India (IND)

### **Moxifloxacin ophthalmic solution**



## **MOXIFLOXACIN OPHTHALMIC SOLUTION**

moxifloxacin ophthalmic solution/ drops

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:50090-4781(NDC:62332-505)
<b>Route of Administration</b>	OPHTHALMIC		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>MOXIFLOXACIN HYDROCHLORIDE</b> (UNII: C53598599T) (MOXIFLOXACIN - UNII:U188XYD42P)	MOXIFLOXACIN	5 mg in 1 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>BORIC ACID</b> (UNII: R57ZH85D4)	
<b>HYDROCHLORIC ACID</b> (UNII: QTT17582CB)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>WATER</b> (UNII: 059QF0KOOR)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50090-4781-0	1 in 1 CARTON	12/05/2019	
1		3 mL in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209469	02/13/2019	

**Labeler** - A-S Medication Solutions (830016429)

## Establishment

Name	Address	ID/FEI	Business Operations
A-S Medication Solutions		830016429	RELABEL(50090-4781)