

KROGER DAYTIME COLD AND FLU SOFTGEL- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

The Kroger Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Kroger DayTime Cold and Flu Softgel

Drug Facts

Active ingredients (in each Softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Purposes

Pain reliever / Fever reducer

Cough suppressant

Nasal decongestant

Uses:

temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

Warnings:

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is

accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • liver disease • heart disease • thyroid disease • diabetes • high blood pressure • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if • you get nervous, dizzy or sleepless • pain, nasal congestion or cough get worse or last more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

Directions:

- take only as directed - see Overdose warning • do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information:

- store at room temperature between 20-25 °C (68-77 °F)

Inactive ingredients:

Edible white ink, FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan solution

Questions or comments:

1-800-632-6900

COMPARE TO the active ingredients of VICKS® DAYQUIL®

COLD & FLU RELIEF LIQUICAPS® *See bottom panel

POWERFUL NON-DROWSY RELIEF

OUR PHARMACIST RECOMMENDED

Multi-Symptom Relief

TAMPER EVIDENT: DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

READ AND KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

**DISTRIBUTED BY
THE KROGER CO.
CINCINNATI, OHIO 45202**

MADE IN UAE

For More Product Information, Scan UPC

Using Your **Kroger App** or Call 800-632-6900

*This product is not manufactured or distributed by The Proctor & Gamble Company. Vicks®, DayQuil®, and LiquiCaps® are registered trademarks of The Proctor & Gamble Company.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Our Brands, Our Guarantee.

Love It or Your Money Back.

www.kroger.com

Packaging



COMPARE TO the active ingredients of VICKS® DAYQUIL® COLD & FLU RELIEF LIQUICAPS® *See bottom panel

NDC 30142-087-48

POWERFUL NON-DROWSY RELIEF

Kroger

Cold & Flu

DayTime

Multi-Symptom Relief

Acetaminophen Dextromethorphan HBr Phenylephrine HCl

OUR PHARMACIST RECOMMENDS

ACTUAL SIZE

Pain Reliever
Fever Reducer
Cough Suppressant
Nasal Decongestant

48 SOFTGELS

Kroger

Cold & Flu

DayTime

Multi-Symptom Relief

Acetaminophen Dextromethorphan HBr Phenylephrine HCl

Pain Reliever
Fever Reducer
Cough Suppressant
Nasal Decongestant

48 SOFTGELS

PARENTS: Learn about them medicine abuse. www.stoolmediclinebus.com

Our Brands. Our Guarantee. Love It or Your Money Back.

TAMPER EVIDENT: DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

Drug Facts

Active ingredients (in each Softgel)

Active ingredients (in each Softgel)	Purposes
Acetaminophen 325 mg	Pain reliever / Fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses: temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat & bronchial irritation
- headache
- minor aches & pains
- sore throat
- fever

Warnings:

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses in 24 hrs, which is the maximum daily amount for this product

with other drugs containing acetaminophen

3 or more alcoholic drinks daily while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have liver disease, heart disease, thyroid disease, diabetes, high blood pressure, trouble urinating due to enlarged prostate gland, cough that occurs with too much phlegm (mucus), persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if you get nervous, dizzy or sleepless, pain, nasal congestion or cough get worse or last more than 7 days, fever gets worse or lasts more than 3 days, redness or swelling is present, new symptoms occur, cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

Other information: store at room temperature between 20-25 °C (68-77 °F)

Inactive ingredients: Edible white ink, FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan solution

Questions or comments: 1-800-632-6900

Drug Facts (continued)

Directions:

adults & children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information:

store at room temperature between 20-25 °C (68-77 °F)

Inactive ingredients: Edible white ink, FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan solution

Questions or comments: 1-800-632-6900

READ AND KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

DRUG FACTS TABLE

Drug Facts	
Active ingredients (in each Softgel)	Purposes
Acetaminophen 325 mg	Pain reliever / Fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant
Uses:	
temporarily relieves common cold/flu symptoms:	
nasal congestion	cough due to minor throat & bronchial irritation
headache	minor aches & pains
	sore throat
	fever
Warnings:	
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses in 24 hrs, which is the maximum daily amount for this product	
with other drugs containing acetaminophen	
3 or more alcoholic drinks daily while using this product	
Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:	
skin reddening	blisters
	rash
If a skin reaction occurs, stop use and seek medical help right away.	
Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
Ask a doctor before use if you have liver disease, heart disease, thyroid disease, diabetes, high blood pressure, trouble urinating due to enlarged prostate gland, cough that occurs with too much phlegm (mucus), persistent or chronic cough as occurs with smoking, asthma, or emphysema	
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.	
When using this product, do not use more than directed.	
Stop use and ask a doctor if you get nervous, dizzy or sleepless, pain, nasal congestion or cough get worse or last more than 7 days, fever gets worse or lasts more than 3 days, redness or swelling is present, new symptoms occur, cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children.	
Overdose warning: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.	

Drug Facts (continued)	
Directions:	
adults & children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use
Other information:	
store at room temperature between 20-25 °C (68-77 °F)	
Inactive ingredients:	
Edible white ink, FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol sorbitan solution	
Questions or comments: 1-800-632-6900	

KROGER DAYTIME COLD AND FLU SOFTGEL

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:30142-087
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	807
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30142-087-48	4 in 1 CARTON	08/31/2022	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/31/2022	

Labeler - The Kroger Company (006999528)

Revised: 7/2023

The Kroger Company