

**FRIDABABY CHILDREN PAIN AND FEVER ACETAMINOPHEN 160MG PER 5ML VIAL 10CT- acetaminophen 160mg per 5ml solution**  
Fridababy LLC

-----  
**Fridababy Children Pain and Fever Acetaminophen 160mg per 5mL vial 10ct**

***Active Ingredient (in each 5mL)***

Acetaminophen, 160mg

***Purpose***

Pain reliever/fever reducer

***Uses***

temporarily:

- reduces fever
- relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

***Warnings***

**Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes ■ more than 5 doses in 24 hours, which is the maximum daily amount for this product

- with other drugs containing acetaminophen.

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** if sore throat is severe, persistent for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting. Consult a doctor promptly.

**Do not use**

with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if your child**

has liver disease

**Ask a doctor or pharmacist before use if your**

child is taking the blood thinning drug warfarin

### **Stop use and ask a doctor if**

■ pain gets worse or lasts more than 5 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

### **Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical even if you do not notice any signs or symptoms.

### **Directions**

■ **this product does not contain directions or complete warnings for adult use**

■ do not give more than directed (see overdose warning)

■ use the chart below to confirm dose. If possible, use weight to dose, otherwise, use age.

■ give the recommended number of vials below. To open, follow the instructions under How to Use.

■ if needed, repeat dose every 4 hours while symptoms last.

■ do not give more than 5 times in 24 hours

■ do not give for more than 5 days unless directed by a doctor

---

<b>Weight (lbs)</b>	<b>Age (Years)</b>	<b>Dose</b>
under 24 lbs.	under 2 years	Ask a doctor
24-35 lbs.	2 - 3 years	1 Single Use Vial (5 mL)
36-47 lbs.	4-5 years	Ask a doctor
48-71 lbs.	6-10 Years	2 Single Use Vial (10 mL)
72-95 lbs	11 years	3 Single Use Vial (15 mL)

---

### **Other Information**

■ Each Single Use Vial contains 8mg of sodium

■ Store at 68-77°F (20-25°C)

■ Tamper Evident: do not use if the pouch is damaged, open or broken. Do not use if any vial is open or broken.

### **Inactive Ingredient**


citric acid, cherry flavor, edetate disodium, glycerin, maltitol, polyethylene glycol,

propylene glycol, purified water, sodium benzoate, sodium metabisulfite, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum.

### Questions or Comments?

1-833-WTF-FRIDA (1-833-983-3743)

### Package



**Back Panel**

**frida baby®**

**Easy dose it.**

These pre-measured vials deliver the right dose in three easy steps, at home or on-the-go

**Pain + Fever**

Reduce fever and get pain relief from cold + flu symptoms

**How To Use**



Tap the vial to get rid of air bubbles



Twist the top off the vial



Squeeze all liquid in vial into your kid's mouth

No parabens, dyes, alcohol or high fructose corn syrup

Love it? Leave a review. [fridababy.com](http://fridababy.com)  
Medicine Questions? [frida.com/help](http://frida.com/help)

Distributed by: Fridababy, LLC  
82 NE 26th St. Miami, FL 33437



8 10028 77312 7

**Drug Facts**

Active Ingredient (in each 5mL)	Purpose
Acetaminophen, 160mg	Pain reliever/fever reducer

**Uses temporarily:** ■ reduces fever ■ relieves minor aches and pains due to:  
■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

**Warnings**

**Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes ■ more than 5 doses in 24 hours, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** if sore throat is severe, persistent for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly

**Do not use** with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if your child** has liver disease.

**Ask a doctor or pharmacist before use if your child is** taking the blood thinning drug warfarin

**Stop use and ask a doctor if** ■ pain gets worse or lasts more than 5 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical even if you do not notice any signs or symptoms.

**Directions** ■ **this product does not contain directions or complete warnings for adult use** ■ do not give more than directed (see overdose warning) ■ use the chart below to confirm dose. If possible, use weight to dose, otherwise use age ■ give the recommended number of vials below. To open, follow the instructions under How To Use. ■ if needed, repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours ■ do not give for more than 5 days unless directed by a doctor

Weight (lbs.)	Age (Years)	Dose
under 24 lbs.	Under 2 years	Ask a doctor
24-35 lbs.	2-3 years	1 Single Use Vial (5mL)
36-47 lbs.	4-5 years	Ask a doctor
48-71 lbs.	6-10 years	2 Single Use Vials (10mL)
72-95 lbs.	11 years	3 Single Use Vials (15mL)

**Other Information** ■ Each single use vial contains 8mg of sodium ■ Store at 68-77°F (20-25°C) ■ Tamper Evident: do not use if pouch is damaged, open or broken. Do not use if any vial is open or broken.

**Inactive Ingredients** citric acid, cherry flavor, edetate disodium, glycerin, maltitol, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium metabisulfite, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum

**Questions or Comments?** 1-833-WTF-FRIDA (1-833-983-3743)

PL-0013v0

NDC: 72705-124-10

**WARNING: Choking Hazard. Small parts. Keep out of reach of children.**

075

**FRIDABABY CHILDREN PAIN AND FEVER ACETAMINOPHEN 160MG PER 5ML VIAL 10CT**

acetaminophen 160mg per 5ml solution

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:72705-124
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

**Inactive Ingredients**

Ingredient Name	Strength
MALTITOL (UNII: D65DG142WK)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	

<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>SODIUM METABISULFITE</b> (UNII: 4VON5FNS3C)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72705-124-10	10 in 1 POUCH	08/16/2022	
1		5 mL in 1 VIAL; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/16/2022	08/31/2024

**Labeler** - Fridababy LLC (783729598)

Revised: 12/2023

Fridababy LLC