

SODIUM BICARBONATE- sodium bicarbonate powder
Pharma Nobis, LLC

Private Label Sodium Bicarbonate

Drug Facts

Active Ingredient

Sodium Bicarbonate, USP

Purpose

Antacid

Use

For relief of heartburn, acid indigestion, and upset stomach associated with these symptoms.

Warnings

NOT FOR INJECTIONS

Except under supervision of a doctor do not administer to children under 6 years of age.

Do not take more than six, 1/2 tsp. per person up to 60 years old, or three 1/2 tsp. per person 60 years or older in a 24 hour period.

Do not use this product if you are on a sodium restricted diet (each 1/2 tsp. contains 30 mEq (0.7 g) Sodium).

Do not use the maximum dose more than 2 weeks.

Ask a doctor or pharmacist before use if

you are currently taking a prescription drug. Antacids may interact with certain prescription drugs.

Keep out of reach of children.

In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

Directions

Adult and children 6 yrs of age and older:

Take 1/2 tsp. in 1/2 g;ass (4 fl oz) of water every 2 hrs. up to maximum dosage or as directed by doctor.

Inactive ingredients

none

Premier Value Label

4 oz



***Warning:** Do not use if Tamper Evident Seal imprinted "Sealed For Your Protection" is broken or missing. This product is sealed with either a shrink band around cap or foil seal under cap.

Drug Facts	
Active ingredient: Bicarbonate of Soda USP.....	Purpose Antacid
Use For relief of heartburn, acid indigestion, and upset stomach associated with these symptoms.	
Warnings NOT FOR INJECTIONS. Except under supervision of a doctor: ■ do not administer to children under 6 years of age ■ do not take more than 6, 1/2 tsps. per person up to 60 years old, or 3, 1/2 tsps. per person 60 years or older in a 24 hour period. ■ do not use this product if you are on a sodium restricted diet (each 1/2 tsp. contains 30mEq. (0.7g) Sodium). ■ do not use the maximum dose for more than 2 weeks.	
Keep out of reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.	
Directions Adults and children 6 yrs. of age and older	
Take 1/2 level tsp. in 1/2 glass (4 fl oz) of water every 2 hrs. up to maximum dosage or as directed by a doctor.	


If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

8 40986 01749 0

RO11510RLG

Dist. by: Chain Drug Consortium, LLC.
2300 NW Corporate Blvd., Suite 115
Boca Raton, FL 33431

16 oz



***Warning:** Do not use if Tamper Evident Seal imprinted "Sealed For Your Protection" is broken or missing. This product is sealed with either a shrink band around cap or foil seal under cap.

Drug Facts	
Active ingredient: Bicarbonate of Soda USP.....	Purpose Antacid
Use For relief of heartburn, acid indigestion, and upset stomach associated with these symptoms.	
Warnings NOT FOR INJECTIONS. Except under supervision of a doctor: <ul style="list-style-type: none"> ■ do not administer to children under 6 years of age ■ do not take more than 6, 1/2 tsp. per person up to 60 years old, or 3, 1/2 tsp. per person 60 years or older in a 24 hour period. ■ do not use this product if you are on a sodium restricted diet (each 1/2 tsp. contains 30mEq. (0.7g) Sodium). ■ do not use the maximum dose for more than 2 weeks. 	
Keep out of reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.	
Directions Adults and children 6 yrs. of age and older	
Take 1/2 level tsp. in 1/2 glass (4 fl oz) of water every 2 hrs. up to maximum dosage or as directed by a doctor.	

If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.
 Distributed by
 Chain Drug Consortium, LLC.
 2300 NW Corporate Blvd., Suite 115
 Boca Raton, FL 33431

8 40986 01751 3
 R011510R1G

Vida Mia Label

*CON SELLO DE SEGURIDAD

*SAFETY SEALED



mi salud

SODIUM BICARBONATE ORAL POWDER USP

BICARBONATO DE SODIO POLVO ORAL USP

ANTACID • ANTIÁCIDO

(Sodium Bicarbonate) USP and Food Grade
(Bicarbonato de Sodio) USP Calidad Alimentaria

CONTENTS 16 OZ (1LB) 454g • CONTENIDO 16 onzas (1 lb) 454g

*WARNINGS: Do not use if Tamper Evident Seal imprinted "Sealed for your protection" is broken or missing. This product is sealed with either a shrink band around cap or foil seal under cap.

*ADVERTENCIAS: No utilizar si el sello impreso de Evidencia de Alteración "Sellado para su Protección" está roto o hace falta. Este producto está sellado con una banda alrededor de la tapa o con un sello de aluminio debajo de la tapa.

Drug Facts

Active ingredient	Purpose
Sodium Bicarbonate USP	Antacid

Use For relief of heartburn, acid indigestion, and upset stomach associated with these symptoms.

Warnings NOT FOR INJECTIONS.

Except under supervision of a doctor:

- Do not administer to children under 6 years of age.
- Do not take more than six, ½ tsp. per person up to 60 years old, or three, ½ tsp. per person 60 years or older in a 24 hour period.
- Do not use this product if you are on a sodium restricted diet (each ½ sp. contains 30mEq, 3.7g Sodium).
- Do not use the maximum dose for more than 2 weeks.
- Ask a doctor or pharmacist before use if you are presently taking a prescription. Antacids may interact with certain prescription drugs.

Keep out of reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

Directions

Adults and children 6 yrs. of age or older	Take ½ level tsp. in ½ glass (4 fl oz) of water every 2 hrs. up to maximum dosage or as directed by doctor.
--	---

Distributed by/Distribuido por:

R070213.RLG

MAGE LAN
DISTRIBUTION SOLUTIONS

Medley, Florida 35178

Questions or Comments? (31-800-662-3435



8 47717 06076 3



3/8" TIEDOWN



mi salud

BICARBONATO DE SODIO POLVO ORAL USP

ANTIÁCIDO

(Bicarbonato de Sodio) USP Calidad Alimentaria

*ADVERTENCIAS: No utilizar si el sello impreso de Evidencia de Alteración "Sellado para su Protección" está roto o hace falta. Este producto está sellado con una banda alrededor de la tapa o con un sello de aluminio debajo de la tapa.

Información sobre el Medicamento

Ingreiente Activo	Propósito
Bicarbonato de Sodio USP	Antiácido

Uso Para el alivio de la acidez estomacal, la indigestión ácida y el malestar estomacal asociado con estos síntomas.

Advertencias NO DEBE SER INYECTADO.

Excepto bajo supervisión de un médico:

- No se debe administrar a niños menores de 6 años de edad.
- En un período de 24 horas, no tome más de 6 dosis de ½ cucharilla por persona si tiene hasta 60 años de edad o 3 dosis de ½ cucharilla por persona si tiene más de 60 años.
- No utilice este producto si usted está en una dieta restringida de sodio (cada ½ cucharilla contiene 30mEq, 3.7g de Sodio).
- No use la dosis máxima durante más de 2 semanas.
- Consulte con un médico o farmacéutico si está tomando algún medicamento de prescripción. Los antiácidos pueden interactuar con algunos medicamentos de prescripción.

Mantener fuera del alcance de los niños. En caso de sobredosis accidental, busque asistencia profesional o contacte un Centro de Control de Envenenamiento inmediatamente.

Instrucciones

Adultos y niños de 6 años de edad o mayores:	Tome ½ cucharadita en medio vaso de agua (4 onzas líquidas) cada 2 horas hasta la dosis máxima o según las indicaciones del médico.
--	---

¿Preguntas o comentarios? (31-800-662-3435

A34135 07 2022



BICARBONATO DE SODA POLVO INGERIBLE U.S.P.

Sodium Bicarbonate Oral Powder U.S.P.
(BICARBONATE OF SODA) U.S.P. AND FOOD GRADE
NOT FOR MAKING INJECTIONS.

USE: Anti-Acid, and for baking.
INDICATIONS: For the relief of Heartburn, Sour Stomach, and/or Acid Indigestion.
DOSE: 0.3 to 2 grams 1 to 4 times daily. Should not be given to young children (to age six), unless prescribed by their Physician.
WARNING: KEEP OUT THE REACH OF CHILDREN. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

CONTENTS 4 OUNCES (113g.)
DIST. BY HUMCO, TEXARKANA, TEXAS 75501

WARNING: Do not use if Tamper Evident Seal imprinted "Sealed For Your Protection" is broken or missing. This product is sealed with either a shrink band around cap or foil seal under cap.
WARNING: Do not take more than 2 grams in a 24 hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a physician.
WARNING: Do not use this product except under the advice of a Physician if you are on a Sodium Restricted Diet. Contains 270 mg. Sodium per Gram. As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product.



USP: Indicado y para usar con agua.
Indicaciones: Para el alivio de Acidez Estomacal, Hiperacidez, Dolor. Dosis: 0.3 a 2 gramos 1 a 4 veces al día. No se debe dar a niños menores de 6 años, excepto por orden médica.
ADVERTENCIA: Manténgase lejos del alcance de los niños. En caso de sobredosis accidental, consulte a un médico o pongase en contacto inmediatamente con un Centro de Control de Intoxicaciones.
ADVERTENCIA: No se recomienda el uso de este producto por más de 2 semanas, excepto bajo la supervisión y vigilancia médica.
ADVERTENCIA: Use este producto solamente bajo indicación médica si usted está en una dieta restringida de sodio. Contiene 270 mg. de sodio por gramo. Como con cualquier medicamento, si usted está embarazada o está amamantando un bebé, consulte a un médico o profesional de la salud antes de usar este producto.

NO USAR PARA INYECCIONES

Bicarbonato de Sodio Polvo Ingerible U.S.P.

SELLO DE SEGURIDAD ADVERTENCIA: No usar si la Banda de Seguridad impresa "Sealed For Your Protection" está rota o falta. Este producto está sellado con una banda adherida de la tapa o con una hoja sellada de metal bajo la tapa.

ROTBORSALES

SODIUM BICARBONATE

sodium bicarbonate powder

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:82645-904
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM BICARBONATE (UNII: 8MDF5V39QO) (SODIUM CATION - UNII:LYR4MONH37)	SODIUM BICARBONATE	1000 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82645-904-01	454 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/27/2017	
2	NDC:82645-904-94	113 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/27/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	10/27/2017	

Labeler - Pharma Nobis, LLC (118564114)

Registrant - Pharma Nobis, LLC (118564114)

Establishment

Name	Address	ID/FEI	Business Operations
Pharma Nobis, LLC		118564114	label(82645-904) , manufacture(82645-904) , pack(82645-904) , analysis(82645-904)

Revised: 12/2023

Pharma Nobis, LLC