

JUNIOR STRENGTH ACETAMINOPHEN- acetaminophen tablet, chewable

Target Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Target 44-449

Active ingredient (in each tablet)

Acetaminophen 160 mg

Purpose

Pain Reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - sore throat
 - flu
 - headache
 - the common cold
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has

liver disease.

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days

- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- dissolve in mouth or chew before swallowing
- if needed, repeat dose every 4 hours while symptoms last
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor

Weight (lb)	Age (yr)	Tablets
Under 48	Under 6	Ask a doctor
48-59	6-8	2
60-71	9-10	2 1/2
72-95	11	3

Other information

- see end flap for expiration date and lot number
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients

citric acid, crospovidone, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrans hydrated, ethylcellulose, FD&C blue #1 aluminum lake, flavors, magnesium stearate, mannitol, polyethylene, stearic acid, sucralose

Questions?

Call 1-800-910-6874

Principal Display Panel

NDC 11673-449-08

Compare to active ingredient in Junior Strength Tylenol® Meltaways®*

**junior strength
acetaminophen**

tablets, 160 mg

pain reliever/fever reducer

rapid tabs melt in mouth

aspirin free, ibuprofen free

GRAPE FLAVOR

AGE 6-11 YEARS

24 TABLETS

up&up[®]

24 TABLETS

*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Junior Strength Tylenol[®] Meltaways[®]. 50844 REV0213F44908

094 01 0603 ID201416

Distributed by Target Corporation

Minneapolis, MN 55403

© 2013 Target Brands, Inc.

All Rights Reserved

Shop Target.com

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS
TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

junior strength acetaminophen
tablets, 160 mg

up&up

Compare to active ingredient in
Junior Strength Tylenol[®] Meltaways[®]

NDC 11673-449-08

junior strength
acetaminophen
tablets, 160 mg

pain reliever/fever reducer
rapid tabs melt in mouth
aspirin free, ibuprofen free

GRAPE
FLAVOR

AGE
6-11
YEARS

24
TABLETS

24 TABLETS

8067777805
5

Questions? Call 1-800-910-6874

see end flap for expiration date and lot number

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS
TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by McNeil
Consumer Healthcare, owner of the registered trademark Junior
Strength Tylenol[®] Meltaways[®]. 50844 REV0213F44908

094 01 0603 ID201416
Distributed by Target Corporation
Minneapolis, MN 55403
© 2013 Target Brands, Inc.
All Rights Reserved
Shop Target.com

B-1920-449-08
REV0213F44908



Drug Facts (continued)																
<ul style="list-style-type: none"> do not give more than 5 times in 24 hours do not give for more than 5 days unless directed by a doctor 	<table border="1"> <tr> <th>Weight (lb)</th> <th>Age (yr)</th> <th>Tablets</th> </tr> <tr> <td>Under 48</td> <td>Under 6</td> <td>Ask a doctor</td> </tr> <tr> <td>48-59</td> <td>6-8</td> <td>2</td> </tr> <tr> <td>60-71</td> <td>9-10</td> <td>2 1/2</td> </tr> <tr> <td>72-95</td> <td>11</td> <td>3</td> </tr> </table>	Weight (lb)	Age (yr)	Tablets	Under 48	Under 6	Ask a doctor	48-59	6-8	2	60-71	9-10	2 1/2	72-95	11	3
Weight (lb)	Age (yr)	Tablets														
Under 48	Under 6	Ask a doctor														
48-59	6-8	2														
60-71	9-10	2 1/2														
72-95	11	3														
Drug Facts (continued)																
<ul style="list-style-type: none"> TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) 	<p>Other Information</p> <p>inactive ingredients: citric acid, croscopolone, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrans hydrate, ethylcellulose, FD&C blue #1 aluminum lake, flavors, magnesium stearate, mannitol, polyethylene, stearic acid</p>															

Drug Facts (continued)	
<p>Warnings</p> <ul style="list-style-type: none"> Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount. with other drugs containing acetaminophen Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. 	<p>Do not use</p> <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if your child is allergic to acetaminophen or any of the inactive ingredients in this product
Drug Facts (continued)	
<p>Uses</p> <ul style="list-style-type: none"> temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> flu headache the common cold temporarily reduces fever 	<p>Active ingredient (in each tablet)</p> <p>Acetaminophen 160 mg. . . . Pain reliever/fever reducer</p>
Drug Facts (continued)	
<p>Directions</p> <ul style="list-style-type: none"> this product does not contain directions or complete warnings for adult use dissolve in mouth or chew before swallowing if needed, repeat dose every 4 hours while symptoms last find the right dose on chart below. If possible, use weight to dose; otherwise, use age. 	<p>Purpose</p> <p>FOR COMPLETE PRODUCT INFORMATION</p>
Drug Facts (continued)	
<p>Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.</p> <p>Keep out of reach of children.</p> <p>These could be signs of a serious condition:</p> <ul style="list-style-type: none"> redness or swelling is present new symptoms occur fever gets worse or lasts more than 3 days gain gets worse or lasts more than 5 days <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> recommended dosage. When using this product do not exceed ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin. ask a doctor before use if your child has liver disease. 	<p>Other Information</p> <p>do not give more than 5 times in 24 hours</p> <p>do not give for more than 5 days unless directed by a doctor</p>

Target 44-449

JUNIOR STRENGTH ACETAMINOPHEN

acetaminophen tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-449
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
D&C RED NO. 27 (UNII: 2LRS185U6K)	
D&C RED NO. 30 (UNII: 2S42T2808B)	
ETHYLCELLULOSES (UNII: 7Z8S9VYZ4B)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

MANNITOL (UNII: 3OWL53L36A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics			
Color	PURPLE	Score	2 pieces
Shape	ROUND	Size	16mm
Flavor	GRAPE	Imprint Code	44;449
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-449-08	4 in 1 CARTON	01/28/2005	08/04/2019
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part343	01/28/2005	08/04/2019

Labeler - Target Corporation (006961700)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(11673-449)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(11673-449)