LEVOCETIRIZINE DIHYDROCHLORIDE- levocetirizine dihydrochloride tablet, coated

KROGER COMPANY

Drug Facts

Active ingredient (in each tablet)

Levocetirizine dihydrochloride USP, 5 mg

Purpose

Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings

Do not use

- if you have kidney disease
- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine

Ask a doctor before use if you have

· ever had trouble urinating or emptying your bladder

When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask doctor if

- you have trouble urinating or emptying your bladder
- an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding:

• if breast-feding: not recommended

• if pregnant: ask a health professional before use

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

adults 65 years of age and older	ask a doctor
adults and children 12-64 years of age	 take 1 tablet (5 mg) once daily in the evening do not take more than 1 tablet (5 mg) in 24 hours 1/2 tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms
children 6-11 years of age	 take 1/2 tablet (2.5 mg) once daily in the evening do not take more than 1/2 tablet (2.5 mg) in 24 hours
children under 6 years of age	• do not use
consumers with kidney disease	• do not use

Other information

- store between 20° and 25°C (68° and 77°F)
- (Bottle only)- Safety Sealed: Do not use if carton was opened or if printed foil inner seal on bottle is torn or missing
- (Blister only)- Satey Sealed: Do not use if seal is broken or if individual blister unit is open or torn

Inactive ingredients

colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

Questions or comments?

Call 1-888-375-3784

Carton Label

Carton Label



Bottle Label

Bottle Label



Active ingredient (in each tablet) Purpose
Levocetirizine dihydrochloride USP, 5 mg.......Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

Warnings Do not use ■ if you have kidney disease ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine Ask a doctor before use if you have ■ ever had trouble urinating or emptying your bladder When using this product ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery Stop use and ask a doctor if ■ you have trouble urinating or

(Confinued On Back of Label)
DISTRIBUTED BY
THE KROGER CO.
CINCINNATI, OHIO 45202
MADE IN INDIA
QUALITY GUARANTEE
800-632-6900 | www.kroger.com
150090908 REV 10/22





emptying your bladder ■ an allergic reaction to this product occurs. Seek medical help right away. If pregnant or breast-feeding: ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions adults 65 years of age and older ■ ask a doctor adults and children 12-64 years of age ■ take 1 tablet (5 mg) once daily in the evening ■ do not take more than 1 tablet (5 mg) in 24 hours ■ 1/2 tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms children 6-11 years of age ■ take 1/2 tablet (2.5 mg) once daily in ▶

the evening ■ do not take more than 1/2 tablet (2.5 mg) in 24 hours **children under 6 years of age** ■ do not use **consumers** with kidney disease ■ do not use

Other information ■ store between 20° and 25°C (68° and 77°F) ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing

Inactive ingredients colloidal silicon dioxide, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

Questions and comments? call 1-800-632-6900

LEVOCETIRIZINE DIHYDROCHLORIDE

levocetirizine dihydrochloride tablet, coated

D	1
Product	Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:30142-717(NDC:43598-735)

Route of Administration ORAL

Active Ingredient/Active Moiety Ingredient Name Basis of Strength Ievocetirizine dihydrochloride (UNII: SOD6A38AGA) (levocetirizine UNII:6U5EA9RT2O) Basis of Strength Ievocetirizine dihydrochloride 5 mg

Inactive Ingredients	
Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

Product Characteristics					
Color	white	Score	2 pieces		
Shape	OVAL	Size	9mm		
Flavor		Imprint Code	L		
Contains					

P	Packaging						
#	Item Code	Package Description	Marketing Start Date	Marketing End Date			
1	NDC:30142-717- 35	1 in 1 CARTON	09/30/2018				
1		35 in 1 BOTTLE; Type 0: Not a Combination Product					
2	NDC:30142-717- 16	2 in 1 CARTON	09/30/2018				
2	NDC:30142-717- 80	80 in 1 BOTTLE; Type 0: Not a Combination Product					

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA210375	09/30/2018		

Labeler - KROGER COMPANY (006999528)

Revised: 10/2023 KROGER COMPANY