TOPCARE STOMACH RELIEF ORIGINAL STRENGTH- bismuth subsalicylate tablet TopCo Associates, LLC

Drug Facts

Active ingredient

Bismuth subsalicylate 262 mg (each tablet contains 102 mg salicylate)

Purpose

Antidiarrheal/Antacid

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink including:
- heartburn
- indigestion
- nausea
- gas
- belching
- fullness

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer
- a bleeding problem

bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if

you are taking any drug for

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

If pregnant or breast-feeding

ask a health professional before use.

Keep out of reach of children

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

• chew or dissolve in mouth

Adults and children 12 years and over: 2 tablets every ½ to 1 hour as needed

- do not exceed 16 tablets in 24 hours
- use until diarrhea stops but not more than 2 days

Children under 12 years of age: ask a doctor

• drink plenty of clear fluids to help prevent dehydration caused by diarrhea


Other information

- **Phenylketonurics:** contains phenylalanine 1.1mg per tablet
- Calcium content per tablet: 73mg
- save carton for full directions and warnings
- store at room temperature 15°- 30°C (59°- 86°F)
- do not use if cellophane unit is torn

Inactive ingredients

acacia gum, aspartame, calcium carbonate, D&C red #27 aluminum lake, dextrates, magnesium stearate, maltodextrin, microcrystalline cellulose, peppermint flavor, silicon dioxide

Package/Label Principal Display Panel



TOPCARE STOMACH RELIEF ORIGINAL STRENGTH bismuth subsalicylate tablet					
Product Information					
Product Type	HUMAN OTC DRUG	Item Code (Sou	rce)	NDC:7616	2-001
Route of Administration	ORAL				
Active Ingredient/Active Moiety					
Ingr	edient Name		Basis of S	trength	Strength

Inactive Ingredients			
Ingredient Name	Strength		
ACACIA (UNII: 5C5403N26O)			
ASPARTAME (UNII: Z0H242BBR1)			
CALCIUM CARBONATE (UNII: H0G9379FGK)			
D&C RED NO. 27 (UNII: 2LRS185U6K)			
DEXTRATES (UNII: G263MI44RU)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
MALTODEXTRIN (UNII: 7CVR7L4A2D)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
PEPPERMINT (UNII: V95R5KMY2B)			

Product Characteristics				
Color	pink	Score	no score	
Shape	ROUND	Size	17mm	
Flavor		Imprint Code	RH046	
Contains				

Packaging					
# Item Code	Package Description	Marketing Start Date	Marketing End Date		
NDC:76162-001- 85	6 in 1 CARTON	12/23/2014			
1	5 in 1 BLISTER PACK; Type 0: Not a Combination Product				

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC Monograph Drug	M008	12/23/2014			

Labeler - TopCo Associates, LLC (006935977)

Revised: 12/2018 TopCo Associates, LLC