

NICOTINE POLACRILEX COATED ICE MINT- nicotine polacrilex gum, chewing
CVS Pharmacy

Nicotine Polacrilex Coated Ice Mint 2 mg and 4 mg CVS

Drug Facts

Active ingredient (in each chewing piece) - 2 mg

Nicotine Polacrilex 2 mg (nicotine)

Active ingredient (in each chewing piece) - 4 mg

Nicotine Polacrilex 4 mg (nicotine)

Purpose

Stop smoking aid

Use

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions - 2 mg

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day

- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Directions - 4 mg

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information - 2 mg

- **each piece contains:** calcium 115 mg, sodium 8 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

Other information - 4 mg

- **each piece contains:** calcium 120 mg, sodium 8 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

Inactive ingredients - 2 mg

acacia, acesulfame potassium, carnauba wax, flavors, gum base, hydroxypropyl

cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, talc, titanium dioxide and xylitol.

Inactive ingredients - 4 mg

acacia, acesulfame potassium, carnauba wax, D&C yellow #10 aluminum lake, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, talc, titanium dioxide and xylitol.

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal display panel 2 mg

Compare to the active ingredient in Nicorette® Gum*

Sugar Free

Nicotine Gum

Nicotine Polacrilex Gum USP, 2 mg (nicotine)

STOP SMOKING AID

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 minutes of waking up, use Nicotine Polacrilex Gum USP, 4 mg

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

TO INCREASE YOUR SUCCESS IN QUITTING.

1. You must be motivated to quit.
2. Use Enough - Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
3. Use Long Enough - Use nicotine Polacrilex Gum for the full 12 weeks.

Use with a support program as directed in the enclosed User's Guide.

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum.

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

Distributed by: CVS Pharmacy, Inc.

One CVS Drive, Woonsocket, RI 02895

Package label 2 mg

Drug Facts (continued)

Other information

- each piece contains: calcium 115 mg, sodium 8 mg
- store at 20-25°C (68-77°F)
- protect from light

Inactive ingredients

acacia, aceulfame potassium, carnauba wax, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, talc, titanium dioxide, xylitol

Questions or comments?
Call 1-877-753-3935
Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum.

THEFT SURVEILLANCE TAG AREA

Distributed by:
CVS Pharmacy, Inc.
One CVS Drive
Woonsocket, RI 02895
© 2021 CVS/pharmacy
CVS.com®
1-800-SHOP CVS
V-16402

CVS Quality
Money Back Guarantee
PLD-D616A FC008095

minute clinic
START TO STOP
CAN HELP YOU QUIT

CVS.com/StartToStop

Visit MinuteClinic® for personal counseling with a licensed provider.

CVS Health.

Compare to the active ingredient in Nicorette® Gum*

NDC 69842-216-15

Sugar Free

Nicotine Gum

2 mg

Nicotine Polacrilex Gum USP, 2 mg (nicotine)

STOP SMOKING AID

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette **WITHIN 30 MINUTES** of waking up, use Nicotine Polacrilex Gum USP, 4 mg.

Actual Size

Coated Ice Mint Flavor

20 PIECES, 2 mg EACH

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

Push gum through foil.

Peel off backing starting at corner with blunt edge.

To remove the gum, tear off single unit.

TO INCREASE YOUR SUCCESS IN QUITTING:

- You must be motivated to quit.
- Use Enough – Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
- Use Long Enough – Use Nicotine Polacrilex Gum for the full 12 weeks.
- Use with a support program as directed in the enclosed User's Guide.

#978270

0 50428 53503 5

CVS HEALTH Sugar Free Nicotine Gum Coated Ice Mint Flavor

Principal display panel 4 mg

Compare to the active ingredient in Nicorette® Gum
Sugar Free
Nicotine Gum

Nicotine Polacrilex Gum USP, 4 mg (nicotine)

STOP SMOKING AID

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.

If you smoke your first cigarette MORE THAN 30 minutes after waking up, use Nicotine Polacrilex Gum USP 2 mg

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use Enough - Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
3. Use Long Enough - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. Use with a support program as directed in the enclosed User's Guide.

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum.

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

Distributed by: CVS Pharmacy, Inc.

One CVS Drive, Woonsocket, RI 02895

Package label 4 mg

minute clinic

START TO STOP[®]
CAN HELP YOU QUIT

CVS.com/StartToStop

Visit MinuteClinic[®] for
personal counseling with
a licensed provider.

CVS
Health.

Compare to the
active ingredient in
Nicorette[®] Gum*

NDC 69842-217-15

Sugar Free Nicotine Gum

Nicotine Polacrilex
Gum USP, 4 mg
(nicotine)

**4
mg**

STOP SMOKING AID

**FOR THOSE WHO SMOKE
THEIR FIRST CIGARETTE WITHIN
30 MINUTES OF WAKING UP.**

If you smoke your first
cigarette **MORE THAN
30 MINUTES** after
waking up, use
Nicotine Polacrilex
Gum USP, 2 mg.

 Actual
Size
**Coated Ice Mint Flavor
20 PIECES, 4 mg EACH**

Drug Facts (continued)

Other information

- each piece contains:
calcium 120 mg, sodium 8 mg
- store at 20-25°C (68-77°F)
- protect from light

Inactive ingredients

acacia, acesulfame potassium,
carnauba wax, D&C yellow #10
aluminum lake, flavors, gum base,
hydroxypropyl cellulose,
magnesium oxide, sodium
bicarbonate, sodium carbonate,
talc, titanium dioxide, xylitol

Questions or comments?

Call 1-877-753-3935
Monday-Friday 9AM-5PM EST

*This product is not
manufactured or
distributed by
GlaxoSmithKline
Consumer Healthcare,
L.P., distributor of
Nicorette[®] Gum.

THEFT SURVEILLANCE
TAG AREA

Distributed by:
CVS Pharmacy, Inc.
One CVS Drive
Woonsocket, RI 02895
© 2021 CVS/pharmacy
CVS.com[®]
1-800-SHOP CVS
V-16402

 **CVS Quality**
Money Back Guarantee

PLD-D622A FC08106

**BLISTER PACKAGED FOR YOUR PROTECTION. DO
NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.**

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where
proof of age cannot be verified



- TO INCREASE YOUR SUCCESS IN QUITTING:**
1. You must be motivated to quit.
 2. Use Enough – Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
 3. Use Long Enough – Use Nicotine Polacrilex Gum for the full 12 weeks.
 4. Use with a support program as directed in the enclosed User's Guide.

Drug Facts

Active ingredient (in each chewing piece) Purpose
Nicotine polacrilex..... 4 mg (nicotine)..... Stop smoking aid

Use ■ reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have
■ a sodium-restricted diet ■ heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate. ■ high blood pressure not controlled with medication. Nicotine can increase blood pressure.
■ stomach ulcer or diabetes ■ history of seizures

Ask a doctor or pharmacist before use if you are
■ using a non-nicotine stop smoking drug ■ taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if
■ mouth, teeth, or jaw problems occur ■ irregular heartbeat or palpitations occur
■ you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
■ you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
■ if you are under 18 years of age, ask a doctor before use
■ before using this product, read the enclosed User's Guide for complete directions and other important information ■ begin using the gum on your quit day
■ if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum ■ if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece ■ to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks ■ if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

#979096



CVS HEALTH Sugar Free Nicotine Gum Coated Ice Mint Flavor

NICOTINE POLACRILEX COATED ICE MINT

nicotine polacrilex gum, chewing

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69842-216
Route of Administration	BUCCAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
MAGNESIUM OXIDE (UNII: 3A3U0GI71G)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM CARBONATE (UNII: 45P3261C7T)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
XYLITOL (UNII: VCQ006KQ1E)	

Product Characteristics

Color	white	Score	no score
Shape	SQUARE	Size	14mm
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-216-44	100 in 1 CARTON	05/01/2016	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:69842-216-15	20 in 1 CARTON	05/01/2016	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:69842-216-16	160 in 1 CARTON	05/01/2016	
3		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA204794	05/01/2016	

NICOTINE POLACRILEX COATED ICE MINT

nicotine polacrilex gum gum, chewing

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69842-217
Route of Administration	BUCCAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	4 mg

Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
MAGNESIUM OXIDE (UNII: 3A3U0GI71G)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM CARBONATE (UNII: 45P3261C7T)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
XYLITOL (UNII: VCQ006KQ1E)	

Product Characteristics

Color	yellow	Score	no score
Shape	SQUARE	Size	14mm
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-217-44	100 in 1 CARTON	05/01/2016	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:69842-217-15	20 in 1 CARTON	05/01/2016	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:69842-217-16	160 in 1 CARTON	05/01/2016	
3		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA204833	05/01/2016	

Labeler - CVS Pharmacy (062312574)

Revised: 2/2023

CVS Pharmacy