

MEDIQUE BACK PAIN RELIEF- magnesium salicylate tetrahydrate tablet, film coated
Unifirst First Aid Corporation

Medique Back Pain Relief

Drug Facts

Active ingredient (in each caplet)

Magnesium Salicylate Tetrahydrate (NSAID*) 580 mg
(equivalent to 467.2 mg of anhydrous magnesium salicylate)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

For the temporary relief of minor backache pain

Warnings

Reye's syndrome:

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:

Magnesium salicylate may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcohol drinks every day while using this product

■ take more or for a longer time than directed

Do not use if you are allergic to salicylates (including aspirin) or any other pain reliever/fever reducer.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
- if ringing in the ears or loss of hearing occurs
- new symptoms occur
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use magnesium salicylate at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose

Adults and children: (12 years and over)

Take 2 caplets every 6 hours, not to exceed 8 caplets in 24 hours.

Children under 12 years:

Do not use unless directed by a doctor

Other information

- each caplet contains: magnesium 40 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- tamper evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, stearic acid, titanium dioxide, triacetin

Questions or comments? 1-800-634-7680

Medique®

Back Pain Relief


Pain Reliever/ Anti-Inflammatory

Pull to Open

Magnesium Salicylate Tetrahydrate (NSAID) 580 mg

80 Caplets (40 x 2)

Tamper Evident Unit Dose Packets



MEDIQUE BACK PAIN RELIEF

magnesium salicylate tetrahydrate tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-074
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
MAGNESIUM SALICYLATE (UNII: 41728CY7UX) (SALICYLIC ACID - UNII:O414PZ4LPZ)		MAGNESIUM SALICYLATE	580 mg	
Inactive Ingredients				
Ingredient Name			Strength	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
POLYDEXTROSE (UNII: VH2XOU12IE)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
TRIACETIN (UNII: XHX3C3X673)				
Product Characteristics				
Color	white	Score	no score	
Shape	CAPSULE (Caplet)	Size	17mm	
Flavor		Imprint Code	44;338	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-074-80	40 in 1 BOX	09/26/2022	
1	NDC:47682-074-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-074-36	75 in 1 BOX	12/12/2022	
2		2 in 1 PACKET; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013		09/26/2022	

Labeler - Unifirst First Aid Corporation (832947092)