

**IDO DISINFECTION ETHANOL- ethanol liquid**  
**IDO PHARM**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

ethanol

Water

First aid to help prevent the risk of infection in minor cuts, scrapes or burns.

Keep out of reach of children

For the external use only

For external use only

- If taken internally, serious gastric disturbances will result
- flammable, keep away from heat or flame

Ask a doctor before use if you have deep or puncture wounds, animal bites or serious burns

When using this product

- do not use in or near eyes. In case of contact with eyes, rinse eyes thoroughly with water
- do not apply over large areas of the body
- do not use longer than one week unless directed by a doctor

Stop use and ask a doctor if irritation or rash appears

Keep out of reach of children. In case of ingestion, get medical help or contact a Poison Control Center 1-800-222-1222 or dial 911 immediately

Clean affected area

Apply small amount of this product on the affected area 1-3 times daily

May be covered with a sterile bandage. If bandaged, let dry first

children under 6 years of age should be supervised when using this product

# IDO DISINFECTION ETHANOL

<b>Drug Facts</b>	
<b>Active Ingredient</b>	<b>Purpose</b>
ethyl Alcohol 83% w/w	Antiseptic
<b>Uses</b> Disinfection of hands and skin, disinfection of surgical site skin	
<b>Warnings</b> <ul style="list-style-type: none"><li>■ For external use only</li><li>■ Flammable, keep away from fire or flames</li></ul>	
<b>Do not Use</b> In the oral cavity, mucous membranes and damaged skin (It has an irritating effect)	
<b>When using this product</b> <ul style="list-style-type: none"><li>■ Be careful not to get it into your eyes.</li><li>■ In case of extensive or long-term use, be careful not to inhale the vapor. (If ethanol vapor is consumed in large amounts or repeatedly, irritation to the mucous membrane, headache, etc. may appear.)</li><li>■ In case of repeated use on the same site, be careful as the skin may become rough due to degreasing, etc.</li><li>■ Since organic substances such as serum and pus weaken the sterilization effect of this product, when using this product to disinfect medical equipment, etc., use this product after thoroughly washing the equipment</li><li>■ Be careful with fire as it is flammable and explosive</li><li>■ It is desirable to avoid using synthetic rubber products, synthetic resin products, optical instruments, glass instruments, painted catheters, etc.</li><li>■ If it is necessary to soak metal instruments for a long time, add 0.2~1% sodium nitrite to prevent corrosion</li></ul>	
<b>Stop use</b> <ul style="list-style-type: none"><li>■ As hypersensitivity symptoms such as rash may appear, in this case, discontinue use</li><li>■ As skin irritation symptoms may appear, so in this case, discontinue use</li></ul>	
<b>Directions</b> <ul style="list-style-type: none"><li>■ Apply to the area to be disinfected</li><li>■ Supervise children under 6 years of age when using this product to avoid swallowing</li></ul>	
<b>Other Information</b> <ul style="list-style-type: none"><li>■ Store in a light-shielding airtight container, avoiding fire</li><li>■ Avoid freezing and excessive heat above 40C (104F)</li></ul>	
<b>Inactive Ingredients</b> Water	
<b>Questions or Comments?</b> Call 82-31-4936545	

**[Manufacturer]** IDO PHARM / 41, Beonyeong-ro 15beon-gil, Danwon-gu, Ansan, Gyeonggido 15616, Korea, South (KOR)

## IDO DISINFECTION ETHANOL

ethanol liquid

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:77039-022
<b>Route of Administration</b>	TOPICAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
ALCOHOL (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M)	ALCOHOL	83 mL in 100 mL

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
WATER (UNII: 059QF0KO0R)	

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:77039-022-01	500 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/01/2022	

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph not final	part333A	05/01/2022	

**Labeler** - IDO PHARM (694853523)

**Registrant** - IDO PHARM (694853523)

## Establishment

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
IDO PHARM		694853523	manufacture(77039-022)

Revised: 8/2022

IDO PHARM