

RAPIDOL ASPIRIN DISPLAY 2X25- aspirin tablet, coated
Pharmadel LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Rapidol Aspirin Display (HPUSA)

Drug Facts

Active ingredient and Purpose

Active ingredient

(in each tablet)

Aspirin 325 mg **(NSAID)***.....

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

For the temporary relief of minor aches and pains due to:

- a cold
- a headache
- a toothache
- muscular aches
- a backache
- minor pains of arthritis

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- shock
- facial swelling
- asthma (wheezing)

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an **NSAID**, which may cause severe stomach bleeding. The chance is higher if you:

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take more or for a longer time than directed
- are age 60 or older
- have 3 or more alcoholic drinks everyday while using this product

Do not use

if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have not been drinking fluids
- you have lost a lot of fluid due to vomiting or diarrhea
- you are taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

- pain persist, gets worse or last for more than 10 days
- ringing in the ears or a loss of hearing occurs, consult a doctor before taking any more of this product
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

It is very important not to use aspirin during the last 3 months of pregnancy unless directed to do so by a doctor, because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- **adults and children 12 years of age and over:**
- take two (2) tablet every 4 hours
- do not take more than 6 doses in 24 hours
- **children under 12 years of age:** consult a doctor

Other information

- store between 68-77°F (20-25°C) in a dry place
- use by expiration date on package

TAMPER EVIDENT: Do not use if individual packets are broken or torn.

Inactive ingredients

corn starch, hypromellose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments?

1-866-359-3478 (M-F) 9AM to 5 PM EST or www.pharmadel.com

Principal Display Panel

NDC 55758-312-50

NDC 55758-312-02

Rapidol

Aspirin/Aspirina

325 mg

Alivia Dolor (AINE*) Pain Reliever (NSAID*)

Relieves / Alivia

- ✓ Headaches/ Dolor de Cabeza
- ✓ Muscular Pain/ Dolores Musculares
- ✓ Minor Arthritis Pain/ Dolores de Artritis Leves

SIN CAFEINA / CAFFEINE FREE

25 Sobres x 2 Tabletas 25 packets of 2 tablets each

2 Coated Tablets

Rapido[®] ASPIRIN

Pain Reliever (NSAID)* 325 mg

Drug Facts

Active ingredient (in each tablet)
Aspirin 325 mg (NSAID)* **Pain reliever**

*nonsteroidal anti-inflammatory drug
Uses

For the temporary relief of minor aches and pains due to: ■ a cold ■ a headache ■ a toothache ■ muscular aches ■ a backache ■ minor pains of arthritis

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reactions. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ take more or for a longer time than directed ■ are age 60 or older ■ have 3 or more alcoholic drinks everyday while using this product **Do not use** if you are allergic to aspirin or any other pain reliever/fever reducer. **Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

Drug Facts (continued)

■ you have not been drinking fluids ■ you have lost a lot of fluid due to vomiting or diarrhea ■ you are taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis
Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding:

- feel faint ■ vomit blood ■ have bloody or black stools
- have stomach pain that does not get better

■ pain persist, gets worse or last for more than 10 days
■ ringing in the ears or a loss of hearing occurs, consult a doctor before taking any more of this product ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is very important not to use aspirin during the last 3 months of pregnancy unless directed to do so by a doctor, because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away

Directions

- do not take more than directed ■ drink a full glass of water with each dose
- adults and children 12 years of age and over:
 - take two (2) tablet every 4 hours
 - do not take more than 6 doses in 24 hours
- children under 12 years of age: consult a doctor

Other information

■ store between 68-77°F (20-25°C) in a dry place

Inactive ingredients

corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

Questions or comments? 1-866-359-3478 (M-F) 9AM to 5 PM EST or www.pharmadel.com

Distributed by:
PHARMADEL LLC
Georgetown, DE 19947

**DO NOT USE IF
POUCH IS BROKEN
OR TORN**

Alivia Dolor (AINE)
325 mg
ASPIRINA
Rapido[®]

Drug Facts

Active ingredient (in each tablet)
Aspirin 325 mg (NSAID)* **Pain reliever**

*nonsteroidal anti-inflammatory drug
Uses

For the temporary relief of minor aches and pains due to: ■ a cold ■ a headache ■ a toothache ■ muscular aches ■ a backache ■ minor pains of arthritis

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reactions. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ take more or for a longer time than directed ■ are age 60 or older ■ have 3 or more alcoholic drinks everyday while using this product **Do not use** if you are allergic to aspirin or any other pain reliever/fever reducer. **Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

Drug Facts (continued)
■ you have not been drinking fluids ■ you have lost a lot of fluid due to vomiting or diarrhea ■ you are taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis
Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding:

- feel faint ■ vomit blood ■ have bloody or black stools
- have stomach pain that does not get better

■ pain persist, gets worse or last for more than 10 days
■ ringing in the ears or a loss of hearing occurs, consult a doctor before taking any more of this product ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is very important not to use aspirin during the last 3 months of pregnancy unless directed to do so by a doctor, because it may cause problems in the unborn child or complications during delivery.

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

Questions or comments? 1-866-359-3478 (M-F) 9AM to 5 PM EST or www.pharmadel.com

Distributed by / Distribuido por:
PHARMADEL LLC
Georgetown, DE 19947



Información de los medicamentos

Ingredientes activos (por cada tableta)
Aspirina 325 mg (AINE)* **Analgésico**

*antiinflamatorio no esteroideo
Usos

Para el alivio temporal de dolores menores y molestias debidas a: ■ un resaca ■ un dolor de cabeza ■ un dolor de muelas ■ dolores musculares ■ un dolor de espalda ■ un dolor de espalda ■ un dolor de espalda

Advertencias

Síndrome de Reye: Niños y adolescentes que hayan o estén recuperándose de sarampión o de la gripe no deben usar este producto. Cuando se use este producto, si se observan cambios en el comportamiento con náuseas y vómitos, consulte a un médico porque estos síntomas podrían ser una señal temprana del síndrome de Reye, una enfermedad rara pero grave.

Alerta de alergia: La aspirina puede causar una reacción alérgica grave, que puede incluir: ■ hives ■ inflamación facial ■ asma (sibilancias) ■ shock
Si ocurre una reacción alérgica, deje de usar el producto y busque ayuda médica de inmediato.

Advertencia de sangrado estomacal: Este producto contiene un AINE, que puede causar sangrado estomacal grave. El riesgo es mayor si usted: ■ tiene antecedentes de úlceras o sangrado estomacal ■ toma otros medicamentos que contengan AINEs (aspirina, ibuprofeno, naproxeno, u otros) ■ toma más o por un tiempo más largo del indicado ■ tiene 60 años o más ■ tiene 3 o más bebidas alcohólicas por día

Si está embarazada o lactando, consulte a su proveedor de la salud antes de usar este medicamento. Es muy importante no usar aspirina durante los últimos 3 meses de embarazo a menos que lo indique un médico, ya que puede causar problemas en el niño o complicaciones durante el parto.

Mantenga fuera del alcance de los niños. En caso de sobredosis, obtenga ayuda médica o llame al Centro de Control de Venenos.

Alivio / Alivia
✓ Headaches / Dolor de Cabeza
✓ Muscular Pain / Dolores Musculares
✓ Minor Arthritis Pain / Dolores de Artritis Leves

Ingredientes Inactivos
maiz almidón, croscarmellose sodio, hipromelosa, microcristalina celulosa, aceite mineral, dióxido de titanio

¿Preguntas o comentarios? 1-866-359-3478 (M-F) 9AM to 5 PM EST or www.pharmadel.com

Información de los medicamentos

Ingredientes activos (por cada tableta)
Aspirina 325 mg (AINE)* **Analgésico**

*antiinflamatorio no esteroideo
Usos

Para el alivio temporal de dolores menores y molestias debidas a: ■ un resaca ■ un dolor de cabeza ■ un dolor de muelas ■ dolores musculares ■ un dolor de espalda ■ un dolor de espalda

Advertencias

Síndrome de Reye: Niños y adolescentes que hayan o estén recuperándose de sarampión o de la gripe no deben usar este producto. Cuando se use este producto, si se observan cambios en el comportamiento con náuseas y vómitos, consulte a un médico porque estos síntomas podrían ser una señal temprana del síndrome de Reye, una enfermedad rara pero grave.

Alerta de alergia: La aspirina puede causar una reacción alérgica grave, que puede incluir: ■ hives ■ inflamación facial ■ asma (sibilancias) ■ shock
Si ocurre una reacción alérgica, deje de usar el producto y busque ayuda médica de inmediato.

Advertencia de sangrado estomacal: Este producto contiene un AINE, que puede causar sangrado estomacal grave. El riesgo es mayor si usted: ■ tiene antecedentes de úlceras o sangrado estomacal ■ toma otros medicamentos que contengan AINEs (aspirina, ibuprofeno, naproxeno, u otros) ■ toma más o por un tiempo más largo del indicado ■ tiene 60 años o más ■ tiene 3 o más bebidas alcohólicas por día

Si está embarazada o lactando, consulte a su proveedor de la salud antes de usar este medicamento. Es muy importante no usar aspirina durante los últimos 3 meses de embarazo a menos que lo indique un médico, ya que puede causar problemas en el niño o complicaciones durante el parto.

Mantenga fuera del alcance de los niños. En caso de sobredosis, obtenga ayuda médica o llame al Centro de Control de Venenos.

Alivio / Alivia
✓ Headaches / Dolor de Cabeza
✓ Muscular Pain / Dolores Musculares
✓ Minor Arthritis Pain / Dolores de Artritis Leves

Ingredientes Inactivos
maiz almidón, croscarmellose sodio, hipromelosa, microcristalina celulosa, aceite mineral, dióxido de titanio

¿Preguntas o comentarios? 1-866-359-3478 (M-F) 9AM to 5 PM EST or www.pharmadel.com

Distributed by / Distribuido por:
PHARMADEL LLC
Georgetown, DE 19947



NDC 55758-212-50

ASPIRINA
325 mg

Alivia Dolor (AINE)
Pain Reliever (NSAID)



Relieves / Alivia

- ✓ Headaches / Dolor de Cabeza
- ✓ Muscular Pain / Dolores Musculares
- ✓ Minor Arthritis Pain / Dolores de Artritis Leves

SIN CAFEINA / CAFFEINE FREE

25 Sobres x 2 Tabletas each
25 packets of 2 tablets each

ASPIRINA
325 mg
Pain Reliever (NSAID)

RAPIDOL ASPIRIN DISPLAY 2X25

aspirin tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55758-362
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POVIDONE (UNII: FZ989GH94E)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	white	Score	score with uneven pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	ASP325
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55758-362-50	25 in 1 CARTON	08/10/2022	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:55758-362-02	2 in 1 PACKET; Type 0: Not a Combination Product	08/10/2022	



Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	08/10/2022	

Labeler - Pharmadel LLC (030129680)

Revised: 8/2022

Pharmadel LLC