EXTRA STRENGTH PAIN RELIEVER- acetaminophen 500 mg, film-coated caplet tablet, film coated Lil' Drug Store Products, Inc.

Pain Reliever, Circle K

Drug Facts

Active ingredient

Active ingredient(in each caplet)

Acetaminophen USP 500 mg

Purpose

Purpose

Pain reliever/fever reducer

Uses

Uses

- temporarily relieves minor aches and pains due to:
- the common cold
- headache
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

skin reddening

- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick

medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Directions

do not take more than directed (see Overdose warning)

adults and children 12 years and over

- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

children under 12 years

ask a doctor

Other information

Other information

store between 20° to 25°C (68° to 77°F)

Inactive ingredients

Inactive ingredients povidone, pregelatinized starch (maize), stearic acid

Questions

Questions or comments?

Call toll-free 1-877-507-6516 (M-F 8AM-4:30PM CST)

**This product is not manufactured or distributed by McNeil Consumer Healthcare Division, distributor of Tylenol® Extra Strength Caplets.

PDP/Package

TEMPORARILY RELIEVES MINOR ACHES & PAINS

Compare to the active ingredient in

Tylenol® Extra Strength Caplets**

[Circle K logo]

extra strength pain reliever 50 caplets Acetaminophen 500 mg Pain Reliever/Fever Reducer

[caplet image]

actual size

50 caplets



TAMPER EVIDENT: DO NOT USE IF CARTON IS OPEN OR IF KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts

Active inaredient (in each caplet)

Purpose

Acetaminophen USP 500 mg...

Pain reliever. fever reduce

- temporarily relieves minor aches and pains
- the common cold
 - headache
 - backache minor pain of arthritis toothache

 - muscular aches premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
skin reddening blisters rash

Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present.
 These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Geep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any

EXTRA STRENGTH PAIN RELIEVER

acetaminophen 500 mg, film-coated caplet tablet, film coated

ORAL

Product Information

Route of Administration

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-5680

Active Ingredient/Active Moiety

POVIDONE K30 (UNII: U725QWY32X)

Ingredient Name	Basis of Strength	Strength
ingredient Name	Basis of Strength	Strengt

ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII: 36209ITL9D) **ACETAMINOPHEN** 500 mg

Inactive Ingredients Ingredient Name Strength STARCH, CORN (UNII: O8232NY3SJ) PALMITOSTEARIC ACID (UNII: Q8Y7S3B85M)

Product Characteristics			
Color	white (white to off-white)	Score	no score
Shape	OVAL (capsule-shaped biconvex)	Size	18mm
Flavor		Imprint Code	N79
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715- 5680-4	1 in 1 CARTON	05/09/2022	
1		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/09/2022	

Labeler - Lil' Drug Store Products, Inc. (093103646)

Revised: 12/2023 Lil' Drug Store Products, Inc.