

**MUCUS RELIEF COLD FLU AND SORE THROAT MAXIMUM STRENGTH-  
acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid  
United Natural Foods, Inc. dba UNFI**

-----  
**Drug Facts**

**Active ingredients (in each 20 mL)**

**Acetaminophen 650 mg**

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCl 10 mg

**Purposes**

**Pain reliever/fever reducer**

Cough suppressant

Expectorant

Nasal decongestant

**Uses**

- temporarily relieves these common cold and flu symptoms:
  - sinus congestion and pressure
  - minor aches and pain
  - nasal congestion
  - cough due to minor throat and bronchial irritation
  - sore throat
  - headache
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

**When using this product,**

**do not use more than directed.**

**Stop use and ask a doctor if**

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed (see overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL every 4 hours while symptoms last
- children under 12 years of age: do not use

### **Other information**

- **each 20 mL contains:** sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

### **Inactive ingredients**

anhydrous citric acid, disodium EDTA,, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

### **Questions or comments?**

Call toll free **1-877-932-7948**

### **Principal Display Panel**

Compare to Maximum Strength Mucinex® Fast-Max® Cold, Flu & Sore Throat active ingredients\*

Maximum Strength

Mucus Relief

Cold, flu & sore throat

acetaminophen 650 mg (pain reliever-fever reducer)

dextromethorphan HBr 20 mg (cough suppressant)

guaifenesin 400 mg (expectorant)

phenylephrine HCl 10 mg (nasal decongestant)

- relieves headache & fever
- controls cough
- relieves nasal & chest congestion
- thins & loosens mucus

for ages 12 & over

FL OZ (mL)

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Cold, Flu & Sore Throat.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

Distributed by Supervalu Inc.

Eden Prairie, MN 55344 USA

[www.supervaluprivatebrands.com](http://www.supervaluprivatebrands.com)

**Package Label**

NDC 41163-035-06

compare to  
**Maximum Strength  
Mucinex® Fast-Max®  
Cold, Flu & Sore Throat**  
active ingredients\*

**EQUALINE®**

maximum strength  
**mucus relief**  
cold, flu & sore throat

**acetaminophen 650 mg**  
(pain reliever- fever reducer)  
**dextromethorphan HBr 20 mg**  
(cough suppressant)  
**guaifenesin 400 mg** (expectorant)  
**phenylephrine HCl 10 mg** (nasal decongestant)

- relieves headache & fever
- controls cough
- relieves nasal & chest congestion
- thins & loosens mucus

for ages 12 & over

PLD-D282C  
LB002648

6 FL OZ (177 mL)

DISTRIBUTED BY SUPERVALU INC., EDEN PRAIRIE, MN 55344 USA

**100% Quality  
GUARANTEED**

Like it or let us make it right.  
**That's our quality promise.**  
877-932-7948  
supervaluprivatebrands.com



**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY  
SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Cold, Flu & Sore Throat.

PLD-C282C LB002647

**PARENTS:**  
Learn about how medicine works  
[www.StopMedicineFromKids.org](http://www.StopMedicineFromKids.org)

| Drug Facts                            |                             |
|---------------------------------------|-----------------------------|
| Active ingredients<br>(in each 20 mL) | Purposes                    |
| Acetaminophen 650 mg.....             | Pain reliever/fever reducer |
| Dextromethorphan HBr 20 mg.....       | Cough suppressant           |
| Guaifenesin 400 mg.....               | Expectorant                 |
| Phenylephrine HCl 10 mg.....          | Nasal decongestant          |

PEEL CORNER FOR MORE DRUG FACTS

**Drug Facts (continued)**

**Uses**

- temporarily relieves these common cold and flu symptoms
  - sinus congestion and pressure
  - minor aches and pains
  - nasal congestion
  - cough due to minor throat and bronchial irritation
  - sore throat
  - headache
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:
- more than 4,000 mg of acetaminophen in 24 hours
  - with other drugs containing acetaminophen
  - 3 or more alcoholic drinks daily while using this product
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
  - blisters
  - rash
- If a skin reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Drug Facts (continued)**

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin.

**When using this product, do not use more than directed.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL every 4 hours while symptoms last
- children under 12 years of age: do not use

**Other information**

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

**Questions or comments?**

Call toll free 1-877-932-7948

**EQUALINE Mucus Relief Cold, Flu & Sore Throat**

**MUCUS RELIEF COLD FLU AND SORE THROAT MAXIMUM STRENGTH**

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid

**Product Information**

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:41163-035 |
| <b>Route of Administration</b> | ORAL           |                           |               |

**Active Ingredient/Active Moiety**

| <b>Ingredient Name</b>  | <b>Basis of Strength</b>         | <b>Strength</b>    |
|---|----------------------------------|--------------------|
| <b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH)<br>(DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN<br>HYDROBROMIDE | 20 mg<br>in 20 mL  |
| <b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)                       | ACETAMINOPHEN                    | 650 mg<br>in 20 mL |
| <b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)                           | GUAIFENESIN                      | 400 mg<br>in 20 mL |
| <b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)         | PHENYLEPHRINE<br>HYDROCHLORIDE   | 10 mg<br>in 20 mL  |

**Inactive Ingredients**

| <b>Ingredient Name</b>                                | <b>Strength</b> |
|---|-----------------|
| <b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)       |                 |
| <b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)            |                 |
| <b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)         |                 |
| <b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)         |                 |
| <b>GLYCERIN</b> (UNII: PDC6A3C0OX)                    |                 |
| <b>PROPYL GALLATE</b> (UNII: 8D4SNN7V92)              |                 |
| <b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)            |                 |
| <b>WATER</b> (UNII: 059QF0KO0R)                       |                 |
| <b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)             |                 |
| <b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K) |                 |
| <b>SORBITOL</b> (UNII: 506T60A25R)                    |                 |
| <b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)                   |                 |
| <b>XANTHAN GUM</b> (UNII: TTV12P4NEE)                 |                 |

**Packaging**

| <b>#</b> | <b>Item Code</b> | <b>Package Description</b>                                     | <b>Marketing Start Date</b> | <b>Marketing End Date</b> |
|----------|------------------|--|-----------------------------|---------------------------|
| 1        | NDC:41163-035-06 | 177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 12/31/2014                  |                           |

**Marketing Information**

| <b>Marketing Category</b> | <b>Application Number or Monograph Citation</b> | <b>Marketing Start Date</b> | <b>Marketing End Date</b> |
|---------------------------|---|-----------------------------|---------------------------|
| OTC Monograph Drug        | M012  | 12/31/2014                  |                           |

**Labeler** - United Natural Foods, Inc. dba UNFI (943556183)

